UWS Academic Portal

Consistent microplastic ingestion by deep-sea invertebrates over the last four decades (1976-2015), a study from the North East Atlantic

Courtene-Jones, Winnie; Quinn, Brian; Ewins, Ciaran; Gary, Stefan F.; Narayanaswamy, Bhavani E.

Published in:
Environmental Pollution

DOI:
[10.1016/j.envpol.2018.10.090](https://doi.org/10.1016/j.envpol.2018.10.090)

Published: 31/01/2019

Document Version
Peer reviewed version

[Link to publication on the UWS Academic Portal](#)

Citation for published version (APA):

Courtene-Jones, W., Quinn, B., Ewins, C., Gary, S. F., & Narayanaswamy, B. E. (2019). Consistent microplastic ingestion by deep-sea invertebrates over the last four decades (1976-2015), a study from the North East Atlantic. *Environmental Pollution*, 244, 503-512. <https://doi.org/10.1016/j.envpol.2018.10.090>

General rights

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact pure@uws.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Consistent microplastic ingestion by deep-sea invertebrates over the last four decades (1976 – 2015), a study from the North East Atlantic

Winnie Courtene-Jones^{1*}, Brian Quinn², Ciaran Ewins², Stefan F. Gary¹, Bhavani E. Narayanaswamy¹

¹ Scottish Association for Marine Science, Scottish Marine Institute, Oban, Argyll, PA37 1QA, Scotland

² Institute of Biomedical and Environmental Health Research (IBEHR), School of Science & Sport, University of the West of Scotland, Paisley, PA1 2BE, Scotland.

Corresponding Author

* E-mail: winnie.courtene-jones@sams.ac.uk

Graphical abstract:

Abstract

Although evidence suggests the ubiquity of microplastics in the marine environment, our knowledge of its occurrence within remote habitats, such as the deep sea, is scarce. Furthermore, long term investigations of microplastic abundances are even more limited. Here we present a long-term study of the ingestion of microplastics by two deep-sea benthic invertebrates (*Ophiomusium lymani* and *Hymenaster pellucidus*) sampled over four decades. Specimens were collected between the years 1976 – 2015 from a repeat monitoring site > 2000 m deep in the Rockall Trough, North East Atlantic. Microplastics were identified at a relatively consistent level throughout and therefore may have been present at this locality prior to 1976. Considering the mass production of plastics began in the 1940s - 50s our data suggest the relatively rapid occurrence of microplastics within the deep sea. Of the individuals examined (n = 153), 45% had ingested microplastics, of which fibres were most prevalent (95%). A total of eight different polymer types were isolated; polyamide and polyester were found in the highest concentrations and in the majority of years, while low-density polystyrene was only identified in 2015. This study provides an assessment of the historic occurrence of microplastics on the deep seafloor and presents a detailed quantification and characterisation of microplastics ingested by benthic species. Furthermore these data advance our knowledge on the long-term fate of microplastic in marine systems.

Keywords

Microplastic, time series, deep sea, marine litter, long-term study

Capsule

This unique dataset reveals consistent levels of microplastics were ingested by deep-sea invertebrates since 1976, indicating the long-term occurrence of microplastic pollution in this region.

Introduction

Since the 1940s - 50s there has been a rapid and world-wide increase in plastic production. The latest data suggest that around 60% of all plastics ever produced are accumulating in landfills or in the environment (Geyer et al., 2017). By their very nature, synthetic plastics are extremely durable (Shah et al., 2008); but under the influence of environmental exposure, including mechanical forces and/or photochemical processes, plastics fragment into microplastics and nanoplastics (Gewert et al., 2015; ter Halle et al., 2016). Several trillion tonnes of plastics are estimated to be afloat in the world's oceans, with microplastics constituting around 92% of this (Eriksen et al., 2014; van Sebille et al., 2015). Plastic is thus a problem of growing environmental concern and has recently been proposed as a planetary boundary threat. This is defined as having planetary-scale occurrence which is not readily reversible and may cause currently unrecognised disruptive effects to vital Earth system processes (Galloway et al., 2017; Jahnke et al., 2017; Rockström et al., 2009).

With the continued and increasing input of plastic to the oceans (Geyer et al., 2017) it might be expected that surface concentrations would also increase over time, however data reveal inconsistencies. The North Pacific Subtropical Gyre has been documented to be accumulating plastics (Goldstein et al., 2012; Lebreton et al., 2018) while no such trends have been observed in other surface waters (Beer et al., 2017; Law et al., 2014, 2010; ter Halle et al., 2016). Global ocean budgeting identifies large discrepancies between expected and observed surface quantities (Cozar et al., 2014; Eriksen et al., 2014). A number of hypotheses may explain these findings. Firstly, the majority of studies employ surface-towed nets, typically with a mesh size of ~300 µm (Barrows et al., 2017; Hidalgo-ruz et al., 2012). Due to the continual fragmentation of plastics in the environment it is possible that a large proportion of small microplastics are not being detected, or have now fragmented to nano-scale which may be missed by these types of sampling. Secondly, wind-driven mixing of the surface layers may vertically redistribute microplastics, resulting in potentially large amounts going unmeasured (Kukulka et al., 2012). Lastly it is thought that the majority of plastic litter does not remain in surface water but sinks until ultimately residing on the seafloor.

Recently Koelmans et al., (2017), undertook a whole ocean mass balance study; their simulations showed that 99.8% of the plastics which had entered the oceans since 1950 had settled below the ocean surface layer and around 8.5 million tonnes of plastics settle annually. The sinking of plastic particles is related to their size, density, physical/chemical properties (Kowalski et al., 2016) and biological interactions. Biofilm formation occurs rapidly in the environment (Lobelle and Cunliffe, 2011; Rummel et al., 2017), altering particle density and hydrophobicity and in turn increasing the rate of sinking. Microplastics may form aggregates with phytoplankton (Long et al., 2015) or marine snow (Porter et al., 2018; Zhao et al., 2016), flocculate and adhere to transparent exopolymers (Engel, 2004; Passow et al., 2001; Summers et al., 2018) or be incorporated into faecal pellets (Cole et al., 2016; Katija et al., 2017). In most cases these processes act to reduce the buoyancy of microplastics and thus increase the sinking rate, redistributing microplastics from surface waters to the ocean floor.

Data quantifying microplastic contamination in the deep sea are sparse; yet those which have, report microplastics in water (Courtenne-Jones et al., 2017a; Kanhai et al., 2018), benthic invertebrates (Courtenne-Jones et al., 2017a; Taylor et al., 2016) and sediments (Fischer et al., 2015; Van Cauwenberghe et al., 2013; Woodall et al., 2014). Indeed some of the highest concentrations of microplastics reported to date are from deep sediments (42–6595 microplastics kg⁻¹) (Bergmann et al., 2017) supporting the hypothesis that the seafloor may accumulate microplastics. There are some long-term studies which monitor large marine litter on the seafloor (Bergmann and Klages, 2012; Chiba et al., 2018; Galgani et al., 2013, 2010; Maes et al., 2018; Schlining et al., 2013), however we are not aware of any such studies for microplastics. These enquiries are needed to elucidate fundamental questions regarding the behaviour and fate of microplastics within marine systems.

The Scottish Association for Marine Science (SAMS) has conducted deep-sea research at a permanent location 2200 m deep in the Rockall Trough since the mid-1970s. Operations ceased during the period 1995 - 2013 due to a lack of funding; however this collection, spanning four decades represents one of the longest time-series of deep ocean specimens. Organisms were originally collected largely for the analysis of community structure and were preserved intact, giving a unique opportunity to study temporal trends in ingested microplastics. The present study assesses microplastic ingestion over four decades by two species of benthic echinoderms; *Ophiomusium lymani* and *Hymenaster pellucidus*, to examine long-term trends of microplastic abundance and polymer types. *O. lymani* is a benthic deposit feeder and facultative predator (Iken et al., 2001; Pearson and Gage, 1984) while *H. pellucidus* predaes on invertebrates and planktonic fallout; both species are mobile and feed at the surface of the sediment. Additionally, both species have previously been reported to ingest microplastics (Courtenne-Jones et al., 2017a) making them suitable for inclusion in this study.

The aims of this research are (i) to determine whether the first occurrence of microplastic pollution can be detected in the Rockall Trough, and (ii) to examine long-term temporal trends in overall microplastic abundance and constituent polymer types. Utilising this unique archival collection we aim to provide insights regarding the long-term fate of marine microplastics and their historical occurrence on the deep seafloor.

Method

Sample locations and species

'Gage Station M' is situated at a depth of 2200 m in the Rockall Trough (57.300°N, 10.383°W), North East Atlantic Ocean (Figure 1). Deep-sea sampling began at this location in 1976 and has been conducted on an annual basis since, except during an 18 year period between the years 1995 to 2013 where no deep-sea operations were undertaken. The methods to collect samples have remained constant during four decades of operation. The primary historical sampling method entails use of a Woods Hole Oceanographic Institution-pattern epibenthic sled, rigged with polyamide (nylon) nets of mesh size 0.5 mm to sample both the epifauna and infauna. Early sampling (1970s) was also conducted with an Agassiz trawl, which is similar to the epibenthic sled (and also utilise a polyamide net), however it is more suitable for the collection of benthic macro/megafauna (Jamieson et al., 2013). Historic samples were all processed in the same way, involving the contents of the nets being washed through stacked stainless steel sieves (mesh sizes 0.42 mm, 0.5 mm and 1 mm) to separate the invertebrates

into different size classes. Each size fraction was transferred to separate lidded buckets and were fixed using 4% buffered formaldehyde before being stored in 70% ethanol. Upon return to the laboratory faunal assemblages were identified to species levels and stored thereafter in 70% ethanol (Gage et al., 1980) in sealed containers.

Macrobenthic species assemblages in this region are dominated by echinoderms and in particular *Ophiomusium lymani* and *Hymenaster pellucidus* are found in relatively high abundance (Gage, 1986). Both species have previously been identified to ingest microplastics (Courtenne-Jones et al., 2017a) thus for these reasons *O. lymani* (n = 90) and *H. pellucidus* (n = 63) were selected to investigate microplastic loads (SI Figure 1). Specimens from the years 1976, 1980, 1985, 1990, 1995, 2013 and 2015 where possible were analysed, with sample sizes varying slightly between years based on the number of individuals available (details summarised in SI Table 1).

Figure 1. Map showing the sampling locations around the long-term monitoring site 'Gage Station M' (green triangle) to the east of the Anton Dohrn seamount (A.D.S.) in the Rockall Trough; West of the United Kingdom. The area shown in the hatched box is depicted in the subsequent panel, and shows the sampling locations of *O. lymani* (black points), *H. pellucidus* (white points) and both

species (grey points). Bathymetry is contoured at 500 m intervals from depths of 500 m to 3500 m (MATLAB R2015b using GEBCO_2014 bathymetry data).

Quality assurance /quality control

Well documented laboratory quality assurance /quality control (QA/QC) procedures were followed as detailed in Courtene-Jones et al., (2017a) and summarised in the supplementary information. All scientific equipment (scalpels, forceps, glass beakers, filter manifold etc.) were covered/wrapped in aluminium foil and examined under the microscope prior to use. All samples remained covered when not in use to reduce risk of contamination from aerial sources.

At the time of field sampling the same rigorous contamination measures implemented today were not enforced. The same equipment (metal epibenthic sled/Agassiz trawl with a polyamide net (white in colour) see SI Figure 2) has been used throughout the four decades of deep-sea operations. The net was sampled and analysed spectroscopically as well as examined closely under the microscope. While we cannot completely rule out historic contamination at the time of collection, thorough washing of each specimen with a flow of deionised water followed by close visual examination prior to dissection was undertaken to remove any external microplastics which may confound results. As only microplastics internalised within the organisms' soft tissues were assessed the likelihood of contamination arising from sampling are reduced.

Enzymatic digestion of invertebrates

Specimens from each sampling year were removed from ethanol and the width of the central disc was measured using metal dial calipers. Individuals were washed thoroughly with a flow of deionised water and placed into clean lidded glass petri dishes. The exoskeleton was carefully examined under a dissecting microscope to ensure there were no microplastics present, and any found were removed with forceps. Next, the central disc was dissected and all soft tissue was removed from the exoskeleton; in the case of *H. pellucius*, soft tissue extending into each of the five arms (part of the digestive and reproductive systems) was also removed. The soft tissue was then weighed using a Sartorius analytic electronic balance (4 d.p.) before being placed into separate 100 ml glass beakers which were covered with aluminium foil. Enzymatic digestions with the proteolytic digestive enzyme trypsin were carried out following the protocol outlined by Courtene-Jones et al., (2017b).

The 52 µm mesh gauze, through which the solution was filtered, was systematically and thoroughly examined under a dissecting microscope three times and any potential microplastics were removed and placed into a small petri dish containing a 30 mm diameter filter paper (Whatman Grade 1). Once all potential microplastics had been transferred, the 30 mm petri dish was sealed and labelled for later spectroscopic analysis.

Microplastics identification

A Perkin-Elmer One Fourier Transformation Infrared (FTIR) microscope in transmission mode was used to identify potential microplastic particles. Samples were placed onto gold coated glass slides and infrared radiation in the wavelengths 600 - 4000 cm⁻¹ was used. Each spectra produced was the average from 16 scans, a variable aperture size was used and the spectral resolution was 4cm⁻¹. Background scans were taken between each sample and were automatically subtracted from the sample's spectrum. Spectra were visualised in OMNIC 9 (Thermo Fisher Scientific Inc.) with use of the inbuilt Hummel polymer library to aid identification. Additional user generated libraries from the

collaborative University of the West of Scotland and Scottish Association for Marine Science 'UWS/SAMS' library and the Alfred Wegener Institute 'AWI' (Primpke et al., 2018) were utilised to enable the comparison with a wider variety of polymers as well as those having undergone environmental weathering. As well as using these three libraries (Hummel, 'UWS/SAMS' and 'AWI'), the characteristic functional group signals from each spectra acquired were examined manually to confirm the identity of the materials being assessed.

Data analyses

The percentage incidence of ingestion was calculated as the number of individuals that ingested one or more microplastic particle / total number of dissected individuals x 100. This was calculated for each species and when species were grouped for each sampling year. To take into account differences in specimen weights, microplastic ingestion was standardised per gram of wet weight (w.w.) tissue; these data were tested for normal variance structure using a Shapiro and Levene's test. Data were subsequently log₁₀ transformed to meet the criteria of normal distribution and homoscedasticity.

Interspecific differences between the numbers of ingested microplastics/g tissue and the abundance of polymer types ingested were examined by two sample t-test. Polymer diversity was computed based on Shannon-Weiner (H') diversity index (log base transformed data) and species specific and grouped diversity indices were used for Spearman's rank correlation test with year. Differences in the abundance of each polymer type ingested/g tissue between years were compared using the Kruskal-Wallis test (as data failed to meet normal variance structure) and where a significant difference was found a post-hoc Dunn's test was carried out. As the polymer types polyvinylchloride-copolymer and polystyrene were only present during one sampling year these polymers were not included. The polymer composition and abundance between years was also compared by hierarchical cluster analysis based on group average linkage of Bray-Curtis similarity of microplastic/g polymer data. All statistical analysis were performed in R Studio V 0.99.892 (R Core Team, 2016) with the use of the packages *car* (John et al., 2017) *vegan* (Oksanen et al., 2018) and *dunn.test* (Dinno, 2017).

Results

Polymer composition

Eight different polymers were identified (SI Figure 3); overall, polyamide and polyester were the most abundant (Figure 2). Three of the polymer types were ingested by both species, namely polyamide, polyester and acrylic. Alkyd and polyethylene were only isolated from *O. lymani* and PVC-copolymer, polystyrene and polyacrylonitrile were only present in *H. pellucidus*. The mean abundance of microplastics/g of each polymer type did not differ significantly between the two species ($t = -1.096$, $df = 18.72$, $p = 0.287$).

Polymer diversity, measured using the Shannon-Weiner index (Table 1), varied between years and species. When species were grouped, the years 1980 followed by 1985 had the greatest polymer diversity ($H' 1.367$ and 1.225 respectively) and 1990 had the lowest polymer diversity ($H' 0.534$). No correlation was found between polymer diversity and year (*H. pellucidus*: $r_s = -0.700$, $p = 0.233$; *O. lymani*: $r_s = 0.321$, $p = 0.498$; species grouped: $r_s = -0.107$, $p = 0.840$). Cluster analysis based on group average linkage of Bray-Curtis dissimilarity further illustrates the lack of trend between polymer type and abundance between years (SI Figure 4).

Figure 2. Total abundance of different polymers and particle types ingested by *O. lymani* and *H. pellucidus* across all years. The quantities of fibres are indicated by the solid filled bars and fragments by the patterned bars.

PAN = polyacrylonitrile, PA = polyamide, PE = polyethylene, PS = polystyrene, co-PVC = polyvinylchloride-copolymer

Interspecific and inter-annual microplastic ingestion

Microplastic ingestion varied between species with *O. lymani* having a higher incidence of ingestion than *H. pellucidus* (51% and 22% across all years respectively, Table 1). Examining each year separately revealed the highest ingestion occurred in 1995 for *O. lymani* (80%) and in 1985 for *H. pellucidus* (40%) and when species were grouped (45%).

Table 1. Sampling information along with the abundance of microplastics per gram of wet weight soft tissue (MP/g w.w.), incidence of microplastic ingestion, polymer diversity (Shannon-Weiner H') and total number of unique polymer types ingested by each species and when species were grouped from the different sampling years.

Values presented after $\pm$ symbol represents standard deviation. n/a indicates the absence of individuals within the sampling year.

Year	Species	No. individuals sampled	Mean tissue w.w. (g)	Incidence of ingestion	Mean MP/g w.w. tissue	No. unique polymers	Polymer diversity (H')
------	---------	-------------------------	----------------------	------------------------	-----------------------	---------------------	------------------------

2015	<i>O. lymani</i>	20	0.547 ± 0.195	30%	2.57 ± 1.53	3	1.053
	<i>H. pellucidus</i>	20	0.773 ± 0.520	20%	1.27 ± 0.73	2	0.462
	Species grouped	40	0.660 ± 0.404	25%	2.11 ± 1.42	4	1.115
2013	<i>O. lymani</i>	20	0.449 ± 0.168	25%	2.93 ± 1.34	2	0.692
	<i>H. pellucidus</i>	20	0.228 ± 0.173	10%	9.10 ± 4.21	2	0.597
	Species grouped	40	0.338 ± 0.202	18%	4.61 ± 3.62	2	0.660
1995	<i>O. lymani</i>	10	0.585 ± 0.139	80%	2.27 ± 0.97	4	1.207
	<i>H. pellucidus</i>	3	1.863 ± 0.211	33%	0.48 ± 0	1	0.000
	Species grouped	13	0.880 ± 0.579	70%	2.11 ± 1.07	4	1.197
1990	<i>O. lymani</i>	10	0.476 ± 0.157	30%	2.48 ± 0.50	2	0.584
	<i>H. pellucidus</i>	n/a	n/a	n/a	n/a	n/a	n/a
	Species grouped	10	0.476 ± 0.157	30%	2.48 ± 0.50	2	0.584
1985	<i>O. lymani</i>	10	0.627 ± 0.130	50%	2.46 ± 1.23	3	0.670
	<i>H. pellucidus</i>	10	0.838 ± 0.231	40%	2.05 ± 1.05	3	1.278
	Species grouped	20	0.732 ± 0.212	45%	2.55 ± 1.59	4	1.225
1980	<i>O. lymani</i>	10	0.346 ± 0.069	50%	3.43 ± 1.35	3	1.063
	<i>H. pellucidus</i>	10	1.039 ± 0.455	30%	1.36 ± 0.27	4	1.371
	Species grouped	20	0.693 ± 0.476	40%	2.51 ± 1.46	5	1.367
1976	<i>O. lymani</i>	10	0.484 ± 0.142	40%	1.96 ± 0.66	2	0.640
	<i>H. pellucidus</i>	n/a	n/a	n/a	n/a	n/a	n/a
	Species grouped	10	0.484 ± 0.142	40%	1.96 ± 0.66	2	0.640

Microplastics were identified in all years and remained at relatively consistent levels throughout the time series. The abundance of ingested microplastics was between 1.96 ± 0.66 - 4.61 ± 3.62 microplastics/g w.w. (mean $\pm$ SD) when species were grouped; and between 1.96 ± 0.66 - 3.43 ± 1.35 microplastics/g w.w. (mean $\pm$ SD) for *O. lymani* and 0.48 ± 0 - 9.10 ± 4.21 microplastics/g w.w. (mean $\pm$ SD) for *H. pellucidus* (Figure 3). Total abundance of ingested microplastics did not statistically differ between years for either *H. pellucidus* ($F(1, 14) = 0.195$, $p = 0.666$), *O. lymani* ($F(1, 38) = 0.047$, $p = 0.83$), or when species were grouped ($F(1, 54) = 0.011$, $p = 0.918$). Also, there were no significant differences in the abundance of ingested polymers types/g w.w. tissue between years (Figure 4).

Figure 3 Mean abundance of microplastics/g of wet weight tissue across the time period for *O. lymani* (black line), *H. pellucidus* (orange line) and when species are grouped (blue line). Error bars show standard deviation. This plot is a summary of values in presented in the 'Mean MP/g w.w. tissue' column of Table 1.

Figure 4. Composition and mean abundance of each polymer type ingested per gram of wet weight soft tissue across years. Error bars show standard deviation and are absent for those polymers where only a single microplastic was present within the sampling year.

PAN = polyacrylonitrile, PA = polyamide, PE = polyethylene, PS = polystyrene, co-PVC = polyvinylchloride-copolymer

Characterisation of microplastics

FTIR was performed on 470 potential microplastic items; of which 74 were identified as synthetic polymers, 294 items were natural (cellulose (n = 292)/calcium carbonate (n = 2)) and 102 items yielded low quality and unusable spectral data. Thus, 16% of items analysed were confirmed to be microplastics. The majority of plastic items were fibres (n = 70, 95%) with only a very low number of fragments identified (n = 4, 5%). A variety of colours were detected (SI Figure 5) and example microplastics are shown in SI Figure 6. Fragments ranged in diameter from a minimum of 0.08 mm to a maximum of 0.67 mm (mean $\pm$ SD: 0.34 mm $\pm$ 0.26 mm) and fibres ranged in length from 0.21 mm – 4.90 mm (mean $\pm$ SD: 1.20 mm $\pm$ 0.98 mm). All microplastics were less than 5 mm and the size distributions varied only marginally between years (SI Figure 6).

QA/QC

No particles were found on any of the 22 atmospheric controls performed. The quantities of microplastics on Tape Lift Screening (n = 22), used to assess background contamination levels, varied; there was 2.25 fibres $\pm$ 1.58 (mean $\pm$ S.D.) after cleaning and prior to commencing laboratory work, and 3.50 fibres $\pm$ 2.27 (mean $\pm$ S.D.) after laboratory work. These fibres were red, blue and white in colour and identified as cellulose/cotton in their composition by FTIR analysis. Additionally none of the ingested fibres were found to match the characteristics (white polyamide) of the Agassiz/epibenthic sled netting.

Discussion

A consistent abundance of microplastics were identified throughout the entire specimen series spanning a period of four decades. The present dataset represents a quantification of historic microplastic ingestion and develops our knowledge of the occurrence of this pollutant on the deep seafloor. While the number of positively identified microplastics from each sampling year was relatively low, no trends were observed between the overall abundance or the polymer types ingested between years, demonstrating not only the consistent levels of ingested microplastics but that this pollutant likely arrived at the study site prior to 1976.

Global production and use of synthetic plastics have increased exponentially since their mass production in the 1950s (Geyer et al., 2017; PlasticsEurope, 2017), resulting in an increase in the flux of plastic waste entering the environment by as much as 120% between the years 1975 - 2010 (Jambeck et al., 2015). These decadal trends have been observed in some surface waters (Goldstein et al., 2012; Lebreton et al., 2018; Thompson et al., 2004) and in coastal sediments around Belgium between the years 1993 – 2008 (Claessens et al., 2011). It is not well established how long it will take for macroplastics to break down into microplastics and subsequently escape surface waters (GESAMP, 2015) but high concentrations of microplastics in marine sediments indicate this as a potential sink for these small particles (Gago et al., 2018; Woodall et al., 2014). Given the uncertainty surrounding microplastic deposition rates, it was initially hypothesised that this study may observe the first occurrence of microplastics on the Rockall Trough seafloor and identify temporal trends in abundance and/or polymer types using benthic invertebrates as a proxy.

It was realised that the occurrence of ‘the plastic age’ (Thompson et al., 2009) within the Rockall Trough pre-dates our earliest sample, dating from 1976; sampling was not undertaken prior to this preventing the analysis of earlier samples. To some degree this is not entirely surprising. Larger plastics have been documented on the seafloor (100 – 200 m depth) as early as the mid-1970s (Holmström, 1975), and recent experimental and modelling data indicate the relatively rapid transit of microplastics through the water column (Long et al., 2015; Zhang, 2017). Theoretical modelling by Kooi et al., (2017) showed that microplastics with densities greater than seawater (i.e negatively buoyant), such as polyamide, polyester and PVC start to settle immediately and always sink to the seafloor. For example, a 0.1 mm PVC particle may only take 10 days to sink to a depth of 4000 m (Kooi et al., 2017). By this rationale most of the plastics entering the oceans since the 1950s may now reside on the seafloor (Koelmans et al., 2017).

The majority of the polymers identified in the present study (polyester, polyamide, acrylic, alkyd, PAN and PVC) had densities greater than seawater (Andrady, 2011; Crawford and Quinn, 2017), corroborating the findings of other benthic research (Bergmann et al., 2017; Courtene-Jones et al., 2017a; Woodall et al., 2014) and indicating these polymers likely sink directly and relatively rapidly to the seafloor, where they are bioavailable to the benthic community. Positively buoyant microplastics were also isolated; non-expanded polystyrene (1.04 g/cm³ (Crawford and Quinn, 2017)), was only present in 2015, and polyethylene which was one of the lowest density polymers identified in this study (density 0.92 - 0.97 g/cm³ (Crawford and Quinn, 2017)) was identified post-1980. This may indicate that low density polymers sink over longer time scales than their high density counterparts. Experimental work showed polystyrene particles decreased their sinking velocity with increasing salinity (Kowalski et al., 2016). In natural marine waters the density of seawater typically increases with increasing depth; in the Rockall Trough seawater potential density (σ_θ) changes from around

1.0272 g/cm³ in the surface layer to 1.0278 g/cm³ at 2500 m depth (Holliday et al., 2000). It has been suggested that microplastic particles will remain in suspension when reaching a depth where water and particle density are similar (Wang et al., 2016). However ocean processes are complex and other factors will influence the sinking of microplastics. Synthetic polymers vary in their elasticity (Young's modulus) from around 0.15 GPa for low density polyethylene to 4.5 GPa for polyester (measured at 20 °C and atmospheric pressure) (Ashby and Jones, 2006). The increasing pressure associated with depth will cause microplastics to compress; the depth at which this occurs and the amount of compression will depend on the specific Young modulus. It is likely that certain microplastic polymers will compress to a greater degree than the surrounding seawater (2.21 GPa at atmospheric pressure at 20 °C) (Liley et al., 2007). This will result in the microplastics increasing in density relative to the surrounding seawater, and in this way facilitate their sinking.

The occurrence of these low density polymers on the seafloor may also be explained by the hetero-aggregation of small plastic particles with other inorganic material. Coagulation is driven by the size, shape and density, along with the surface character of a particle, in turn influencing the fate of these aggregates in marine systems (Filella, 2015). In this study, 95% of the microplastics isolated were fibres. Particles such as fibres and films have high surface area to volume ratios and as such have a higher rate of aggregation/biofouling causing them to sink sooner and over relatively shorter time scales than larger plastic fragments (Ryan, 2015). Theory predicts that small sized particles will coagulate quickly (Filella and Buffle, 1993) and even buoyant material may be transported to the deep ocean relatively quickly in this way (Logan and Hunt, 1987), offering a mechanism by which microplastics may be removed from surface waters (Cozar et al., 2014). Furthermore not all sources of plastic may be introduced from land or at the ocean surface. Shipping activity, fishing equipment, offshore industries such as oil and gas and other maritime activities may introduce microplastics directly into the water column (Browne, 2015) or to the seabed and consequently reduce the time taken for microplastics to be sequestered to the seafloor. Human influences, such as deep-water bottom trawling can be traced back to 1960s in International waters to the west of Rockall Plateau (Basson et al., 2001), and by the mid-1980s the Rockall Trough had become the major deep-water bottom trawl fishing area in the northern North East Atlantic Ocean (Gordon, 2003; Heymans et al., 2011). The deep waters of the Rockall Trough are mainly comprised of Labrador seawater and tend to be slow moving (Holliday et al., 2000). Waters deeper than 1200 m are constrained due to the local bathymetry and can only exit via the southern approach of the trough (New and Smythe-Wright, 2001). These factors may promote the accumulation of microplastics within the Rockall Trough, however further work would be needed to corroborate this.

Interestingly no clear trends in overall ingested abundances or polymer abundances were observed across the time series. There was a slight increase in microplastics in 2013, however this was not statistically significant. The gap in sampling between the years 1995 - 2013 was beyond the control of the present study. To try and overcome this and enable a more in-depth assessment of microplastic load post-sampling break, 20 specimens from each species were analysed from the years 2013 and 2015. While the analysis of a complete time-series would be beneficial in drawing more robust conclusions about microplastic prevalence in the deep sea; the dataset obtained indicates stability in microplastic abundance across sampling years and additional sampling is unlikely to reveal different conclusions. Species distribution in the deep-sea is heterogeneous (Zeppilli et al., 2016) and as such the sample sizes varied between species and years. *O. lymani* dominates macrofauna assemblages in this region (Gage, 1986) enabling ≥ 10 individuals to be analysed for microplastics throughout the

time series. *H. pellucidus* was the next most abundant species collected by the sled trawls (*pers obs.*), however specimens were absent during the years 1976 and 1990 preventing their inclusion in the study and only a low number ($n = 3$) were present in 1995. It is possible that the difference in sample numbers may have influenced overall microplastic abundances or polymer diversity, however these years do not show such reductions, giving confidence to the results reported. No other macroinvertebrate species occurred in high enough abundance across the time period to be included within this study. Additional species and sample numbers would, of course, strengthen the dataset presented and facilitate drawing conclusions on the multi-decadal ingestion of microplastics.

Similar to the results presented here, Beer et al., (2017) found no trends in the abundance of microplastics ingested by pelagic fish from the Baltic sea over a 28 year dataset (1987 - 2015), however the authors did not characterise polymer types preventing comparison with our findings. In fact, the current study appears to be the first long-term (decadal/multi-decadal) investigation to characterise microplastic polymers. A total of eight different polymer types were identified; polymer diversity (H') remained relatively stable throughout the time-series, with two polymer types identified in 1976, 1990 and 2013, four polymer types in 1995 and 2015; and a maximum of 5 polymer types in 1980 and 1985. Three polymers were unique to just one sampling year; polyacrylonitrile in 1980, PVC-copolymer in 1985 and polystyrene in 2015. Acrylic, alkyd and polyethylene were found sporadically within the time series, and both polyamide and polyester fibres were present in all years sampled, with the exception of polyester in 1990. Polyester fibres are documented to be one of the most prevalent and widespread microplastics in the environment (Browne et al., 2011). Polyester and polyamide dominate fibre production, with polyester use in the textile sector seeing an annual growth of 7% since 1990; this polymer now accounts for nearly half of the global fibre market (Carr, 2017; Chemical Economics Handbook, 2016). An estimated 600 metric tons of polyester, polyamide and acrylic fibres have been discarded and are accumulating in landfill sites or the natural environment (Geyer et al., 2017), in addition to this up to 0.1 mg fibres/g polyester textile can be shed during washing (Hernandez et al., 2017), providing a source of fibre release to waste water treatment plants and to the environment (Murphy et al., 2016).

The two species examined have previously been reported to ingest microplastics (Courtene-Jones et al., 2017a). *O. lymani* is a surface deposit feeder and facultative predator on small crustaceans and polychaetes (Iken et al., 2001; Pearson and Gage, 1984), and *H. pellucidus* predaes on small benthic invertebrates and planktonic fallout (Wagstaff et al., 2014); however despite having different feeding modes and diets these factors have not been linked to the abundance of microplastic they ingest (Courtene-Jones et al., 2017a). Additionally, prior work found no detrimental effects to microplastic polymers when using 4% formaldehyde as a fixative, followed by storage in 70% ethanol (Courtene-Jones et al., 2017b), indicating the suitability of these specimens for this study. The abundances of microplastics ingested across years ($1.96 \pm 0.66 - 4.61 \pm 3.62$ microplastics/g (mean $\pm$ SD) depending on sampling year) are within a range comparable to coastal species (Devriese et al., 2015; Foekema et al., 2013; Li et al., 2016; Van Cauwenberghe et al., 2015), providing further evidence for the high levels of microplastics at the seafloor. Sample contamination is a key issue within the field of microplastics research (Wesch et al., 2017); at the time of sampling the same rigorous contamination measures put in place today would not have been used. While we cannot completely rule out historic contamination at the time of collection, thorough washing of each specimen with deionised water followed by visual examination prior to dissection was undertaken, to remove any external microplastics which may confound results. This study only assesses microplastic levels contained within the internal tissues of

these fauna where the chances for contamination arising from sample collection are minimal. Strict protocols were followed in the laboratory where samples were dissected and background controls revealed only low numbers of cellulose fibres were detected, giving confidence to the results reported here.

Only 16% of all the putative microplastics analysed were confirmed to be synthetic; thus the data presented may in fact be an under estimation of ingested quantities. The majority of particles (63%) were identified as natural based on their FTIR spectrum despite having a brightly coloured appearance (primarily blue or red). Other studies have found similar results, attributing spectra to the semi-synthetic polymer Rayon, which has an almost identical FTIR spectrum to cellulose (Lusher et al., 2015, 2014; Sadri and Thompson, 2014). To conclude whether the material is natural or Rayon would require further analysis of these particles; thus to prevent confounding our results, all cellulosic fibres (as identified with FTIR) were discounted from this study.

Microplastics have the potential to be retained and accumulate within the bodies of biota (GESAMP, 2015). Watts et al., (2014) demonstrated that following laboratory exposure crabs retained microplastics in their tissues for up to 14 days, while Browne et al., (2008) found microplastics remained in the circulatory system of mussels 48 days after high-concentration exposure. The longest study of its kind evidenced that langoustine retained microplastic fibres following low-concentration chronic exposure, which were then lost during moult, a process occurring every 6 - 12 months (Welden and Cowie, 2016). With the exception of these studies little work has been undertaken to determine the retention times, clearance rates and associated influencing factors in different organisms. Microplastics clearly have the potential to remain in the bodies of organisms which have been suggested as a short-term sink (Güven et al., 2017) however may not be retained long term due to processes of egestion. Different environmental matrices (sediment/water/fauna) likely store microplastics differently and while sediment and water may continually accumulate microplastics, it is possible that organisms may not accumulate microplastics in the same quantities as in their ambient environment. In fact it has been suggested that there may be a point at which an organism becomes saturated (Ryan et al., 2009) and does not continue to accumulate microplastics internally. This may explain why the abundances of ingested microplastics remains relatively stable in this study, despite the continued inputs of plastics into the marine environment (Geyer et al., 2017; Jambeck et al., 2015); however this is a matter requiring further work.

Conclusions

The dataset shows, for the first time, the long-term prevalence of microplastic pollution in the deep sea and documents a relatively stable abundance over the last four decades. Previous studies from marine systems have tended to provide only a snapshot in time and there is a lack of quantitative long-term data on microplastic pollution. This study provides one such assessment and indicates that microplastics may have been present on the seafloor at this locality prior to 1976. Questions remain regarding the rates of vertical transport and the dynamic processes influencing the sinking of microplastics to the benthos and we suggest this as a future priority area. Furthermore, this study shows the merit of accessing archival specimen collections to determine the historical presence of microplastic pollution. We would therefore encourage those with access to such repositories to

consider their application within this field to broaden our knowledge of the long-term fate and behaviour of microplastics within marine systems.

Funding

WCJ was jointly funded through the Scottish Association for Marine Science and the University of the Highlands and Islands. SFG was supported by NERC National Capability funding (R8-H12-85).

Acknowledgements

Thanks are extended to Fiona Ware and the National Museums Scotland for providing specimens reference NMSZ:2003.173.404 (trawl AT107, from 1976) utilised in this study; and to Dr. Sebastian Primpke for providing access to the Alfred-Wegener Institute FTIR spectral reference library prior to its publication.

References

- Andrady, A.L., 2011. Microplastics in the marine environment. *Mar. Pollut. Bull.* 62, 1596–1605. doi:10.1016/j.marpolbul.2011.05.030
- Ashby, M.F., Jones, D.R.H., 2006. Polymers, in: *Engineering Materials 2*. Butterworth Heinemann, p. 451.
- Barrows, A.P.W., Neumann, C.A., Berger, L., Shaw, S.D., 2017. Grab vs. neuston tow net: a microplastic sampling performance comparison and possible advances in the field. *Anal. Methods* 9, 1446–1453. doi:10.1039/C6AY02387H
- Basson, M., Gordon, J.D.M., Large, P., Lorange, P., Pope, J., Rackham, B., 2001. The Effects of Fishing on Deep-water Fish Species to the West of Britain 1–150.
- Beer, S., Garm, A., Huwer, B., Dierking, J., Nielsen, T.G., 2017. No increase in marine microplastic concentration over the last three decades – A case study from the Baltic Sea. *Sci. Total Environ.* 15, 1272–1279. doi:10.1016/j.scitotenv.2017.10.101
- Bergmann, M., Klages, M., 2012. Increase of litter at the Arctic deep-sea observatory HAUSGARTEN. *Mar. Pollut. Bull.* 64, 2734–2741. doi:10.1016/j.marpolbul.2012.09.018
- Bergmann, M., Wirzberger, V., Krumpfen, T., Lorenz, C., Primpke, S., Tekman, M.B., Gerdts, G., 2017. High quantities of microplastic in Arctic deep-sea sediments from the HAUSGARTEN observatory. *Environ. Sci. Technol.* acs.est.7b03331. doi:10.1021/acs.est.7b03331
- Browne, M.A., 2015. Sources and Pathways of Microplastics to Habitats, in: Bergmann, M. (Ed.), *Marine Anthropogenic Litter*. pp. 229–244. doi:10.1007/978-3-319-16510-3
- Browne, M.A., Crump, P., Niven, S.J., Teuten, E., Tonkin, A., Galloway, T., Thompson, R., 2011. Accumulation of microplastic on shorelines worldwide: sources and sinks. *Environ. Sci. Technol.* 45, 9175–9179. doi:10.1021/es201811s
- Browne, M.A., Dissanayake, A., Galloway, T.S., Lowe, D.M., Thompson, R.C., 2008. Ingested microscopic plastic translocates to the circulatory system of the mussel, *Mytilus edulis* (L.).

475 Environ. Sci. Technol. 42, 5026–5031. doi:10.1021/es800249a

476 Carr, S.A., 2017. Sources and dispersive modes of micro-fibers in the environment. Integr. Environ.
477 Assess. Manag. 13, 466–469. doi:10.1002/ieam.1916

478 Chemical Economics Handbook, 2016. Polyester fibres.

479 Chiba, S., Saito, H., Fletcher, R., Yogi, T., Kayo, M., Miyagi, S., Ogido, M., Fujikura, K., 2018. Human
480 footprint in the abyss: 30 year records of deep-sea plastic debris. Mar. Policy 1–9.
481 doi:10.1016/j.marpol.2018.03.022

482 Claessens, M., De Meester, S., Van Landuyt, L., De Clerck, K., Janssen, C.R., 2011. Occurrence and
483 distribution of microplastics in marine sediments along the Belgian coast. Mar. Pollut. Bull. 62,
484 2199–204. doi:10.1016/j.marpolbul.2011.06.030

485 Cole, M., Lindeque, P.K., Fileman, E., Clark, J., Lewis, C., Halsband, C., Galloway, T.S., 2016.
486 Microplastics alter the properties and sinking rates of zooplankton faecal pellets. Environ. Sci.
487 Technol. 50, 3239–3246. doi:10.1021/acs.est.5b05905

488 Courtene-Jones, W., Quinn, B., Gary, S.F., Mogg, A.O.M., Narayanaswamy, B.E., 2017a. Microplastic
489 pollution identified in deep-sea water and ingested by benthic invertebrates in the Rockall
490 Trough, North Atlantic Ocean. Environ. Pollut. 231, 271–280. doi:10.1016/j.envpol.2017.08.026

491 Courtene-Jones, W., Quinn, B., Murphy, F., Gary, S.F., Narayanaswamy, B.E., 2017b. Optimisation of
492 enzymatic digestion and validation of specimen preservation methods for the analysis of
493 ingested microplastics. Anal. Methods 9, 1437–1445. doi:10.1039/C6AY02343F

494 Cozar, A., Echevarria, F., Gonzalez-Gordillo, J.I., Irigoien, X., Ubeda, B., Hernandez-Leon, S., Palma,
495 A.T., Navarro, S., Garcia-de-Lomas, J., Ruiz, A., Fernandez-de-Puelles, M.L., Duarte, C.M., 2014.
496 Plastic debris in the open ocean. Proc. Natl. Acad. Sci. 111, 10239–10244.
497 doi:10.1073/pnas.1314705111

498 Crawford, C.B., Quinn, B., 2017. Microplastic separation techniques, in: Microplastic Pollutants. pp.
499 203–218. doi:10.1016/B978-0-12-809406-8.00009-8

500 Devriese, L.I., van der Meulen, M.D., Maes, T., Bekaert, K., Paul-pont, I., Frère, L., Robbens, J.,
501 Vethaak, A.D., 2015. Microplastic contamination in brown shrimp (*Crangon crangon*, Linnaeus
502 1758) from coastal waters of the Southern North Sea and Channel area. Mar. Pollut. Bull. 98,
503 179–187. doi:10.1016/j.marpolbul.2015.06.051

504 Dinno, A., 2017. Dunn’s Test of Multiple Comparisons Using Rank Sums.

505 Engel, A., 2004. Distribution of transparent exopolymer particles (TEP) in the northeast Atlantic
506 Ocean and their potential significance for aggregation processes. Deep. Res. Part I Oceanogr.
507 Res. Pap. 51, 83–92. doi:10.1016/j.dsr.2003.09.001

508 Eriksen, M., Lebreton, L.C.M., Carson, H.S., Thiel, M., Moore, C.J., Borrorro, J.C., Galgani, F., Ryan,
509 P.G., Reisser, J., 2014. Plastic pollution in the world’s oceans: more than 5 trillion plastic pieces
510 weighing over 250,000 tons afloat at sea. PLoS One 9, e111913.
511 doi:10.1371/journal.pone.0111913

512 Filella, M., 2015. Questions of size and numbers in environmental research on microplastics:
513 Methodological and conceptual aspects. Environ. Chem. 12, 527–538. doi:10.1071/EN15012

514 Filella, M., Buffle, J., 1993. Factors controlling the stability of submicron colloids in natural waters.

515 Colloids Surfaces A Physicochem. Eng. Asp. 73, 255–273.

516 Fischer, V., Elsner, N.O., Brenke, N., Schwabe, E., Brandt, A., 2015. Plastic pollution of the Kuril–
517 Kamchatka Trench area (NW pacific). Deep Sea Res. Part II Top. Stud. Oceanogr. 111, 399–405.
518 doi:10.1016/j.dsr2.2014.08.012

519 Foekema, E.M., De Gruijter, C., Mergia, M.T., van Franeker, J.A., Murk, A.J., Koelmans, A. a, 2013.
520 Plastic in north sea fish. Environ. Sci. Technol. 47, 8818–24. doi:10.1021/es400931b

521 Gage, J.D., 1986. The benthic fauna of the Rockall Trough: regional distribution and bathymetric
522 zonation. Proc. R. Soc. Edinburgh 888.

523 Gage, J.D., Lightfoot, R.H., Pearson, M., Tyler, P.A., 1980. An introduction to a sample time-series of
524 abyssal macrobenthos: methods and principle sources of variability. Oceanol. Acta 3, 169–176.

525 Gago, J., Carretero, O., Filgueiras, A. V., Viñas, L., 2018. Synthetic microfibers in the marine
526 environment: A review on their occurrence in seawater and sediments. Mar. Pollut. Bull. 127,
527 365–376. doi:10.1016/j.marpolbul.2017.11.070

528 Galgani, F., Hanke, G., Werner, S., Oosterbaan, L., Nilsson, P., Fleet, D., Kinsey, S., Thompson, R.C.,
529 van Franeker, J., Vlachogianni, T., Scoullou, M., Veiga, J.M., Palatinus, A., Matiddi, M., Maes, T.,
530 Korpinen, S., Budziak, A., Leslie, H., Gago, J., Liebezeit, G., 2013. Guidance on Monitoring of
531 Marine Litter in European Seas, EUR – Scientific and Technical Research series, European
532 Commission, EUR 26113 EN.

533 Galgani, F., Oosterbaan, L., Poitou, I., Hanke, G., Thompson, R., Amato, E., Janssen, C., Galgani, F.,
534 Fleet, D., Franeker, J. Van, Katsanevakis, S., Maes, T., 2010. Marine Strategy Framework
535 Directive: Task Group 10 Report Marine Litter., Group. doi:10.2788/86941

536 Galloway, T.S., Cole, M., Lewis, C., Atkinson, A., Allen, J.I., 2017. Interactions of microplastic debris
537 throughout the marine ecosystem. Nat. Ecol. Evol. 1, 0116. doi:10.1038/s41559-017-0116

538 GESAMP, 2015. Sources, fate and effects of microplastics in the marine environment: A global
539 assessment. Reports Stud. GESAMP 90, 96.

540 Gewert, B., Plassmann, M.M., MacLeod, M., 2015. Pathways for degradation of plastic polymers
541 floating in the marine environment. Environ. Sci. Process. Impacts 17, 1513–21.
542 doi:10.1039/c5em00207a

543 Geyer, R., Jambeck, J.R., Law, K.L., 2017. Production, use, and fate of all plastics ever made. Sci. Adv.
544 3, 25–29. doi:10.1126/sciadv.1700782

545 Goldstein, M.C., Rosenberg, M., Cheng, L., 2012. Increased oceanic microplastic debris enhances
546 oviposition in an endemic pelagic insect. Biol. Lett. 8, 817–820. doi:10.1098/rsbl.2012.0298

547 Gordon, J.D.M., 2003. The Rockall Trough, Northeast Atlantic: the cradle of deep-sea biological
548 oceanography that is now being subjected to unsustainable fishing activity. J. Northwest Atl.
549 Fish. Sci. 31, 57–83.

550 Güven, O., Gökdağ, K., Jovanović, B., Kideys, A.E., 2017. Microplastic litter composition of the Turkish
551 territorial waters of the Mediterranean Sea, and its occurrence in the gastrointestinal tract of
552 fish. Environ. Pollut. 223, 286–294. doi:10.1016/j.envpol.2017.01.025

553 Hernandez, E., Nowack, B., Mitrano, D.M., 2017. Polyester textiles as a source of microplastics from
554 households: A mechanistic study to understand microfiber release during washing. Environ. Sci.

Technol. doi:10.1021/acs.est.7b01750

Heymans, J.J., Howell, K.L., Ayers, M., Burrows, M.T., Gordon, J.D.M., Jones, E.G., Neat, F., 2011. Do we have enough information to apply the ecosystem approach to management of deep-sea fisheries? An example from the West of Scotland. *ICES J. Mar. Sci.* 68, 265–280. doi:10.1093/icesjms/fsq065

Hidalgo-ruz, V., Gutow, L., Thompson, R.C., Thiel, M., 2012. Microplastics in the Marine Environment: A Review of the Methods Used for Identification and Quantification. *Environ. Sci. Technol.* 46, 3060–3075. doi:dx.doi.org/10.1021/es2031505

Holliday, N.P., Pollard, R.T., Read, J.F., Leach, H., 2000. Water mass properties and fluxes in the Rockall Trough, 1975–1998. *Deep. Res. Part I Oceanogr. Res. Pap.* 47, 1303–1332. doi:10.1016/S0967-0637(99)00109-0

Holmström, A., 1975. Plastic films on the bottom of the Skagerack. *Nature* 255, 622–623. doi:10.1038/255622a0

Iken, K., Brey, T., Wand, U., Voigt, J., Junghans, P., 2001. Food web structure of the benthic community at the Porcupine Abyssal Plain (NE Atlantic): A stable isotope analysis. *Prog. Oceanogr.* 50, 383–405. doi:10.1016/S0079-6611(01)00062-3

Jahnke, A., Arp, H.P., Escher, B.I., Gewert, B., Gorokhova, E., Kühnel, D., Ogonowski, M., Potthoff, A., Rummel, C.D., Schmitt-Jansen, M., Toorman, E., MacLeod, M., 2017. Reducing uncertainty and confronting ignorance about the possible impacts of weathering plastic in the marine environment. *Environ. Sci. Technol. Lett.* 4, 85–90. doi:10.1021/acs.estlett.7b00008

Jambeck, J.R., Geyer, R., Wilcox, C., Siegler, T.R., Perryman, M., Andrady, A., Narayan, R., Law, K.L., 2015. Plastic waste inputs from land into the ocean. *Science* (80-). 347, 768–771. doi:10.1126/science.1260352

Jamieson, A.J., Boorman, B., Jones, D.O.B., 2013. Deep-Sea Benthic Sampling. *Methods Study Mar. Benthos* 285–347. doi:10.1002/9781118542392.ch7

John, F., Weisberg, S., Adler, D., Bates, D., Baud-Bovy, G., Ellison, S., Firth, D., Friendly, M., Gorjanc, G., Graves, S., Heiberger, R.M., Laboissiere, R., Monette, G., Murdoch, D., Nilsson, H., Ogle, D., Ripley, B., Venables, W., Winsemius, D., Zeileis, A., R Core, 2017. Companion to Applied Regression.

Kanhai, L.D.K., Gårdfeldt, K., Lyashevskaya, O., Hassellöv, M., Thompson, R.C., O'Connor, I., 2018. Microplastics in sub-surface waters of the Arctic Central Basin. *Mar. Pollut. Bull.* 130, 8–18. doi:10.1016/j.marpolbul.2018.03.011

Katija, K., Choy, C.A., Sherlock, R.E., Sherman, A.D., Robison, B.H., 2017. From the surface to the seafloor: How giant larvaceans transport microplastics into the deep sea. *Sci. Adv.* 3, e1700715. doi:10.1126/sciadv.1700715

Koelmans, A.A., Kooi, M., Law, K.L., Seville, E. Van, Quality, W., Group, M., Hole, W., 2017. All is not lost: Deriving a top-down mass budget of plastic at sea. *Environ. Res. Lett.* 12, 114028. doi:10.1088/1748-9326/aa9500

Kooi, M., van Nes, E.H., Koelmans, A.A., 2017. Ups and Downs in the Ocean: Effects of Biofouling on Vertical Transport of Microplastics. *Environ. Sci. Technol.* 51, 7963–7971. doi:10.1021/acs.est.6b04702

596 Kowalski, N., Reichardt, A.M., Waniek, J.J., 2016. Sinking rates of microplastics and potential
597 implications of their alteration by physical, biological, and chemical factors. *Mar. Pollut. Bull.*
598 109, 310–319. doi:10.1016/j.marpolbul.2016.05.064

599 Kukulka, T., Proskurowski, G., Morét-Ferguson, S., Meyer, D.W., Law, K.L., 2012. The effect of wind
600 mixing on the vertical distribution of buoyant plastic debris. *Geophys. Res. Lett.* 39.
601 doi:10.1029/2012GL051116

602 Law, K.L., Moret-Ferguson, S., Maximenko, N. a., Proskurowski, G., Peacock, E.E., Hafner, J., Reddy,
603 C.M., 2010. Plastic Accumulation in the North Atlantic Subtropical Gyre. *Science* (80-.). 329,
604 1185–1188. doi:10.1126/science.1192321

605 Law, K.L., Morét-Ferguson, S.E., Goodwin, D.S., Zettler, E.R., Deforce, E., Kukulka, T., Proskurowski,
606 G., 2014. Distribution of surface plastic debris in the eastern pacific ocean from an 11-year data
607 set. *Environ. Sci. Technol.* 48, 4732–4738. doi:10.1021/es4053076

608 Lebreton, L., Slat, B., Ferrari, F., Sainte-Rose, B., Aitken, J., Marthouse, R., Hajbane, S., Cunsolo, S.,
609 Schwarz, A., Levivier, A., Noble, K., Debeljak, P., Maral, H., Schoeneich-Argent, R., Brambini, R.,
610 Reisser, J., 2018. Evidence that the Great Pacific Garbage Patch is rapidly accumulating plastic.
611 *Sci. Rep.* 8, 4666. doi:10.1038/s41598-018-22939-w

612 Li, J., Qu, X., Su, L., Zhang, W., Yang, D., Kolandhasamy, P., 2016. Microplastics in mussels along the
613 coastal waters of China. *Environ. Pollut.* 214, 177–184. doi:10.1016/j.envpol.2016.04.012

614 Liley, P.E., Traganza, E.D., Tucker, S.P., Gordon, S., Beerbower, A., Schmidt, J.E., Momeny, A.M.,
615 Fuhs, A.E., Olien, N., Schetz, J.A., Fuhs, S.E., 2007. Properties of fluids, in: Schetz, J.A., Fuhs, A.E.
616 (Eds.), *Handbook of Fluid Dynamics and Fluid Machinery: Fundamentals of Fluid Dynamics*. p.
617 130. doi:10.1002/9780470172636

618 Lobelle, D., Cunliffe, M., 2011. Early microbial biofilm formation on marine plastic debris. *Mar.*
619 *Pollut. Bull.* 62, 197–200. doi:10.1016/j.marpolbul.2010.10.013

620 Logan, B.E., Hunt, J.R., 1987. Advantages to microbes of growth in permeable aggregates in marine
621 systems. *Limnol. Oceanogr.* 32, 1034–1048. doi:10.4319/lo.1987.32.5.1034

622 Long, M., Moriceau, B., Gallinari, M., Lambert, C., Huvet, A., Raffray, J., Soudant, P., 2015.
623 Interactions between microplastics and phytoplankton aggregates: Impact on their respective
624 fates. *Mar. Chem.* 175, 39–46. doi:10.1016/j.marchem.2015.04.003

625 Lusher, A.L., Burke, A., O'Connor, I., Officer, R., 2014. Microplastic pollution in the Northeast Atlantic
626 Ocean: Validated and opportunistic sampling. *Mar. Pollut. Bull.* 88, 325–333.
627 doi:10.1016/j.marpolbul.2014.08.023

628 Lusher, A.L., Tirelli, V., O'Connor, I., Officer, R., 2015. Microplastics in Arctic polar waters: the first
629 reported values of particles in surface and sub-surface samples. *Sci. Rep.* 5, 14947.
630 doi:10.1038/srep14947

631 Maes, T., Barry, J., Leslie, H.A., Vethaak, A.D., Nicolaus, E.E.M., Law, R.J., Lyons, B.P., Martinez, R.,
632 Harley, B., Thain, J.E., 2018. Below the surface: Twenty-five years of seafloor litter monitoring
633 in coastal seas of North West Europe (1992–2017). *Sci. Total Environ.* 630, 790–798.
634 doi:10.1016/j.scitotenv.2018.02.245

635 Murphy, F., Ewins, C., Carbonnier, F., Quinn, B., 2016. Wastewater Treatment Works (WwTW) as a
636 Source of Microplastics in the Aquatic Environment. *Environ. Sci. Technol.* 50, 5800–5808.
637 doi:10.1021/acs.est.5b05416

638 New, A.L., Smythe-Wright, D., 2001. Aspects of the circulation in the rockall trough. *Cont. Shelf Res.*
639 21, 777–810. doi:10.1016/S0278-4343(00)00113-8

640 Oksanen, A.J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R., O'Hara,
641 R.B., Simpson, G.L., Solymos, P., Henry, M., Stevens, H., Szoecs, E., Wagner, H., 2018.
642 *Community Ecology Package*. doi:ISBN 0-387-95457-0

643 Passow, U., Shipe, R.F., Murray, A., Pak, D.K., Brzezinski, M.A., Alldredge, A.L., 2001. The origin of
644 transparent exopolymer particles (TEP) and their role in the sedimentation of particulate
645 matter. *Cont. Shelf Res.* 21, 327–346. doi:10.1016/S0278-4343(00)00101-1

646 Pearson, M., Gage, J.D., 1984. Diets of some deep-sea brittle stars in the Rockall Trough. *Mar. Biol.*
647 82, 247–258.

648 PlasticsEurope, 2017. *Plastics – the Facts 2017* An analysis of European plastics production, demand
649 and waste data.

650 Porter, A., Lyons, B.P., Galloway, T.S., Lewis, C., 2018. Role of Marine Snows in Microplastic Fate and
651 Bioavailability. *Environ. Sci. Technol.* 52, 7111–7119. doi:10.1021/acs.est.8b01000

652 Primpke, S., Wirth, M., Lorenz, C., Gerdts, G., 2018. Reference database design for the automated
653 analysis of microplastic samples based on Fourier transform infrared (FTIR) spectroscopy. *Anal.*
654 *Bioanal. Chem.* 410, 5131–5141. doi:10.1007/s00216-018-1156-x

655 R Core Team, 2016. *R: A language and environment for statistical computing*. R Foundation for
656 Statistical Computing.

657 Rockström, J., Steffen, W., Noone, K., Persson, Å., Chapin, F.S., Lambin, E., Lenton, T.M., Scheffer, M.,
658 Folke, C., Schellnhuber, H.J., Nykvist, B., de Wit, C.A., Hughes, T., van der Leeuw, S., Rodhe, H.,
659 Sörlin, S., Snyder, P.K., Costanza, R., Svedin, U., Falkenmark, M., Karlberg, L., Corell, R.W., Fabry,
660 V.J., Hansen, J., Walker, B., Liverman, D., Richardson, K., Crutzen, P., Foley, J., 2009. Planetary
661 boundaries: Exploring the safe operating space for humanity. *Ecol. Soc.* 14. doi:10.5751/ES-
662 03180-140232

663 Rummel, C.D., Jahnke, A., Gorokhova, E., Kühnel, D., Schmitt-Jansen, M., 2017. Impacts of biofilm
664 formation on the fate and potential effects of microplastic in the aquatic environment. *Environ.*
665 *Sci. Technol. Lett.* 4, 258–267. doi:10.1021/acs.estlett.7b00164

666 Ryan, P.G., 2015. The importance of size and buoyancy for long-distance transport of marine debris.
667 *Environ. Res. Lett.* 10, 84019. doi:10.1088/1748-9326/10/8/084019

668 Ryan, P.G., Moore, C.J., van Franeker, J. a, Moloney, C.L., 2009. Monitoring the abundance of plastic
669 debris in the marine environment. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 364, 1999–2012.
670 doi:10.1098/rstb.2008.0207

671 Sadri, S.S., Thompson, R.C., 2014. On the quantity and composition of floating plastic debris entering
672 and leaving the Tamar Estuary, Southwest England. *Mar. Pollut. Bull.* 81, 55–60.
673 doi:10.1016/j.marpolbul.2014.02.020

674 Schlining, K., von Thun, S., Kuhn, L., Schlining, B., Lundsten, L., Jacobsen Stout, N., Chaney, L.,
675 Connor, J., 2013. Debris in the deep: Using a 22-year video annotation database to survey
676 marine litter in Monterey Canyon, central California, USA. *Deep Sea Res. I* 79, 96–105.
677 doi:10.1016/j.dsr.2013.05.006

678 Shah, A.A., Hasan, F., Hameed, A., Ahmed, S., 2008. Biological degradation of plastics: A

comprehensive review. *Biotechnol. Adv.* 26, 246–265. doi:10.1016/j.biotechadv.2007.12.005

Summers, S., Henry, T., Gutierrez, T., 2018. Agglomeration of nano- and microplastic particles in seawater by autochthonous and de novo-produced sources of exopolymeric substances. *Mar. Pollut. Bull.* 130, 258–267. doi:S0025326X18302042

Taylor, M.L., Gwinnet, C., Robinson, L.F., Woodall, L.C., 2016. Plastic microfibre ingestion by deep-sea organisms. *Sci. Rep.* 6. doi:10.1038/srep33997

ter Halle, A., Ladirat, L., Gendre, X., Goudouneche, D., Pusineri, C., Routaboul, C., Tenailleau, C., Duployer, B., Perez, E., Goudouneche, D., Pusineri, C., Routaboul, C., Tenailleau, C., Duployer, B., Perez, E., 2016. Understanding the Fragmentation Pattern of Marine Plastic Debris. *Environ. Sci. Technol.* 50, 5668–5675. doi:10.1021/acs.est.6b00594

Thompson, R.C., Olsen, Y., Mitchell, R.P., Davis, A., Rowland, S.J., John, A.W.G., McGonigle, D., Russell, A.E., 2004. Lost at sea: where is all the plastic? *Science* 304, 838.

Thompson, R.C., Swan, S.H., Moore, C.J., vom Saal, F.S., 2009. Our plastic age. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 364, 1973–1976. doi:10.1098/rstb.2009.0054

Van Cauwenberghe, L., Claessens, M., Vandegehuchte, M.B., Janssen, C.R., 2015. Microplastics are taken up by mussels (*Mytilus edulis*) and lugworms (*Arenicola marina*) living in natural habitats. *Environ. Pollut.* 199, 10–17. doi:10.1016/j.envpol.2015.01.008

Van Cauwenberghe, L., Vanreusel, A., Mees, J., Janssen, C.R., 2013. Microplastic pollution in deep-sea sediments. *Environ. Pollut.* 182, 495–499. doi:10.1016/j.envpol.2013.08.013

van Sebille, E., Wilcox, C., Lebreton, L., Maximenko, N., Hardesty, B.D., van Franeker, J.A., Eriksen, M., Siegel, D., Galgani, F., Law, K.L., 2015. A global inventory of small floating plastic debris. *Environ. Res. Lett.* 10, 124006. doi:10.1088/1748-9326/10/12/124006

Wagstaff, M.C., Howell, K.L., Bett, B.J., Billett, D.S.M., Brault, S., Stuart, C.T., Rex, M.A., 2014. β -diversity of deep-sea holothurians and asteroids along a bathymetric gradient (NE Atlantic). *Mar. Ecol. Prog. Ser.* 508, 177–185. doi:10.3354/meps10877

Wang, J., Tan, Z., Peng, J., Qiu, Q., Li, M., 2016. The behaviors of microplastics in the marine environment. *Mar. Environ. Res.* 113, 7–17. doi:10.1016/j.marenvres.2015.10.014

Watts, A.J.R., Lewis, C., Goodhead, R.M., Beckett, S.J., Moger, J., Tyler, C.R., Galloway, T.S., 2014. Uptake and Retention of Microplastics by the Shore Crab *Carcinus maenas*. *Environ. Sci. Technol.* 48, 8823–30. doi:10.1021/es501090e

Welden, N.A.C., Cowie, P.R., 2016. Environment and gut morphology influence microplastic retention in langoustine, *Nephrops norvegicus*. *Environ. Pollut.* 214, 859–865. doi:10.1016/j.envpol.2016.03.067

Wesch, C., Elert, A.M., Wörner, M., Braun, U., Klein, R., Paulus, M., 2017. Assuring quality in microplastic monitoring: About the value of clean-air devices as essentials for verified data. *Sci. Rep.* 7, 1–8. doi:10.1038/s41598-017-05838-4

Woodall, L.C., Sanchez-Vidal, A., Canals, M., Paterson, G.L.J., Coppock, R., Sleight, V., Calafat, A., Rogers, a. D., Narayanaswamy, B.E., Thompson, R.C., 2014. The deep sea is a major sink for microplastic debris. *R. Soc. Open Sci.* 1, 140317.

Zeppilli, D., Pusceddu, A., Trincardi, F., Danovaro, R., 2016. Seafloor heterogeneity influences the

719 biodiversity-ecosystem functioning relationships in the deep sea. *Sci. Rep.* 6, 1–12.
720 doi:10.1038/srep26352

721 Zhang, H., 2017. Transport of microplastics in coastal seas. *Estuar. Coast. Shelf Sci.* 199, 74–86.
722 doi:10.1016/j.ecss.2017.09.032

723 Zhao, S., Danley, M., Ward, J.E., Li, D., Mincer, T.J., 2016. An approach for extraction,
724 characterization and quantitation of microplastic in natural marine snow using Raman
725 microscopy. *Anal. Methods* 1359–1366. doi:10.1039/C5AY03217B

726