UWS Academic Portal

Potentially toxic elements in urban soils

Modabberi, Soroush; Tashakor, Mahsa; Sharifi Soltani, Neda; Hursthouse, Andrew

Published in:
Environmental Monitoring and Assessment

DOI:
[10.1007/s10661-018-7066-8](https://doi.org/10.1007/s10661-018-7066-8)

Published: 31/12/2018

Document Version
Peer reviewed version

[Link to publication on the UWS Academic Portal](#)

Citation for published version (APA):

Modabberi, S., Tashakor, M., Sharifi Soltani, N., & Hursthouse, A. (2018). Potentially toxic elements in urban soils: source apportionment and contamination assessment. *Environmental Monitoring and Assessment*, 190(12), [715]. <https://doi.org/10.1007/s10661-018-7066-8>

General rights

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact pure@uws.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

This is a post-peer-review, pre-copyedit version of an article published in ENVIRONMENTAL MONITORING AND ASSESSMENT. The final authenticated version is available online at: <https://doi.org/10.1007/s10661-018-7066-8>

Springer Nature terms of reuse for archived author accepted manuscripts (AAMs) of subscription articles, books and chapters: <https://www.springer.com/gp/open-access/authors-rights/aam-terms-v1>

For articles, books and chapters published within the Springer Nature group of companies that have been archived into academic repositories such as institutional repositories, PubMed Central and its mirror sites, where a Springer Nature company holds copyright, or an exclusive license to publish, users may view, print, copy, download and text and data-mine the content, for the purposes of academic research, subject always to the full conditions of use. The conditions of use are not intended to override, should any national law grant further rights to any user.

Conditions of use

Articles, books and chapters published within the Springer Nature group of companies which are made available through academic repositories remain subject to copyright. Any reuse is subject to permission from Springer Nature. The following restrictions on reuse of such articles, books and chapters apply:

Academic research only

1. Archived content may only be used for academic research. Any content downloaded for text based experiments should be destroyed when the experiment is complete.

Reuse must not be for Commercial Purposes

2. Archived content may not be used for purposes that are intended for or directed towards commercial advantage or monetary compensation by means of sale, resale, licence, loan, transfer or any other form of commercial exploitation ("Commercial Purposes").

Wholesale re-publishing is prohibited

3. Archived content may not be published verbatim in whole or in part, whether or not this is done for Commercial Purposes, either in print or online.
4. This restriction does not apply to reproducing normal quotations with an appropriate citation. In the case of text-mining, individual words, concepts and quotes up to 100 words per matching sentence may be reused, whereas longer paragraphs of text and images cannot (without specific permission from Springer Nature).

Moral rights

5. All reuse must be fully attributed. Attribution must take the form of a link - using the article DOI - to the published article on the journal's website.
6. All reuse must ensure that the authors' moral right to the integrity of their work is not compromised.

Third party content

7. Where content in the document is identified as belonging to a third party, it is the obligation of the user to ensure that any reuse complies with copyright policies of the owner.

Reuse at own risk

8. Any reuse of Springer Nature content is at your own risk and Springer Nature accepts no liability arising from such reuse.

Terms checked: 29 August 2018

Soroush Modabberi^{1*}, Mahsa Tashakor², Neda Sharifi Soltani³, Andrew S Hursthouse⁴

Potentially Toxic Elements in Urban Soils: Source Apportionment and Contamination Assessment

*1. School of Geology, College of Science, University of Tehran, Tehran, Iran. modabberi@ut.ac.ir

School of Geology, University of Tehran, Enghelab Avenue, Tehran, Iran. Phone: +98-9122145463,

ORCID: 0000-0001-6300-6475

2. Researcher in Environmental Geochemistry, mahsita.mt@gmail.com, ORCID: 0000-0003-2342-9857

3. PhD candidate, Department of Environmental Sciences, Macquarie University, NSW 2109, Australia,

neda.sharifi-soltani@hdr.mq.edu.au

4. School of Science & Sport, University of the West of Scotland, UK. Andrew.Hursthouse@uws.ac.uk, ORCID:

0000-0003-3690-2957

Abstract

Soils play a vital role in the quality of the urban environment and the health of its residents. City soils and street dusts accumulate various contaminants and particularly potentially toxic elements (PTEs) from a variety of human activities. This study investigates the current condition of elemental concentration in the urban soils of Hamedan, the largest and the fastest-growing city in western Iran. Thirty-four composite soil samples were collected from 0-10 cm topsoil of various land-uses in Hamedan city and were analyzed for total concentration of 63 elements by ICP-MS. The possible sources of elemental loadings were verified using multivariate statistical methods (principal component analysis and cluster analysis) and geochemical indices. The spatial variability of the main PTEs were mapped using geographic information system (GIS) technique. The results revealed a concentration for As, Co, Cr, Mn, Mo, Ni and V in the soil samples comparable to the background values as well as a range of associations among these elements in a single component suggesting geogenic sources related to geological and pedogenic processes. Whilst, the soils mostly presented a moderate to considerable enrichment/contamination of Cd, Zn, Pb, and Sb and moderate enrichment/contamination of Cu, Zn and Mo. It was found that anthropogenic factors, vehicular traffic in particular, control the concentration of a spectrum of elements that are typical of human activities, i.e. Cd, Cu, Hg, Pb, Sb and Zn. Lead and Sb were both the most enriched elements in soils with no correlation with land use highlighting general urban emissions over time and the impact of transport networks directly on soil quality. The highest concentrations of As was recorded in the southern part of the city reflecting the influence of metamorphic rocks. The effect of the geological substrate on the Co and Ni contents was confirmed by their maximum concentrations in the city's marginal areas. However, high spatial variability of urban elements' contents displayed the contribution of various human activities. In particular, the increased concentration of Cd, Sb and Pb was found to be consistent with the areas where vehicular traffic is heaviest.

Keywords: *Urban geochemistry; Soil Contamination; Environmental Geochemistry; Multivariate Statistics; Enrichment Factor; Contamination Factor.*

1. Introduction

Urban areas are the most densely-populated areas with the geographical focus of contamination discharging from various anthropogenic sources (Acosta et al. 2009; Pickett et al. 2011; Tume et al. 2018). These sources can include

one or more of: transport sources (motor exhausts, brake pads, tire wear); commercial and industrial emissions (energy production, metallurgical industry, electronics, chemical plant, fuel combustion, incinerators, etc.); domestic activities (construction and demolition, waste disposal, wastewater); and agricultural operations (application of fertilizers, pesticides, wastewater irrigation) (Wei and Yang 2010; Patinha et al. 2015; Rate 2018; Argyraki and Kelepertzis 2014; Hu et al. 2013; Li et al. 2017). Emission of harmful substances, in particular Potentially Toxic Elements (PTEs) has become a worldwide environmental concern in urban areas as a result of rapid industrialization and urbanization (Cheng et al. 2014; Johnson and Ander 2008; Manta et al. 2002; Su 2014; Yuan et al. 2014).

Although some elements are essential micronutrients and are considered beneficial for human and plant health and growth, their enrichment may reach levels which become a potential health risk (Fordyce and Ander 2003). In contrast to organic contaminants, PTEs are persistent in the environment and their concentrations remain stable or may accumulate over time (Lee et al. 2006). The prolonged presence of PTEs represents a significant challenge for the health of the urban population (Woszczyk et al. 2018; Micó et al. 2006; Wuana and Okieimen 2011; Guney et al. 2017; Lu et al. 2014; Tang et al. 2013). Direct inhalation, ingestion, dermal contact, drinking of contaminated water and food chain are the principal pathways of human exposure to PTE contaminants in urban environments (Abrahams 2002; McLaughlin et al. 2000; Poggio et al. 2009).

As an essential compartment of the urban ecosystem and sink for PTEs, soils contribute directly or indirectly to the general citizens' quality of life (Biasioli et al. 2006; Christoforidis and Stamatis 2009). Potentially toxic elements from local geogenic sources and from atmospheric deposition of dusts and suspended particles may also be incorporated into urban soil compartment (Norra 2009; Xia et al. 2011). Many researchers have focused on the necessity for a better understanding of urban soil and dust pollution (Hu et al. 2013; Li et al. 2017; Manta et al. 2002). Urban soils are known to have distinctive features such as heterogeneity, high spatial variability of chemical, physical, and biological properties, unpredictable layering, rapid change in land use, poor structure, fragmented distribution and high content of trace elements (Ajmone-Marsan et al. 2010; Kabata-Pendias and Mukherjee 2007; Manta et al. 2002; Xia et al. 2011). For this reason, the traditional approach in environmental survey, used in intact natural environment may not be appropriate in city-wide assessments (Wong et al. 2006). In addition, population, time, location, urban sprawl, intensity of human activities and micro-environmental parameters such as topography, wind direction, and urban runoff may give rise to micro-environmental variation (Amato et al. 2009; Chen et al. 2005).

The characterization of soil and dusts for metal contamination assessment is an effective scientific tool which provides an insight into the source(s) of contamination and enables policy-makers to more effectively manage urban sites in order to protect public and ecosystem health.

This study investigates the geochemistry of urban soils in Hamedan city of Iran where is believed to have an active and increasing pollution burden on its basic infrastructure due to increasing population and industrial growth in previous two decades. The main objectives of this study are: 1) to determine the total concentration of potentially toxic elements (focusing on those with environmental implications) in urban soil under different land uses from Hamedan city; 2) to assess the degree of Hamedan soil pollution in comparison with uncontaminated sub-urban soils and soils from other cities throughout the world as well as applying contamination factor (CF) and enrichment factor (EF); 3) to identify the possible sources (natural and/or anthropogenic) of PTEs in soils and spatial distribution of the accumulated elements.

2. Materials and methods

2.1 Study Area

Hamedan city, in west of Iran (34°48' N, 48°31' E), is the official and political center of Hamedan province of Iran. Hamedan is an ancient city with a history of over 3000 years and one of the oldest capitals in the world. Hamedan was the capital city of Median Empire which was conquered in 550 B.C by Cyrus the Great (Jabarivasal 2012). The municipality spreads over an area of about 60 km² with a population of about 600,000 according to a national census in 2016. Considering the relative density of population, Hamedan ranks sixth in the country and it is known as a metropolis in February 2010 (Shojaimehr and Zakerhaghighi, 2013). Hamedan is an important center of economy, transportation and manufacturing. Industrial zones of the city are particularly located at a distance of at least 10km eastward from the city border mainly consisting of small-scale downstream metal industries such as slabs, electrics, car spares, stone polishing etc. Hamedan has experienced a significant urbanization rate during the past 20 years. Considering the ratio of families having cars in the country, it is estimated that Hamedan contains approximately 85000 private vehicles. According to Hamedan news agencies, the local vehicular fleet has increased to more than 4000 public vehicles of which about 390 are public buses and remaining are taxis.

Geographically, Hamedan is located in the northern foothills of the Mt. Alvand at an altitude of 1813 meters above sea level (Shojaimehr and Zakerhaghighi, 2013). The climate of Hamedan is cold and semi-arid with an

average annual rainfall of about 330 mm and a mean temperature of 11°C. The prevailing wind direction is from southeast to northwest.

From the geological point of view, Hamedan is located in the Sanandaj-Sirjan structural zone (SSZ) of the Zagros Mountain Ranges. This area is best characterized by Alvand intrusive complex composed of granite to granodiorite bodies (Fig. 1) outcropped at the surface toward the southwestern side of the city, where Ganjnameh and Abbas-Abad valleys are on view. Metamorphic processes related to Alvand magmatism has produced various facies of slate, phyllite, schist and hornfels rocks which are exposed in the southern part of the city. Hamedan is underlain by the Quaternary alluvial deposits and alluvial fans derived from weathering of the Alvand granitic and metamorphic rocks (Fig. 1).

2.2 Sample Collection and Analytical Methods

Topsoil samples (0–10 cm depth) were randomly collected from all open, unrestricted lands in Hamedan. Samples were taken at 31 locations from residential (8 samples), green spaces (11 samples), agricultural lands (6 samples), parks (4 samples), and playgrounds (2 samples). Four street dust samples were collected from pavements with small plastic brush and pan. In addition, three samples were taken from suburb far from any pollution sources to represent the natural background concentrations. The location of sampling points is shown in Fig. 1. The uppermost topsoil layer is of particular interest in urban geochemistry studies because of the likelihood of atmospheric aerosol deposition from anthropogenic sources. Composite samples were obtained at each sampling site by thoroughly mixing five sub-samples collected at corners and center of a 2×2 m square by using a stainless steel shovel to have a representative sample with a weight of 2kgs. Vegetation debris and coarse-grained particles were manually removed before packing the samples. Soil samples were stored in polyethylene zip-lock bags and transported to laboratory. Soils were dried at room temperature and were then disaggregated to pass through 2 mm nylon sieve mesh for analysis.

The values of pH, EC and TDS were measured using a calibrated instrument in an aqueous suspension of 10g soil and 25ml distilled water according to BS 1377: Part 3:9.0 (1990b) in Zamin Kavan Jonoub Geotechnical Lab, Tehran, Iran.

The organic matter content was measured by Walkley-Black titration method based on the oxidation of organic matter by $K_2Cr_2O_7$ (Sims and Wolf 1995).

The grain size of soil samples was determined by dry sieve analysis based on the particle size distribution procedure provided by BS 1377: Part 2: 9.4 (1990).

The concentration of major and trace elements in soil samples was measured using aqua regia extractions. A 0.5g sample was digested at 90°C in a microprocessor controlled digestion block for 2h. Digested samples were diluted and analyzed for 63 elements by Perkin Elmer Sciex ELAN ICP/MS in ActLabs analytical laboratory, Canada. For quality control and quality assurance purposes, three duplicate and one blank samples were analyzed. The analytical data represents reasonable and good precision mostly less than 8% and never higher than 10%.

2.3 Statistical Methods

The geochemical results were analyzed for descriptive and multivariate statistics using R3.3.1 and SPSS 23.0 software. First of all, the distribution of data was tested with the Shapiro-Wilk's test ($p < 0.05$). Due to the non-normal distribution of the data, nonparametric statistics was used. Nonparametric kruskal-wallis test was conducted to determine the differences of PTEs concentration among the five different land uses. The degree of correlation between the measured elements was determined with Spearman correlation matrix ($p < 0.05$, $p < 0.01$). Additionally, principal component analysis (PCA) was employed to elucidate the possible sources of PTEs in the Hamedan surface soils. In the PCA, different groups of elements were categorized based on the concept of communality for each element. PCA is a powerful tool to differentiate pollution. The varimax rotation method was used in this analysis to improve the value of factor matrix. Hierarchical cluster analysis (CA) was applied for the same variables to differentiate the geochemical groups using the Ward's linkage method and the results were presented in a dendrogram with Squared Euclidean distances.

2.4 Geochemical Indices

Enrichment factor (EF) is widely used to identify the contribution of anthropogenic sources in soils (Li et al. 2017). EF of individual elements in soils is calculated as:

$$EF = [C_i/C_{Sc}]_{\text{sample}} / [C_i/C_{Sc}]_{\text{background}}$$

in which:

C_i and C_{Sc} are concentrations of the target and reference element in soil sample and in the background sample. The most commonly used reference elements are Al, Fe, Me, Mn, Sc, Ti (Reimann and Caritat 2000; Yongming et al.

2006). Scandium was preferred to Al and Fe and was selected as the reference element in this study since it has no industrial and/or geogenic origin.

Five levels of enrichment can be categorized on the basis of EF (Qingjie et al. 2008):

- $EF < 2$ deficiency or minimal enrichment,
- $2 < EF < 5$ moderate enrichment,
- $5 < EF < 20$ significant enrichment,
- $20 < EF < 40$ very high enrichment,
- $EF > 40$ extremely high enrichment.

Concentration factor (CF) is the quotient of concentration of each element in soil sample (C_n) and the concentration of the same element in reference soil (B_n) (Dung et al. 2013).

$$CF = C_n / B_n$$

Reference soil can be either the world's mean concentration or determined locally from an unpolluted site. The pollution status of elements is classified into four levels from unpolluted to very polluted as below (Abraham and Parker 2008):

- $CF < 1$ low contamination;
- $1 \leq CF < 3$ moderate contamination;
- $3 < CF < 6$ considerable contamination;
- $CF > 6$ very high contamination

Since the composition of soil samples is varied from site to site, in this study local background values were used to calculate the corresponding degree of EF and CF.

The spatial distribution patterns (geochemical maps) showing the hot spots of elevated concentrations were produced using ArcGIS 10.5 software.

3. Results and Discussion

3.1 Physico-Chemical Parameters

The physico-chemical parameters were determined for Hamedan urban soils. The pH values range from 6 to 7.25 with a mean value of 6.87, which suggest circum-neutral acidity for all the topsoils. Despite the fact that soils of Iran

are generally alkaline, the pH of urban soils of Hamedan indicates the influence of the Alvand granite complex. The urban soils of Hamedan have a pH similar to the background with low variation between different land uses.

The electrical conductivity (EC) ranges broadly with a median of 2650 μ S/cm. The largest EC was measured in residential soils (67300), decreasing in green spaces (EC_{Ave} =5076), parks (EC_{Ave} =2143) and agricultural lands (EC_{Ave} =1938). Playgrounds showed the lowest EC values (EC_{Ave} =1429). The high EC values of soils in residential areas are related to the high calcium content in cement and concrete and gypsum which are commonly found in demolition and construction wastes. Organic matter contents vary from 0.4% to 12.9% with a median of 2.8%. The maximum and the minimum values belong to parks and residential lands, respectively, reflecting active use of soil as a growing medium. Sand is the predominant fraction in 78% of the urban soil samples, showing a silty-sand texture. The rest 22% of the samples show a sandy-silt texture. The soil samples contain 9.89 to 29.9% clay size fraction. Particle size distribution of Hamedan urban soils follows the land use classes of the city.

3.2 PTEs Concentrations

Sixty-three elements were analyzed in all soil samples (Appendix 1); of these, 13 elements were selected for further analysis based on their known significance in the urban environment (Charlesworth et al. 2011; Chen et al. 2005; Giusti 2011; Ljung et al. 2006; Wong et al. 2006; Yesilonis et al. 2008). The mean concentrations of As, Ba, Co, Cr, Mn, Mo, Ni, V and Zn in the soil samples were 18.9, 124, 13.1, 56.4, 683, 0.77, 54.9, 61.1 and 115.2 mg kg⁻¹, respectively. These values were compared to the background values of the elements in soils from remote suburb of Hamedan as well as the mean values of the elements from studies in different cities around the world, as shown in Table 1. It is universally admitted that comparing concentrations of elements in urban soils with values measured in other cities is an advantageous practice to better estimate the status of contamination (Duzgoren-Aydin, 2007).

When compared with the background concentration of the PTEs in the suburban soil of Hamedan, As, showed mean concentrations comparable to the background values, suggesting a geogenic source for this element. Mean concentrations of As in urban and suburban soils of Hamedan are 9 times greater than those in the uncontaminated soils worldwide reported by Kabata-Pendias and Mukherjee (2007). Moreover, it is observed that the content of As measured in urban and background soil samples of Hamedan is above the mean values when compared to 11 other cities around the world. This can be attributed to the natural enrichment of arsenic in Sanandaj-Sirjan plutono-metamorphic belt. Concentrations of Co, Cr, Ni and V in the soils of Hamedan city are below the observed values in suburban areas indicating a geogenic source for these elements. On the other hand, the elements with higher

concentration in the urban soil samples compared to the background (e.g. Cd, Cu, Hg, Pb, Sb and Zn) can be possibly connected to certain anthropogenic activities in addition to the original content from the parent soils.

In comparison to the background soil value (0.07 mg kg^{-1}), Cd shows 2 fold rise in urban soils (0.14 mg kg^{-1}), though, this is still below the worldwide values for Cd (0.53 mg kg^{-1}). In a recent study, Solgi et al. (2016) reported a concentration of 0.23 to 2.30 mg kg^{-1} for Cd in urban park soils of Hamedan. The average content of Cu in Hamedan soils, both city (39 mg kg^{-1}) and background (36.5 mg kg^{-1}), are greater than those in most of the reported values in other cities. This indicates that in addition to geogenic sources, extra origin(s) from human activities added copper to the urban soils. Similarly, slight increase is observed for the amount of Zn in Hamedan city soils (115 mg kg^{-1}) in comparison to the background (90.6 mg kg^{-1}) and uncontaminated soils (100 mg kg^{-1}) values, suggesting possible anthropogenic sources of Zn in the soils. As can be seen in Table 1, mean concentration of Hg in urban soils is about 11 times higher than the local background levels, representing a significant anthropogenic source for Hg in Hamedan city. Pb and Sb are similarly enriched up to 3 times in the urban soil samples.

The four street dust samples revealed higher concentrations of some PTEs, so that the highest concentration of Cd (0.35 mg kg^{-1}) and Hg (1.340 mg kg^{-1}) were found in SD17 and the highest content of Mo (1.15 mg kg^{-1}) and Pb (289 mg kg^{-1}) were measured in SD21 dust samples.

As can be noted from the mean values listed in Table 2, variation in land use showed little effect on distribution of elements in urban soils of Hamedan, confirmed statistically (kruskal-wallis H) that there is no significant difference in concentration of PTEs in various land use types ($p\text{-value} > 0.05$). However, it is observed that agricultural land contains high contents of As, Co, Cr, Cu and Ni. Pb and Sb are slightly more concentrated in green space soils. Zn and V are dominant elements in parks and residential areas, respectively.

3.3 Evaluation of the Strength of Soil Pollution

Enrichment factors for PTEs in urban soils of Hamedan are calculated and illustrated in the boxplots Fig. 2.

Almost all of the EF values for As, Co, Cr, Mn, Ni and V in Hamedan soils are below 2 confirming the deficiency or minimal enrichment of these elements and the influence of the local geology. Cadmium has moderately enriched in 32% of the soil samples ($2.22 < \text{EF} < 4.84$), while 19% of the samples are significantly enriched in Cd ($5.23 < \text{EF} < 11.26$). The mean EF value for Cd is 3.23. Seven of the 31 analyzed soils fall in the category of “moderate enrichment” for Cu, with the EF values ranging from 2.14 to 3.95. However, the mean value

of EF for Cu (1.75) shows “minimal enrichment”. Similar results are shown for Mo with 10 moderately enriched soil samples (32%) and the mean EF value of 1.81. Lead appeared to be the most serious enriched element in soils of Hamedan city with the EF mean value of 6.82. More than half of the soil samples are moderately enriched and 26% are significantly enriched in Pb ($2.59 < EF < 4.83$ and $5.34 < EF < 14.44$, respectively). The enrichment of Pb and Cd in Hamedan urban soils has already been identified in previous papers (Solgi et al. 2016; Yeganeh et al. 2012).

Almost all of the analyzed soils (97%), present various levels of enrichment in antimony. 51% of the samples fall into the category of moderate Sb enrichment with the minimum and the maximum observed EF values of 2.04 and 4.62, respectively. Significant enrichment in Sb is observed in 45% of the samples including the EF values ranging from between 5-13.48. One of the samples was very highly Sb enriched, showing the EF value of 22. In average, the enrichment factor of Hamedan soils for Sb is 6.15. Zinc is enriched in 42% of the samples (13 soil samples). Among these, 31% (11 samples) show moderate and the rest show significant enrichments.

Dust street samples including one collected from downtown Shir-Sangi square of Hamedan show the highest EF levels, so that it was extremely enriched showing the EF value of 54.31 for Pb.

The assessment of contamination factor for PTEs of Hamedan urban soils is depicted in Fig. 3. Similar to the results of enrichment factor analysis, the geogenic elements of As, Co, Cr, Mn, Ni and V group together and show the least contamination with the average contamination factor of less than 1. Contamination factor for Cd in about 42% of the samples ranges from 1.2 to 2.9 which indicate moderate contamination for this element. Five of the 31 soil samples with CF values between 3.07 and 3.87 are considerably contaminated by cadmium. For Cu, a moderate contamination is observed in 45% of the samples where the CF ranges between 1 and 2.17. More than half of the samples are moderately contaminated by Mo showing the minimum and maximum CF values of 1.01 and 1.63. The highest contamination factor is observed for Pb and Sb with the same average CF of 3.7. About 58 and 55% of the samples fall in the category of “moderate contamination” for Pb and Sb, respectively. Pb in 26% and Sb in 29% of the analyzed samples are considerably high by showing the respective CF values of 3.15-5.85 and 3.26-4.36. Very high contamination of Pb ($5.49 < CF < 21.09$) and Sb ($6.13 < CF < 13.64$) is observed in five and four of the analyzed soil samples, respectively. With respect to Zn, the majority of samples (67%) are within the category of “moderate contamination” and the remaining are low.

The elements with some degrees of enrichment or contamination with respect to the background level probably suggest the influence of human interventions.

3.4 Interrelationship of PTEs

3.4.1. Bivariate statistics

The correlation between the elements can provide some information on their origin in a metropolitan area like Hamedan. Prior to correlation tests, normal distribution of elements was checked by shapiro-wilk test, as correlation coefficient depends on the linear combinations of the variables. The result of Spearman's correlation matrix of the PTEs for the surface soil of Hamedan city is shown in Table 3. As it is indicated, the correlation coefficient between Co and Cr is 0.880, which represents a strong, positive, linear correlation at the 0.01 significance level which shows their close association in most soil samples. In addition, Co shows strong positive correlation with both Ni (0.893) and V (0.717). Cr-Ni and Cr-V form two other highly correlated pairs with a correlation coefficient of 0.836 and 0.766, respectively. In addition, moderately positive correlations are shown between As and Co (0.566), Cr (0.473), Ni (0.526) and V (0.492). As it is mentioned before, As, Co, Cr, Ni and V occur naturally at abundant levels in suburban soils of Hamedan, thus the association of these elements can be a reflection of their common source. Cu-Pb and Pb-Zn show the same considerable positive correlation of 0.825. A significant correlation between Cu with Zn (0.724) was observed due to the coexistence of these elements as a result of human activities in Hamedan city. The element pairs Cd-Zn (0.652), Cu-Hg (0.671), Hg-Pb (0.649), Hg-Sb (0.531), Hg-Zn (0.475) and Ni-V (0.532) had high to moderate positive linear correlation at 0.01 significance level, indicating the similar influential factor. Pb and Mo (0.433) and Cu and Sb (0.426) were moderately correlated at 0.05 significance level.

3.4.2. Multivariate statistics

In addition to the correlation coefficient analysis, multivariate statistics was used by principal component analysis method for assessment of the sources of PTEs in Hamedan city. Varimax rotation with Kaiser Normalization was applied. Five principal components were extracted from the available dataset accounting for over 80% of the total variance and considering the eigenvalues greater than 1. The results of the factor loading and communalities are shown in Table 4. Factor loadings bigger than 0.71 are regarded as excellent, and those smaller than 0.32 are considered as very poor. The KMO (Kaiser–Meyer–Olkin) test showed a reliable value (0.615) indicating that PTEs concentration data of Hamedan soils are suitable for principal component analysis.

The first factor (F1) explains 28.17% of the total variance and shows positive association between As, Co, Cr, Ni and V with the factor loadings of 0.635, 0.862, 0.814, 0.895, and 0.664, respectively. This factor can be an

285 indication of the influence of lithogenic inputs into the urban soils as the dominant elements are typical lithophile
 286 elements according to the Goldschmidt's classification and mineralogical components of rocks in the area. Cr and Ni
 287 derived from pedogenic materials are reported from other cities in the world (Zheng et al. 2008; Argyraki and
 288 keleptziz 2014). Association of the major elements of Al (0.844), Fe (0.698), K (0.545), Mg (0.783) confirms a
 289 geogenic control and reflects a reasonable grouping based on the similarities in the geochemical behavior of these
 290 elements. The basement of Hamedan is composed of pre-Jurassic rocks consisting of schist, hornfels and phyllite
 291 (Baharifar et al. 2004; Sepahi 2008). Association of As which is a chalcophile metalloid in F1 group elements, is
 292 referred to the Sanandaj-Sirjan plutono -metamorphic belt. This zone of the Zagros Orogen is well-known for As
 293 enrichment due to the strong relationship of arsenic with gold mineralization in this particular tectonic settings. This
 294 has been reflected in elevated amounts of As in background as well as in the urban soils of Hamedan. Khaleghian
 295 and Modabberi, (2014) and Yeganeh et al. (2012) reported the high concentration of As in agricultural soils and
 296 potatoes, wheats and Maize grown in agricultural lands around the city. Similar result obtained from urban study of
 297 Estarreja, north-western Portuguese coast where high concentration of As in the city soil were attributed to the
 298 presence of pyrite in the mineralogical suite of the region (Cachada et al. 2012). Vanadium (0.627) and Fe (0.489)
 299 are also partially presented in the second factor (F2, 12.83% of total variance) along with Mn (0.884) and Ti (0.804).
 300 The influence of parent material and natural processes is deduced for association of these elements. The third factor
 301 (F3) explaining 11.83% of the total variance is dominated by Ba, Cu, Mo and Pb with the factor loading values of
 302 0.712, 0.515, 0.503, 0.831, respectively. The F3 elements, according to many previous studies on urban soil quality,
 303 are known as urban elements and are indicators of traffic-related contamination (Adachi and Tainosho 2004; Hays et
 304 al. 2011; Rodríguez-Seijo et al. 2017; Saeedi et al. 2012; Zechmeister et al. 2005). These elements contribute in the
 305 fabrication of automobiles or as additive for engine operation or fuel or resulting from corrosion and deterioration of
 306 vehicle metal parts (Kuang et al. 2004; Dresel 2007). The concentration of Ba, for example, can be related to brake
 307 linings in cars and its emission occurs as a result of frictional processes (Fernández-Espinosa and Ternero-
 308 Rodríguez 2004). Copper compounds serve as anti-wear additive and friction-reduction agent in lubricants (Okorie
 309 et al. 2012). Even though lead used to be a significant component applying as anti-knock agent in petrol (Callender
 310 and Rice 2000), it has been phased out from gasoline in 1990s in Iran. Considering the long residence time of Pb and
 311 non-biodegradability and cumulative tendency of this element (Mielke et al. 2010), it is believed that enrichment of
 312 Pb in current urban soils represents footprints of prevalent use of tetraethyl lead in the past (Carrero et al. 2013;

Cicchella et al. 2008; Dean et al. 2017; Francek 1992; Galušková et al. 2014; Lau and Othman et al. 1997; Lin et al. 1998; MacKinnon et al. 2011; Poňavič et al. 2018; Sutherland et al. 2000; Wong 1982) and its reworking through circulation of urban dusts by local winds. The fourth factor (F4), which accounts for 9.79% of the total variance, is dominated by Cd (0.899), Hg (0.468) and Zn (0.817). The association of these elements can be mainly attributed to transportation means. Vehicle tires are main source for cadmium and zinc which may lead to contamination of urban soils (Davis et al. 2001; Mitchell et al. 2014; Smolders and Degryse 2002; Adachi and Tainosho 2004; Turner and Rice 2010; van Beers and Graedel 2007; van Bohemen and van De Laak 2003; Zheng et al. 2008). Zinc can also be derived from wearable parts on vehicle brakes (e.g. disc pads). Moreover, cadmium is typically an anthropogenic element derived from a wide range of sources in cities including tire wear, paints, pigments, electroplating, plastic stabilizer, solid wastes and wastewaters and phosphate fertilizers (Kisku et al. 2000; Romic and Romic 2003). The association of volatile element of Hg in this PC fits with a model of wider anthropogenic sources in urban environments. Yeganeh et al. (2012) has reported high concentration of Hg in Hamedan soils, water and food crops which has caused significant potential risk for the city population. The exact source of Hg in Hamedan soils is unknown to the authors. Even though copper presented in factor 3, its highest factor loading (0.702) appeared in factor 5 (F5) along with Sb with the factor loading of 0.772. Partial representation of Cu in another factor implies an ambiguous behavior and probably the presence of different anthropogenic inputs. The mean ratio of Cu:Sb which is commonly used to identify the source of antimony, differs significantly in Hamedan soils with the reported diagnostic criteria (Sternbeck et al. 2002) for brake wear particles (4.6 ± 2.3). Therefore, attribution of Sb together with Cu in brake linings is unexpected. Although the particular source of antimony in Hamedan city soils is still not clear, its significant concentration in urban soils compared to the background together with elevated EF and CF values confirm an anthropogenic source for this element. An extra source might be horticulture. As according to Pietrzak and McPhail (2004), this contributes plant macronutrients such as phosphorus in addition to certain metals like copper, as emerged in F5. Factor 6 (F6) explain 9.64% of the total variance and accounts considerable concentration of Mo (0.648). As mentioned before, molybdenum is also weakly presented in F3 displaying its diffuse origin due to the contribution of a wide range of natural and urban parameters (Peltola and Åström 2003).

To illustrate the interrelationship of the elements, the first three factors are plotted in three-dimensional space in Fig. 4, where the associations between elements can be seen.

Cluster analysis was used to visualize the classification of PTEs according to their plausible origin. The dendrograms (Fig. 5) confirm the attributions explained by PCA analysis. Two distinct clusters were identified for the potentially toxic elements in Hamedan urban soils indicating two main sources:

i) Natural: the cluster of geogenic origin which is divided in two subcategories; the upper sub-cluster is made up of V, Ti, K and Mn displaying the geochemical contribution of underlying Alvand batholith. Whilst, the other sub-cluster includes the elements (e.g. Al, Fe, Mg, As, Co, Cr, Ni) which are mainly associated with the metamorphic formations surrounding the Alvand Batholith.

ii) Mixed: in addition to the influence of the local geology in supplying a fraction of trace elements in soil samples, the other extra non-natural sources have loaded certain elements in the studied urban soils of Hamedan. Two sub-clusters were identified with typical anthropogenic elements including Cd, Pb, Zn, Cu, Sb and Hg. As it is found from the bivariate statistical analysis, the concentration of each element presented in the mixed cluster is positively and significantly correlated with the other elements belonging to the same cluster. This fact can be an indication of a common source. The divided components containing Hg-Cd-Zn, Sb-Cu and Ba-Pb-Mo in the PCA analysis is confirmed with the distinct sub-cluster are shown in the CA (Fig. 5). Moreover, as it is explained, anthropogenic elements reveal higher mean concentrations in urban soils compared to the background soils in Hamedan. These evidence along with the calculated elevated values for the enrichment and contamination factors, highlight the input of PTEs from human activities in Hamedan. This group of elements has previously been interpreted as anthropogenic by many scholars (Rodríguez-Seijo et al. 2017; Costa and Jesus-Rydin 2001; Rate 2018).

3.5 Spatial Distribution

The spatial distributions of concentrations of a number of elements in Hamedan city are visualized by a set of geochemical maps (Fig. 6) prepared by ArcGIS. The segment division displayed in the geochemical maps are according to concentrations of 5%, 25%, 75%, 95% and 100%.

In agreement to Spearman, PCA and CA results, the spatial distribution analysis suggests that the concentration of As, Co, Cr, Ni and V is most tightly controlled by soil parent materials. As it is mentioned before, Hamedan is built upon the old alluvial fans and young terraces which are derived from weathering of the metamorphic rocks from short distances in south and southwest of the city. The piedmont alluvial deposits occur along the valley margins toward the Hamedan plain and form the foothills bordered by Alvand metamorphic and plutonic highlands.

In the north, the cultivated land, consists of clay alluvial soils probably originated from metamorphic bedrocks. The development and thickness of sediments increase toward the north.

Concentration of arsenic accords with the metamorphic rock units with the highest values in the southern part of Hamedan city (Fig.6). Toward the north, due to the less influence of metamorphic rocks and continuous leaching process the content of As decreases.

As seen from the maps (Fig. 6), variations in the levels of Co and Ni demonstrate similar pattern where the influence of parent materials has been involved. The total content of Co in the study area is lower compared to Ni, though both elements show hotspots in the city's marginal areas. Despite the fact that Cr displays a spatial distribution different to Co and Ni, the strong statistical correlation among these elements can be attributed to their common natural source.

Two critical regions for Cd are recognized (Fig. 6): the high-value distribution area in north of the city may be related to the agricultural lands and the influence of fertilizers in addition to the impacts of the nearby highway. While, the hotspots in central and southwestern of the urban areas may indicate the direct influence of vehicular traffic. Ganjnameh and Abbas-Abad are roads to recreational and touristic areas in the southwest of the city with significant traffic density. The latter may be the contributing factor in higher concentration of Sb, as a traffic-generated element, appearing in the vicinity of Ganjnameh road. The valley may funneled the wind and prevented dissipation of traffic-related contaminants in the plain. As it is illustrated in the map (Fig. 6), lead is characterized by higher concentrations along the outer ring road of the city which serves both city and regional traffic with constant congestion. This is correlated with traffic volume and association of some infrastructures (garages, auto-truck service and repair shops) which are generally located around the main entrances of the city.

High spatial variability of Cu, Hg, Mo and Zn even in the short range indicates the spatial heterogeneity caused by various human activities (Liu et al. 2014). This is due in part to the natural background soil but, mainly the result of long-term complex anthropogenic processes.

4. Conclusion

Geochemical study of surface soil in Hamedan city of Iran revealed that the concentration of As, Co, Cr, Ni and V are close to the regional background values, so they may have been originated from the natural parent materials. This result is reinforced by correlation coefficients, PCA and CA analyses identifying geochemical association of

these elements which were also spatially related. Moreover, the information obtained from this study showed that Pb followed by Sb and Cd are the most enriched elements in Hamedan city soils. These elements along with Cu, Hg, Mo and Zn showed anthropogenic influence mainly related to vehicular traffic and automobile exhaust. It is deduced that the control of industrial activity is minor probably due to the remote location of Hamedan industrial zones (Bou-Ali Industrial Town) in the north of the city at a distance of about 10 km downwind. Land use showed to have little or no effect on the enrichment of PTEs in Hamedan surface soils. Spatial distribution maps displayed the contribution of metamorphic parent rocks in accumulation of arsenic in both natural and city soils. An identical pattern of special distribution was observed for Co, Ni revealing the influence of geogenic factor. In contrast, the urban elements display large variability throughout the city which indicates the interruption caused by human activities. Lead concentration, however, accords with the main highway network of the city inherited from past usage of Pb in gasoline which has been preserved in soils. Due to the traffic density in Hamedan roadways and dense network of roads the potential for ongoing inputs of PTEs exists. The study of bioavailability of concerned elements and their risk assessment is recommended for further research.

Acknowledgment

The authors would like to express their sincere gratitude to Mr. Korehei, the former deputy minister and director of the Geological Survey of Iran for financial support of this research. We are also thankful of the Zamin Kavan Jonoub company and especially Miss Sahel Sepand for sample preparation and soil quality tests.

References

- Abrahams, P., (2002). Soils: their implications to human health. *Science of the Total Environment*, 291, 1-32.
- Abraham, G., & Parker, R., (2008). Assessment of heavy metal enrichment factors and the degree of contamination in marine sediments from Tamaki Estuary, Auckland, New Zealand. *Environmental monitoring and assessment* 136, 227-238.
- Acosta, J. A., Cano, A. F., Arocena, J. M., Debela, F., & Martínez-Martínez, S. (2009). Distribution of metals in soil particle size fractions and its implication to risk assessment of playgrounds in Murcia City (Spain). *Geoderma*, 149(1-2), 101-109.
- Acosta, J., Faz, A., Martínez-Martínez, S., & Arocena, J (2011). Enrichment of metals in soils subjected to different land uses in a typical Mediterranean environment (Murcia City, southeast Spain). *Applied Geochemistry* 26, 405-414.
- Adachi, K., & Tainosho, Y. (2004). Characterization of heavy metal particles embedded in tire dust. *Environment international*, 30(8), 1009-1017.

425 Ajmone-Marsan, F., & Biasioli, M. (2010). Trace elements in soils of urban areas. *Water, Air, & Soil Pollution*,
426 213(1-4), 121-143.

427 Amato, F., Pandolfi, M., Viana, M., Querol, X., Alastuey, A., & Moreno, T. (2009). Spatial and chemical patterns of
428 PM10 in road dust deposited in urban environment. *Atmospheric Environment* 43, 1650-1659.

429 Andersson, M., Ottesen, R., & Langedal, M. (2010). Geochemistry of urban surface soils—monitoring in
430 Trondheim, Norway. *Geoderma* 156, 112-118.

431 Argyraki, A., & Kelepertzis, E. (2014). Urban soil geochemistry in Athens, Greece: the importance of local geology
432 in controlling the distribution of potentially harmful trace elements. *Science of the Total Environment* 482, 366-377.

433 Baharifar, A., Moinevaziri, H., Bellon, H., & Piqué, A. (2004). The crystalline complexes of Hamadan (Sanandaj–
434 Sirjan zone, western Iran): metasedimentary Mesozoic sequences affected by Late Cretaceous tectono-metamorphic
435 and plutonic events. *Comptes Rendus Geoscience* 336, 1443-1452.

436 Biasioli, M., Barberis, R., & Ajmone-Marsan, F. (2006). The influence of a large city on some soil properties and
437 metals content. *Science of the Total Environment* 356, 154-164.

438 British Standard Institution (1990). *Methods of Test for soils for civil Engineering Purposes-BS 1377-Part 2 9.4.*
439 *Particle size analysis*, London

440 Cachada, A., Pereira, M. E., Da Silva, E.F., & Duarte, A.C. (2012). Sources of potentially toxic elements and
441 organic pollutants in an urban area subjected to an industrial impact. *Environmental monitoring and assessment*,
442 184(1), 15-32.

443 Callender, E., & Rice, K. C. (2000). The urban environmental gradient: anthropogenic influences on the spatial and
444 temporal distributions of lead and zinc in sediments. *Environmental Science & Technology*, 34(2), 232-238.

445 Cannon, W., & Horton, J.D. (2009). Soil geochemical signature of urbanization and industrialization—Chicago,
446 Illinois, USA. *Applied Geochemistry* 24, 1590-1601.

447 Carrero, J.A., Arrizabalaga, I., Bustamante, J., Goienaga, N., Arana, G., & Madariaga, J.M. (2013). Diagnosing the
448 traffic impact on roadside soils through a multianalytical data analysis of the concentration profiles of traffic-related
449 elements. *Science of the Total Environment*, 458–460, 427–434.

450 Charlesworth, S., De Miguel, E., & Ordóñez, A. (2011). A review of the distribution of particulate trace elements in
451 urban terrestrial environments and its application to considerations of risk. *Environmental geochemistry and health*
452 33, 103-123.

453 Chen, T.B., Zheng, Y.M., Lei, M., Huang, Z.C., Wu, H.T., Chen, H., Fan, K.K., Yu, K., Wu, X., & Tian, Q.Z.
454 (2005). Assessment of heavy metal pollution in surface soils of urban parks in Beijing, China. *Chemosphere* 60,
455 542-551.

456 Cheng, H., Li, M., Zhao, C., Li, K., Peng, M., Qin, A., & Cheng, X. (2014). Overview of trace metals in the urban
457 soil of 31 metropolises in China. *Journal of Geochemical Exploration* 139, 31-52.

458 Christoforidis, A., & Stamatis, N. (2009). Heavy metal contamination in street dust and roadside soil along the
459 major national road in Kavala's region, Greece. *Geoderma* 151, 257-263.

460 Cicchella, D., De Vivo, B., Lima, A., Albanese, S., McGill, R., & Parrish, R.R. (2008). Heavy metal pollution and
461 Pb isotopes in urban soils of Napoli, Italy. *Geochemistry: Exploration, Environment, Analysis* 8, 103-112.

462 Costa, C., & Jesus-Rydin, C. (2001). Site investigation on heavy metals contaminated ground in Estarreja—
463 Portugal. *Engineering Geology*, 60(1-4), 39-47.

464 Davis, A. P., Shokouhian, M., & Ni, S. (2001). Loading estimates of lead, copper, cadmium, and zinc in urban
465 runoff from specific sources. *Chemosphere*, 44(5), 997-1009.

466 Dean, J.R., Elom, N.I., & Entwistle JA (2017). Use of simulated epithelial lung fluid in assessing the human health
467 risk of Pb in urban street dust. *Science of the Total Environment*, 579, 387-395.

468 Dresel, W. (Ed.). (2007). *Lubricants and lubrication*. John Wiley & Sons.

469 Dung, T.T.T., Cappuyns, V., Swennen, R., & Phung, N.K. (2013). From geochemical background determination to
470 pollution assessment of heavy metals in sediments and soils. *Reviews in Environmental Science and*
471 *Bio/Technology* 12, 335-353.

472 Duzgoren-Aydin, N. S. (2007). Sources and characteristics of lead pollution in the urban environment of
473 Guangzhou. *Science of the Total Environment*, 385(1-3), 182-195.

474 Fernandez-Espinosa, A.J., & Ternero-Rodríguez, M. (2004). Study of traffic pollution by metals in Seville (Spain)
475 by physical and chemical speciation methods. *Analytical and bioanalytical chemistry*, 379(4), 684-699.

476 Fordyce, F.M.; & Ander, E.L. (2003). Urban soils geochemistry and GIS-aided interpretation: a case study from
477 Stoke-on-Trent. Nottingham, UK, British Geological Survey, 183pp

478 Francek, M. A. (1992). Soil lead levels in a small town environment: a case study from Mt Pleasant, Michigan.
479 *Environmental Pollution*, 76(3), 251-257.

480 Galušková, I., Mialjevič, M., & Strnad, L. (2014). Lead isotope composition and risk elements distribution in urban
481 soils of historically different cities Ostrava and Prague, the Czech republic. *Journal of Geochemical*
482 *Exploration*, 147, 215–221

483 Giusti, L. (2011). Heavy metals in urban soils of Bristol (UK). Initial screening for contaminated land. *Journal of*
484 *soils and sediments* 11, 1385-1398.

485 Guney, M., Bourges, C. M. J., Chapuis, R. P., & Zagury, G. J. (2017). Lung bioaccessibility of As, Cu, Fe, Mn, Ni,
486 Pb, and Zn in fine fraction (< 20 µm) from contaminated soils and mine tailings. *Science of the Total Environment*,
487 579, 378-386.

488 Hays, M.D, Cho, S.H., Baldauf, R., Schauer, J.J., & Shafer, M. (2011). Particle size distributions of metal and non-
489 metal elements in an urban near-highway environment. *Atmospheric Environment*, 45(4), 925-934.

490 Hu, Y., Liu, X., Bai, J., Shih, K., Zeng, E.Y., & Cheng, H. (2013). Assessing heavy metal pollution in the surface
491 soils of a region that had undergone three decades of intense industrialization and urbanization. *Environmental*
492 *Science and Pollution Research* 20, 6150-6159.

493 Jabarivasal, N. (2012). Indoor atmospheric radon in Hamadan, Iran. *Atmospheric radon indoors and around*
494 *Hamadan city in Iran* (Doctoral dissertation, University of Bradford).

495 Johnson, C.C., & Ander, E.L. (2008). Urban geochemical mapping studies: how and why we do them.
496 *Environmental Geochemistry and Health* 30, 511.

497 Kabata-Pendias, A. (2010). *Trace elements in soils and plants*. CRC press

498 Kabata-Pendias, A., & Mukherjee, A.B. (2007). *Trace elements from soil to human*. Springer Science & Business
499 *Media*

500 Khaleghian, A., & Modabberi, S. (2014). Arsenic contamination in soils and potatoes around Hamedan. 18th
501 *Symposium of the Geological Society of Iran*, Tarbiat Modares Univeristy, Tehran, Iran

502 Kisku, G., Barman, S., & Bhargava, S. (2000). Contamination of soil and plants with potentially toxic elements
503 irrigated with mixed industrial effluent and its impact on the environment. *Water, air, and soil pollution* 120, 121-
504 137.

505 Kuang, C., Neumann, T., Norra, S., & Stüben, D. (2004). Land use-related chemical composition of street sediments
506 in Beijing. *Environmental Science and Pollution Research*, 11(2), 73.

507 Lau, W.M., & Wong, H.M. (1982). An ecological survey of lead contents in roadside dusts and soils in Hong Kong.
508 *Environmental Research*, 28(1), 39-54.

509 Lee, C.S.-l., Li, X., Shi, W., Cheung, S.C.-n., & Thornton, I. (2006). Metal contamination in urban, suburban, and
510 country park soils of Hong Kong: a study based on GIS and multivariate statistics. *Science of the Total Environment*
511 356, 45-61.

512 Li, X., Wu, T., Bao, H., Liu, X., Xu, C., Zhao, Y., Liu, D., & Yu, H. (2017). Potential toxic trace element (PTE)
513 contamination in Baoji urban soil (NW China): Spatial distribution, mobility behavior, and health risk.
514 *Environmental Science and Pollution Research* 24, 19749-19766.

515 Lin, Z., Harsbo, K., Ahlgren, M., & Qvarfort, U. (1998). The source and fate of Pb in contaminated soils at the
516 urban area of Falun in central Sweden. *Science of the Total Environment*, 209(1), 47-58.

517 Liu, E., Yan, T., Birch, G., & Zhu, Y. (2014). Pollution and health risk of potentially toxic metals in urban road dust
518 in Nanjing, a mega-city of China. *Science of the Total Environment*, 476, 522-531.

519 Ljung, K., Selinus, O., & Otabbong, E. (2006). Metals in soils of children's urban environments in the small
520 northern European city of Uppsala. *Science of the Total Environment* 366, 749-759.

521 Lu, X., Wu, X., Wang, Y., Chen, H., Gao, P., & Fu, Y. (2014). Risk assessment of toxic metals in street dust from a
522 medium-sized industrial city of China. *Ecotoxicology and environmental safety*, 106, 154-163.

523 Maas, S., Scheifler, R., Benslama, M., Crini, N., Lucot, E., Brahmia, Z., Benyacoub, S., & Giraudoux, P. (2010).
524 Spatial distribution of heavy metal concentrations in urban, suburban and agricultural soils in a Mediterranean city
525 of Algeria. *Environmental pollution* 158, 2294-2301.

526 Manta, D.S., Angelone, M., Bellanca, A., Neri, R., & Sprovieri, M. (2002). Heavy metals in urban soils: a case study
527 from the city of Palermo (Sicily), Italy. *Science of the total environment* 300, 229-243.

528 MacKinnon, G., MacKenzie, A.B., Cook, G.T., Pulford, I.D., Duncan, H.J., & Scott, E.M. (2011). Spatial and
529 temporal variations in Pb concentrations and isotopic composition in road dust, farmland soil and vegetation in
530 proximity to roads since cessation of use of leaded petrol in the UK. *Science of the Total Environment*, 409, 5010–
531 5019.

532 McLaughlin, M.J., Hamon, R., McLaren, R., Speir, T., & Rogers, S. (2000). A bioavailability-based rationale for
533 controlling metal and metalloid contamination of agricultural land in Australia and New Zealand. *Soil Research* 38,
534 1037-1086.

535 Mielke, H.W., Gonzales, C.R., Smith, M.K., & Mielke, P.W. (1999). The urban environment and children's health:
536 soils as an integrator of lead, zinc, and cadmium in New Orleans, Louisiana, USA. *Environmental research*, 81(2),
537 117-129.

538 Micó, C., Recatalá, L., Peris, M., & Sánchez, J. (2006). Assessing heavy metal sources in agricultural soils of an
539 European Mediterranean area by multivariate analysis. *Chemosphere* 65, 863-872.

540 Mitchell, R.G., Spliethoff, H.M., Ribaud, L.N., Lopp, D.M., Shayler, H.A., Marquez-Bravo, L.G., ... & McBride,
541 M.B. (2014). Lead (Pb) and other metals in New York City community garden soils: factors influencing
542 contaminant distributions. *Environmental Pollution*, 187, 162-169.

543 Norra, S. (2009). The astysphere and urban geochemistry—a new approach to integrate urban systems into the
544 geoscientific concept of spheres and a challenging concept of modern geochemistry supporting the sustainable
545 development of planet earth. *Environmental Science and Pollution Research* 16, 539-545.

546 Odewande, A.A., & Abimbola, A.F. (2008). Contamination indices and heavy metal concentrations in urban soil of
547 Ibadan metropolis, southwestern Nigeria. *Environmental Geochemistry and Health* 30, 243-254.

548 Okorie, A., Entwistle, J., & Dean, J. R. (2012). Estimation of daily intake of potentially toxic elements from urban
549 street dust and the role of oral bioaccessibility testing. *Chemosphere*, 86(5), 460-467.

550 Othman, I., Al-Oudat, M., & Al-Masri, M.S. (1997). Lead levels in roadside soils and vegetation of Damascus city.
551 *Science of the Total Environment*, 207(1), 43-48.

552 Patinha, C., Reis, A. P., Dias, A. C., Abduljelil, A. A., Noack, Y., Robert, S., ... & Da Silva, E. F. (2015). The
553 mobility and human oral bioaccessibility of Zn and Pb in urban dusts of Estarreja (N Portugal). *Environmental*
554 *geochemistry and health*, 37(1), 115-131.

555 Peltola, P., & Åström, M. (2003). Urban geochemistry: a multimedia and multielement survey of a small town in
556 northern Europe. *Environmental Geochemistry and Health* 25, 397-419.

557 Pickett, S.T., Cadenasso, M.L., Grove, J.M., Boone, C.G., Groffman, P.M., Irwin, E., ... & Pouyat, R.V. (2011).
558 Urban ecological systems: Scientific foundations and a decade of progress. *Journal of Environmental Management*,
559 92(3), 331-362.

560 Pietrzak, U., & McPhail, D.C. (2004). Copper accumulation, distribution and fractionation in vineyard soils of
561 Victoria, Australia. *Geoderma*, 122(2-4), 151-166.

562 Poggio, L., Vrščaj, B., Schulín, R., Hepperle, E., & Marsan, F.A. (2009). Metals pollution and human
563 bioaccessibility of topsoils in Grugliasco (Italy). *Environmental Pollution* 157, 680-689.

564 Qingjie, G., Jun, D., Yunchuan, X., Qingfei, W., & Liqiang, Y. (2008). Calculating pollution indices by heavy
565 metals in ecological geochemistry assessment and a case study in parks of Beijing. *Journal of China University of*
566 *Geosciences* 19, 230-241.

567 Rate, A.W. (2018). Multielement geochemistry identifies the spatial pattern of soil and sediment contamination in an
568 urban parkland, Western Australia. *Science of The Total Environment*, 627, 1106-1120.

569 Reimann, C., & Caritat, P. (2000). Intrinsic flaws of element enrichment factors (EFs) in environmental
570 geochemistry. *Environmental Science & Technology* 34, 5084-5091.

571 Rodríguez-Seijo, A., Andrade, M. L., & Vega, F. A. (2017). Origin and spatial distribution of metals in urban soils.
572 *Journal of Soils and Sediments*, 17(5), 1514-1526.

573 Romic, M., & Romic, D. (2003). Heavy metals distribution in agricultural topsoils in urban area. *Environmental*
574 *geology* 43, 795-805.

575 Saeedi, M., Li, L.Y., & Salmazadeh, M. (2012). Heavy metals and polycyclic aromatic hydrocarbons: pollution and
576 ecological risk assessment in street dust of Tehran. *Journal of hazardous materials*, 227, 9-17.

577 Salonen, V.P., & Korkka-Niemi, K. (2007). Influence of parent sediments on the concentration of heavy metals in
578 urban and suburban soils in Turku, Finland. *Applied Geochemistry* 22, 906-918.

579 Sepahi, A.A. (2008). Typology and petrogenesis of granitic rocks in the Sanandaj-Sirjan metamorphic belt, Iran:
580 with emphasis on the Alvand plutonic complex. *Neues Jahrbuch für Geologie und Paläontologie-Abhandlungen* 247,
581 295-312.

582 Shi, G., Chen, Z., Xu, S., Zhang, J., Wang, L., Bi, C., & Teng, J. (2008). Potentially toxic metal contamination of
583 urban soils and roadside dust in Shanghai, China. *Environmental Pollution* 156, 251-260.

584 Shojaeimehr, S., & Zakerhaghighi, K. (2013). Explaining principles & characteristics of night spaces for the
585 purpose of strategic planning of leisure spaces within metropolises (The case study of Hamedan; Eram Area). *J.*
586 *Basic. Appl. Sci. Res.* 3(8), 747-756.

587 Sims, J., & Wolf, A. (1995). Recommended soil testing procedures for the Northeastern United States. *Northeast*
588 *Regional Bulletin* 493:4756 Sharpe.

589 Smolders, E., & Degryse, F. (2002). Fate and effect of zinc from tire debris in soil. *Environmental Science &*
590 *Technology*, 36(17), 3706-3710.

591 Solgi, E., Roohi, N., & Kouroshi-Gholampour, M. (2016). A comparative study of metals in roadside soils and
592 urban parks from Hamedan metropolis, Iran. *Environmental Nanotechnology, Monitoring & Management*, 6, 169-
593 175.

594 Sternbeck, J., Sjödin, Å., & Andréasson, K. (2002). Metal emissions from road traffic and the influence of
595 resuspension—results from two tunnel studies. *Atmospheric Environment* 36, 4735-4744.

596 Su, C. (2014). A review on heavy metal contamination in the soil worldwide: Situation, impact and remediation
597 techniques. *Environmental Skeptics and Critics* 3, 24.

598 Sutherland, R.A., & Tolosa, C.A. (2000). Multi-element analysis of road-deposited sediment in an urban drainage
599 basin, Honolulu, Hawaii. *Environmental pollution*, 110(3), 483-495.

600 Tang, R., Ma, K., Zhang, Y., & Mao, Q. (2013). The spatial characteristics and pollution levels of metals in urban
601 street dust of Beijing, China. *Applied geochemistry*, 35, 88-98.

602 Tume, P., Roca, N., Rubio, R., King, R., & Bech, J. (2018). An assessment of the potentially hazardous element
603 contamination in urban soils of Arica, Chile. *Journal of Geochemical Exploration*, 184, 345-357.

604 Turner, A., & Rice, L. (2010). Toxicity of tire wear particle leachate to the marine macroalga, *Ulva lactuca*.
605 *Environmental Pollution*, 158(12), 3650-3654.

606 Van Beers, D., & Graedel, T. (2007). Spatial characterisation of multi-level in-use copper and zinc stocks in
607 Australia. *Journal of Cleaner Production* 15, 849-861.

608 Van Bohemen, H., & Van De Laak, W.J. (2003). The influence of road infrastructure and traffic on soil, water, and
609 air quality. *Environmental management* 31, 0050-0068.

610 Wei, B., & Yang, L. (2010). A review of heavy metal contaminations in urban soils, urban road dusts and
611 agricultural soils from China. *Microchemical Journal* 94, 99-107.

612 Wong, C.S., Li, X., & Thornton, I. (2006). Urban environmental geochemistry of trace metals. *Environmental*
613 *Pollution* 142, 1-16.

614 Woszczyk, M., Spychalski, W., & Boluspaeva, L. (2018). Trace metal (Cd, Cu, Pb, Zn) fractionation in urban-
615 industrial soils of Ust-Kamenogorsk (Oskemen), Kazakhstan—implications for the assessment of environmental
616 quality. *Environmental monitoring and assessment*, 190(6), 362.

617 Wuana, R.A., & Okieimen, F.E. (2011). Heavy metals in contaminated soils: a review of sources, chemistry, risks
618 and best available strategies for remediation. *Isrn Ecology* 2011.

619 Xia, X., Chen, X., Liu, R., & Liu, H. (2011). Heavy metals in urban soils with various types of land use in Beijing,
620 China. *Journal of Hazardous Materials* 186, 2043-2050.

621 Yeganeh, M., Afyuni, M., Khoshgoftarmansh, A.H., Soffianian, A.R., & Schulin, R. (2012). Health risks of metals
622 in soil, water, and major food crops in Hamedan Province, Iran. *Human and Ecological Risk Assessment: An*
623 *International Journal*, 18(3), 547-568.

624 Yesilonis, I., Pouyat, R., & Neerchal, N. (2008). Spatial distribution of metals in soils in Baltimore, Maryland: role
625 of native parent material, proximity to major roads, housing age and screening guidelines. *Environmental Pollution*
626 156, 723-731.

627 Yongming, H., Peixuan, D., Junji, C., & Posmentier, E.S. (2006). Multivariate analysis of heavy metal
628 contamination in urban dusts of Xi'an, Central China. *Science of the total environment* 355, 176-186.

629 Yuan, G.L., Sun, T.H., Han, P., Li, J., & Lang, X.X. (2014). Source identification and ecological risk assessment of
630 heavy metals in topsoil using environmental geochemical mapping: typical urban renewal area in Beijing, China.
631 *Journal of Geochemical Exploration*. 136, 40-47.

632 Zechmeister, H.G., Hohenwallner, D., Riss, A., & Hanus-Illy, A. (2005). Estimation of element deposition derived
633 from road traffic sources by using mosses. *Environmental Pollution*, 138(2), 238-249.

634 Zhang, C. (2006). Using multivariate analyses and GIS to identify pollutants and their spatial patterns in urban soils
635 in Galway, Ireland. *Environmental pollution* 142, 501-511.

636 Zheng, Y-M., Chen, T.B., & He, J.Z. (2008). Multivariate geostatistical analysis of heavy metals in topsoils from
637 Beijing, China. *Journal of Soils and Sediments* 8, 51-58.

638

639 **Table Captions**

640 **Table 1.** PTEs concentrations (mg kg^{-1}) in surface soils of Hamedan city in comparison to Hamedan background
641 soils, uncontaminated soils and the average values in different cities of the world.

642 **Table 2.** The result of kruskal-wallis H test and mean values of PTEs (mg kg^{-1}) in 31 soil and street dust samples
643 from various land uses of Hamedan city

644 **Table 3.** Spearman's correlation matrix for PTEs in the surface soils and street dusts of Hamedan city

645 **Table 4.** The rotated component matrix of PTEs in urban soils from Hamedan

646 **Figure captions**

647 **Fig. 1.** Geological map of the Hamedan city in West of Iran and the sampling locations (Redrawn from Hamedan
648 1:100000 scale geological map, Geological Survey of Iran)

649 **Fig. 2.** Box plots of PTE enrichment factor (EF) in urban soils of Hamedan; the band near the middle of each box
650 represents the median. The bottom and top of the box are the first and third quartiles, respectively. Whiskers (the
651 vertical lines) are the 1.5 interquartile ranges of the lower and upper quartiles.

652 **Fig. 3.** Box plots of PTE contamination factor (CF) in urban soils of Hamedan; the band near the middle of each box
653 represents the median. The bottom and top of the box are the first and third quartiles, respectively. Whiskers (the
654 vertical lines) are the 1.5 interquartile ranges of the lower and upper quartiles.

655 **Fig. 4.** Principal component analysis results in the three-dimensional space: plot of loading of the first three factors

656 **Fig.5.** Dendrogram of the cluster analysis (CA) of the urban topsoils of Hamedan based on the PTEs concentrations

657 **Fig. 6.** Geochemical maps showing the spatial distribution of certain geogenic (As, Co, Ni) and anthropogenic (Cd,
658 Pb, Sb) elements in the urban soils of Hamedan city. The map legends are as per Fig. 1. The diameters of circles are
659 proportional to the concentration of given elements.

Table 1

		As	Cd	Co	Cr	Cu	Hg	Mo	Ni	Pb	Sb	V	Zn
Hamedan Urban soils	Min	11.1	0.01	9.1	42	22.7	0.01	0.48	36.7	17.7	0.57	39	67
	Max	27	0.35	17.1	76	79.3	1.34	1.15	70.6	289	5.25	79	322
	Mean	18.94	0.14	13.08	56.42	39.05	0.15	0.77	54.86	51.77	1.47	61.13	115.16
	Median	18.8	0.11	13.1	58	35.4	0.09	0.72	54	37.9	1.12	62	101
	SD	3.94	0.1	1.97	7.23	13.27	0.23	0.18	9.02	50.43	1.03	10.41	47.21
	Skewness	-.07	.53	-.02	.19	1.65	4.85	.613	.056	3.7	2.41	-.29	3.0
	Kurtosis	-.55	-.82	-.34	.70	2.98	25.46	-.454	-.881	16.8	6.12	-.39	12.0
Hamedan Background soils (mg kg ⁻¹)		21.9	0.075	18.2	72.5	36.5	0.012	0.7	62.4	13.7	0.39	105	90.6
¹ Uncontaminated soil		2.5	0.53	12	83	24	0.07	1.5	34	44	0.3	67	100
² Galway (Ireland)		8		6	35	27			22	58			85
³ Napoli (Italy)		11.9	0.37	6.3	11.2	74			8.9	141	2		158
⁴ Trondheim (Norway)		3.3	0.12		58	32			43	32			80
⁵ Annaba (Algeria)			0.3		28.3	23.8				42.3			64.7
⁶ Ibadan (Nigeria)		3	0.15		55.5	32			16.5	47			93.5
⁷ Murcia (Spain)					16.3	8.9			11.1	21.9			16.6
⁸ Baltimore (USA)			0.89		38.3	35.2			18.4	89.3			80.7
⁹ Chicago (USA)		13.2		11	65	59			31	198			235
¹⁰ Shanghai (China)			0.52		107.9	59.25			31.14	70.69			301.4
¹¹ Turku (Finland)			0.2		37	19.15			12.45	20			72.5

¹Kabata-Pendias (2010), ²Zhang (2006), ³Cicchella et al. (2008), ⁴Andersson et al. (2010), ⁵Maas et al. (2010), ⁶Odewande and Abimbola (2008), ⁷Acosta et al. (2011), ⁸Yesilonis et al. (2008), ⁹Cannon and Horton (2009), ¹⁰Shi et al. (2008), ¹¹Salonen and Korkka-Niemi (2007)

665
666

Table 2

Land use	As	Cd	Co	Cr	Cu	Hg	Mn	Mo	Ni	Pb	Sb	V	Zn
AG	21.3	0.1	14.1	62.0	43.1	0.1	663.7	0.7	63.1	49.0	1.6	63.8	107.4
GS	17.3	0.2	12.0	52.5	40.1	0.2	666.0	0.8	49.3	61.5	1.9	57.7	114.6
PK	17.1	0.2	13.5	57.0	38.5	0.1	719.3	0.9	55.0	58.2	1.1	60.8	170.0
PG	17.7	0.1	12.6	54.0	27.4	0.0	610.5	0.6	56.7	25.2	1.4	54.5	88.9
RS	20.8	0.1	13.8	58.0	37.8	0.1	723.8	0.8	55.8	44.0	1.0	65.6	101.0
chi-squared	6.42	6.62	6.19	6.62	3.32	2.54	4.65	7.45	9.77	2.71	3.51	3.00	4.62
df	4	4	4	4	4	4	4	4	4	4	4	4	4
p-value	0.17	0.16	0.19	0.16	0.51	0.64	0.32	0.11	0.04	0.61	0.48	0.56	0.33

Abbreviations: agricultural land (AG), green space (GS), park (PK), playground (PG) and residential (RS)

667
668

669
670

Table 3

	As	Cd	Co	Cr	Cu	Hg	Mn	Mo	Ni	Pb	Sb	V	Zn
As	1												
Cd	-0.257	1											
Co	.566**	0.088	1										
Cr	.473**	-0.035	.880**	1									
Cu	-0.211	0.3	-0.018	0.201	1								
Hg	-0.003	.386*	-0.016	0.102	.671**	1							
Mn	-0.076	-0.072	.388*	.397*	0.056	0.127	1						
Mo	-0.276	0.006	-.463**	-0.294	.361*	0.261	0.017	1					
Ni	.526**	0.198	.893**	.836**	0.112	0.061	0.115	-.484**	1				
Pb	-0.283	.466**	-0.184	-0.07	.825**	.649**	0.021	.433*	-0.069	1			
Sb	0.127	0.254	-0.016	-0.001	.426*	.531**	0.046	-0.058	0.058	.426*	1		
V	.492**	-0.118	.717**	.766**	-0.078	0.095	.468**	-0.209	.532**	-0.331	-0.105	1	
Zn	-.450*	.652**	-0.127	-0.087	.724**	.475**	-0.073	0.324	-0.026	.825**	.382*	-0.34	1

******. Correlation is significant at the 0.01 level (2-tailed).

*****. Correlation is significant at the 0.05 level (2-tailed).

Table 4

Elements	Rotated Component Matrix					
	F1	F2	F3	F4	F5	F6
As	0.635	-0.042	-0.359	-0.306	-0.009	0.245
Ba	-0.234	-0.075	0.712	-0.067	-0.272	-0.171
Cd	-0.065	-0.068	-0.093	0.899	0.057	-0.087
Co	0.862	0.327	-0.143	0.044	-0.109	-0.234
Cr	0.814	0.413	0.194	-0.005	0.073	-0.079
Cu	0.042	0.013	0.515	0.169	0.702	0.113
Hg	-0.502	0.184	-0.314	0.468	0.309	0.066
Mn	0.128	0.884	0.087	-0.028	0.023	-0.172
Mo	-0.326	0.032	0.503	0.215	0.058	0.648
Ni	0.895	0.063	-0.132	0.196	-0.003	-0.238
Pb	-0.318	-0.013	0.831	0.117	0.171	0.065
Sb	-0.086	0.057	-0.107	-0.007	0.772	-0.409
V	0.664	0.627	-0.225	-0.115	-0.046	0.123
Zn	0.058	-0.089	0.34	0.817	0.119	0.051
Al	0.844	0.298	-0.221	-0.208	0.008	-0.156
Fe	0.698	0.489	-0.175	-0.049	-0.179	-0.312
K	0.545	0.447	-0.271	-0.181	0.309	0.376
Mg	0.783	0.051	-0.276	-0.015	0.053	0.055
Na	-0.154	-0.025	-0.147	-0.118	0.022	0.831
P	0.008	-0.055	-0.116	0.166	0.77	0.395
Ti	0.429	0.804	-0.088	-0.04	0.022	0.205
Eigenvalue	7.811	2.680	2.241	1.897	1.340	1.263
% variance explained	28.172	12.836	11.834	9.793	9.776	9.644
Cumulative % variance	28.172	41.009	52.843	62.636	72.411	82.055

Extraction Method: Principal Component Analysis.

Rotation Method: Varimax with Kaiser Normalization.

Rotation converged in 9 iterations.

697 **Figure 2.**

698

699

700 **Figure 3.**

701

702

703

Figure 4.

704

705

706 **Figure 5.**

707

708 **Figure 6.**

709

710

711 **Supplementary Material**
712 The results of Sixty-three elements analyzed in urban topsoils of Hamedan
713

	Unit	DI	S1	S2	S3	S4	S5	S6	S7	S8	S9	S10	S11	S12	S13	S14	S15	S16	S17	S18	S19	S20	S21	S22	S23	S24	S25	S26	S27	S28	S29	S30	S31	S32	S33	S34
Ti	%	0.001	0.197	0.162	0.123	0.141	0.09	0.088	0.083	0.068	0.122	0.125	0.141	0.1	0.111	0.171	0.113	0.019	0.083	0.074	0.118	0.134	0.071	0.111	0.129	0.179	0.17	0.133	0.129	0.142	0.143	0.134	0.18	0.22	0.331	0.166
S	%	1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
P	%	0.001	0.114	0.215	0.213	0.329	0.18	0.119	0.129	0.12	0.177	0.118	0.176	0.155	0.248	0.135	0.168	0.086	0.236	0.273	0.232	0.126	0.147	0.135	0.19	0.105	0.115	0.273	0.319	0.176	0.161	0.277	0.132	0.109	0.076	0.289
Li	ppm	0.1	45.7	43.6	38.4	39.8	43.9	38.8	35.5	32.7	41.5	41.4	37.8	31.2	36.2	52.4	35.6	40.4	27.5	28.7	40.5	41.3	24.1	30.7	37.6	55.5	50.1	39.6	36.7	41	34.2	35.9	41.6	44.6	55.8	44.8
Be	ppm	0.1	1.1	1.1	1.1	1.2	1	1.2	1	0.9	1.1	1	1	0.8	1.2	1.2	1	0.9	0.7	0.8	1.3	1	0.9	0.9	0.8	1.3	1.1	1.2	1	1.1	1	0.9	1.6	1	1	1.1
B	ppm	1	16	19	22	20	22	21	23	19	20	14	21	24	27	14	25	10	16	27	18	22	15	16	20	14	14	20	22	22	18	22	10	14	7	21
Na	%	0.001	0.059	0.052	0.049	0.078	0.062	0.058	0.064	0.066	0.084	0.071	0.069	0.137	0.135	0.065	0.066	0.086	0.094	0.092	0.067	0.094	0.078	0.078	0.166	0.07	0.142	0.363	0.128	0.106	0.082	0.075	0.096	0.055	0.057	0.077
Mg	%	0.01	0.94	0.86	0.93	0.93	0.95	0.99	0.83	0.69	0.91	0.68	0.9	0.83	0.92	0.86	0.91	0.71	0.65	0.75	0.77	0.84	0.59	0.77	0.93	0.85	0.85	0.78	0.91	0.87	0.85	0.81	0.72	1.11	1.09	1.1
Al	%	0.01	2.75	2.51	2.4	2.48	2.36	2.49	1.97	1.73	2.25	1.91	2.37	1.73	2.08	2.46	2.14	2.33	1.46	1.55	2.09	1.88	1.41	1.91	2.2	2.29	2.25	2	2.21	2.18	2.08	2.16	2.4	2.92	3.08	2.87
K	%	0.01	1	1.07	0.87	1.04	0.78	0.69	0.65	0.61	0.83	0.77	0.95	0.64	0.89	1.07	0.91	0.58	0.56	0.7	0.85	0.73	0.52	0.69	0.82	0.93	0.87	1.09	1.03	1.16	0.76	0.88	0.77	1.14	1.42	1.23
Bi	ppm	0.02	0.29	0.35	0.34	0.33	0.29	0.24	0.29	0.19	0.28	0.28	0.45	0.23	0.27	0.28	0.24	0.21	0.22	0.26	0.28	0.41	0.23	0.28	0.25	0.25	0.25	0.21	0.25	0.28	0.28	0.42	0.13	0.31	0.39	0.39
Ca	%	0.01	2.41	1.65	5.81	2.59	4.63	6.12	5.53	6.68	5.9	2.27	2.26	4.76	5.7	2.1	4.15	2.92	4.19	4.31	3.24	5.67	4.89	5.35	5.2	4.31	3.04	4.07	4.99	4.32	3.21	3.84	0.91	1.2	0.32	3.69
Sc	ppm	0.1	10.5	9.6	9.2	9.6	7.1	7.5	6.2	5.3	8	7.2	8.5	6.3	7.1	10.9	8	4.9	5.5	4.3	7.4	8	4.7	6.9	7.7	10	9.6	7.9	8.7	8.6	8.1	8.1	9.4	10.3	13.9	10.4
V	ppm	1	79	75	66	67	57	59	53	48	64	61	70	50	57	72	64	45	43	45	57	62	39	56	61	77	69	66	62	67	66	60	71	82	128	78
Cr	ppm	1	67	60	64	63	59	61	50	44	59	48	65	48	58	61	58	50	42	45	54	51	55	53	57	61	55	52	58	60	57	58	50	73	72	76
Mn	ppm	1	883	778	714	765	672	538	625	539	688	603	653	753	691	798	628	593	717	563	662	548	694	699	705	836	789	551	710	681	780	650	661	817	849	690
Fe	%	0.01	4.2	3.82	3.57	3.42	3.54	3.29	3.41	2.81	3.59	3.22	3.66	3.11	3.39	3.77	3.25	3.56	2.84	2.66	3.16	2.98	2.64	3.28	3.37	3.57	3.47	2.91	3.19	3.34	3.3	3.03	3.88	4.12	4.36	3.78
Co	ppm	0.1	17.1	16.2	14.8	13.9	16	14.6	12.8	10.7	14.6	12	14.8	12.6	13.6	14	13.6	11.5	9.5	10.5	11.1	11.6	9.1	12.1	13.6	13.7	12.8	10.6	12.8	13.8	13.1	12.3	14.1	17.7	18.7	16.1
Ni	ppm	0.1	67.3	62.2	68.5	60.6	64.7	69.3	54.8	46.1	60	43.8	67.6	44.2	58.5	54	60.3	53.1	44	47	45.1	45.3	36.7	52.5	59.4	53.7	49.4	44.3	54	59.2	49.6	55	36.9	70.8	54	70.6
Cu	ppm	0.01	31.6	34.8	34.4	40.3	40	29.5	29.4	22.7	35.4	25	46.8	33	40.7	32.4	30.8	24	35.5	38.8	75.5	33.3	56	55.4	52	27.8	27.7	36.4	43.8	39.2	33.3	79.3	25.2	30.2	42.8	45.9
Zn	ppm	0.1	93.4	101	88.3	103	111	87.6	95.3	88.2	106	89	322	134	129	83.5	97.2	80.6	165	170	120	111	146	167	101	67	76.8	69.9	100	102	95.1	134	88.9	85.6	95.6	136
Ga	ppm	0.02	11.1	10.1	9.59	9.93	9.05	8.99	7.66	6.34	8.98	7.98	9.85	6.91	8.05	9.95	8.9	8.01	6.21	6.3	8.62	7.95	5.39	7.53	8.38	9.59	9.4	8.46	8.73	8.88	8.47	8.98	13.5	11.3	13.8	11.4
Ge	ppm	0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1
As	ppm	0.1	20.7	21.2	21.5	24.9	23.2	24.5	24.2	18.1	22.8	23.5	21.4	13.9	15.4	18.8	20.9	14.4	12.8	12.4	15.8	15.1	11.1	17.1	19.4	20.7	17.9	27	18.5	19.7	17.5	16.5	7.8	22.1	21.7	18
Rb	ppm	0.1	68.1	67.7	56.1	62.5	49	45.7	39.4	38.8	52.1	75.9	62	45.6	50.4	72.9	60	26.6	34.2	35.5	62.5	66.2	32.6	41.6	47.2	71.1	65.7	65.9	54	58.2	52.9	57.6	62.2	73	116	74.8
Sr	ppm	0.5	58.1	57	109	80.8	134	175	184	158	135	58.3	75.1	164	156	75.3	117	82.7	141	155	108	145	124	138	158	117	96.5	193	159	124	99.7	112	43.9	35.9	23	121
Y	ppm	0.01	11.9	10.1	9.02	10.2	12.1	10.7	10.2	10.2	11.1	9.35	8.49	12	8.09	10.9	7.79	6.52	10.2	7.2	10.6	7.63	8.98	9.27	10.7	8.52	10.3	7.43	9.26	9.11	10.1	8.48	14.8	9.61	8.13	9.14
Zr	ppm	0.1	2.4	1.4	1.4	1.2	1	1.2	0.8	0.9	1.2	0.9	1.5	0.9	1.1	1.7	1.2	1.7	0.6	1.1	0.7	1.1	0.7	1.1	1.7	2	2	1.3	1.8	1.1	1.5	1.1	2	2.4	2.3	1.4
Nb	ppm	0.1	3.1	2.9	2.5	2.6	1.9	1.7	2.2	1.8	1.9	3	3.6	2.9	3.3	2.2	2.4	0.8	2.5	2.7	3.1	3.9	2.1	1.9	2.1	2.1	2.1	3.2	2.7	3.2	3.4	2.8	2.9	2.8	2.8	3.3
Mo	ppm	0.01	0.7	0.55	0.64	0.57	0.58	0.58	0.66	0.64	0.74	0.87	0.97	0.88	0.86	0.48	0.63	0.55	0.82	0.87	0.67	0.68	1.15	1.12	0.72	0.9	0.76	1.12	0.89	0.71	0.81	1.06	0.48	0.93	0.48	0.58
Ag	ppm	0.002	0.284	0.397	0.484	0.781	0.43	0.289	0.338	0.234	0.651	0.392	0.581	0.323	0.424	0.418	0.46	0.177	1.04	0.543	0.729	0.409	0.328	0.545	0.514	0.282	0.259	0.607	0.749	0.554	0.62	0.892	0.236	0.247	0.144	0.724
In	ppm	0.02	0.05	0.04	0.04	0.04	0.02	0.03	0.03	0.03	0.03	0.04	0.03	0.03	0.04	0.03	0.03	0.03	0.03	0.02	0.03	0.03	0.02	0.03	0.03	0.04	0.04	0.03	0.03	0.03	0.04	0.04	0.05	0.04	0.07	0.04
Sn	ppm	0.05	3.51	5.58	4.87	4.97	16.5	3.18	3.16	2.33	5.29	3.6	7.99	4.18	5.68	4.33	3.29	1.49	4.98	5.58	5.88	5.9	>200	9.46	5.25	3.56	3.65	6.24	6.37	5.21	4.45	17.6	3.43	2.54	3.62	6.47
Sb	ppm	0.02	0.79	3.62	3.79	1.7	1.27	0.65	1.36	1	1.04	1.37	1.37	0.86	1.01	1.03	2.11	0.73	2.39	1.45	5.25	0.57	1.51	1.36	1.62	0.77	0.63	1.12	0.95	1.05	1.08	1	0.19	0.52	0.26	1.15
			<	<		<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<
Te	ppm	0.02	0.02	0.02		0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.02
Cs	ppm	0.02	7.11	8.06	5.59	5.72	5.58	4.55	4.17	4.07	5.37	14.3	6.99	4.08	4.72	6.81	6.14	1.82	3.45	3.15	5.81	6.35	3.02	4.16	4.97	6.41	5.62	12.4	5.05	5.78	6.16	5.43	7.68	7.25	11.2	7.2
Ba	ppm	0.5	127	117	131	114	123	120	105	202	121	103	121	139	128	102	122	115	114	107	99.2	109	219	139	114	118	150	101	114	109	10					

Tb	ppm	0.1	0.5	0.4	0.4	0.4	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.4	0.3	0.3	0.3	0.3	0.4	0.3	0.3	0.3	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.3	0.6	0.5	0.5	0.4	
Dy	ppm	0.1	2.3	2	1.7	2	2.2	1.9	1.8	1.8	2	1.9	1.6	2	1.5	2	1.5	1.3	1.6	1.2	1.9	1.4	1.5	1.6	1.9	1.6	1.9	1.4	1.7	1.7	1.8	1.6	2.8	2	2	1.8	
Ho	ppm	0.1	0.5	0.4	0.4	0.4	0.5	0.4	0.4	0.4	0.5	0.4	0.4	0.5	0.3	0.5	0.3	0.3	0.4	0.3	0.4	0.3	0.3	0.4	0.4	0.4	0.4	0.3	0.4	0.4	0.4	0.3	0.6	0.4	0.4	0.4	
Er	ppm	0.1	1.2	1	0.9	1.1	1.3	1.1	1.1	1.1	1.2	1	0.9	1.3	0.8	1.1	0.8	0.7	1	0.7	1.1	0.8	1	1	1.2	0.9	1.1	0.8	1	0.9	1	0.8	1.4	1	0.8	0.9	
Tm	ppm	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.1	<0.1	0.1	<0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	0.1	<0.1	0.1	
Yb	ppm	0.1	1	0.8	0.8	0.9	1	0.9	0.9	0.9	1	0.8	0.7	1.2	0.7	0.9	0.7	0.5	0.9	0.6	1	0.6	0.9	0.9	1	0.8	1	0.7	0.8	0.8	0.9	0.7	1.1	0.8	0.5	0.7	
Lu	ppm	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2	<0.1	0.1	0.1	<0.1	0.1	<0.1	0.1	<0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	<0.1	0.1	0.1	<0.1	0.1	
Hf	ppm	0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	<0.1	
Ta	ppm	0.05	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	
W	ppm	0.1	0.2	0.3	0.2	0.5	0.2	0.1	0.2	0.2	0.2	0.3	0.3	12.9	0.4	0.3	0.2	<0.1	0.3	0.3	0.3	0.4	0.4	0.7	0.4	0.4	0.4	0.4	0.3	0.3	0.7	0.3	0.1	0.4	0.2	<	
Re	ppm	0.001	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	<	
Au	ppb	0.5	<0.5	7	1.7	9.2	22	<0.5	<0.5	<0.5	<0.5	3.9	19.9	4.6	13.9	2.5	7.7	<0.5	203	10	17.3	9.5	8.7	17	16.3	<0.5	<0.5	9.8	24.7	8.1	5.8	127	<0.5	<0.5	<0.5	30.8	
Tl	ppm	0.02	0.39	0.39	0.31	0.33	0.29	0.29	0.24	0.22	0.27	0.33	0.35	0.22	0.29	0.4	0.33	0.15	0.19	0.21	0.34	0.32	0.18	0.24	0.26	0.38	0.35	0.27	0.27	0.33	0.28	0.28	0.29	0.45	0.6	0.37	
Pb	ppm	0.01	23.5	29.7	30.4	80.2	56.5	22	22.1	25.2	37.5	24.5	58.5	103	42.9	21.1	32.5	17.9	75.2	43.1	40.5	29.9	289	107	62.1	17.7	21.1	37.9	47	39.6	28.2	89.1	10.7	14.7	12.7	50.1	
Th	ppm	0.1	11.3	10	6.5	8.8	8	7	7	6	8	9.3	5.8	4.7	4.5	11	5.4	7.8	4.5	1.9	7.9	5.5	3.3	6.5	7.4	7.3	9.9	6	7.4	7	7.2	7.3	16	11.4	15	8.2	
U	ppm	0.1	1.4	1.6	1	1.4	1	1.1	1.3	1	1.1	1.3	1.2	1.2	1.1	1.2	1	0.8	1	0.9	1.3	1.1	0.9	1	1.1	1	1.2	1.1	1.1	1.1	1.1	1.1	1.1	1.6	1.4	1.9	1.4
Hg	ppb	10	70	150	150	220	90	20	30	50	80	70	150	100	90	60	80	10	1340	150	150	70	70	80	110	60	50	310	250	110	90	200	20	20	<10	140	