UWS Academic Portal

Assessing the impacts of the introduced channel catfish *Ictalurus punctatus* using the comparative functional response approach

Faria, Larissa; Alexander, Mhairi E.; Vitule, Jean R.S.

Published in:
Fisheries Management and Ecology

DOI:
[10.1111/fme.12353](https://doi.org/10.1111/fme.12353)

Published: 16/11/2019

Document Version
Peer reviewed version

[Link to publication on the UWS Academic Portal](#)

Citation for published version (APA):

Faria, L., Alexander, M. E., & Vitule, J. R. S. (2019). Assessing the impacts of the introduced channel catfish *Ictalurus punctatus* using the comparative functional response approach. *Fisheries Management and Ecology*, 26(6), 570-577. <https://doi.org/10.1111/fme.12353>

General rights

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact pure@uws.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

"This is the peer reviewed version of the following article: Faria, L., Alexander, M. E., & Vitule, J. R. S. (2019). Assessing the impacts of the introduced channel catfish *Ictalurus punctatus* using the comparative functional response approach. *Fisheries Management and Ecology*, which has been published in final form at <https://doi.org/10.1111/fme.12353>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions."

<https://authorservices.wiley.com/author-resources/Journal-Authors/licensing/self-archiving.html>

Assessing the impacts of the introduced channel catfish *Ictalurus punctatus* using the Comparative Functional Response approach

Abstract

In the metric “Relative Impact Potential” (RIP), the functional response (FR) of a non-native species can be compared to that of a native analogue and combined with the species abundance to predict its environmental impact. Here, using the River Guaraguaçu (Brazil) as a case study, this methodology was implemented to identify the impacts of the non-native channel catfish *Ictalurus punctatus* compared to a native species *Rhamdia quelen* towards small prey fish. Both species exhibited Type II FRs, however handling times were lower for *I. punctatus*, resulting in a greater maximum feeding rate in this species. Consequently, an $RIP > 1$ was found, indicating that *I. punctatus* represents a superior impact to prey when compared to its native analogue. These results demonstrate that *I. punctatus* is a potential threat to small endangered fish species. Therefore, policies to avoid escapes from aquaculture should be created and the abundance of *I. punctatus* controlled.

Keywords

alien species, aquaculture, fish, freshwater, management, Relative Impact Potential

1 Introduction

The introduction of non-native species to novel regions is one of the main drivers of species extinction and is considered the second major cause of global biodiversity loss after habitat destruction (Bellard, Cassey & Blackburn, 2016; Clavero & García-Berthou, 2005; Vitousek, D'Antonio, Loope & Westbrooks, 1996). Predation by non-native species can devastate native prey populations as non-native predators are often more efficient consumers of resources compared to their native counterparts (Paolucci, MacIsaac & Ricciardi, 2013; Salo, Korpimäki, Banks, Nordström & Dickman, 2007; Simberloff & Vitule, 2014). Compounding this, native prey often do not exhibit effective anti-predator behaviour due to a lack of shared evolutionary history with the new predator (Cox & Lima, 2006). These mechanisms can explain the establishment success of non-native predators and their negative impact in the recipient community (Gallardo, Clavero, Sánchez & Vilà, 2016; Latini & Petreere, 2004; Sharpe, De León, González & Torchin, 2017; Simberloff et al., 2013; Trumpickas, Mandrak & Ricciardi, 2011).

Prediction and quantification of impacts of non-native species is of great importance for conservation of biodiversity, although there is no agreed standard methodology for doing so (Kumschick et al., 2015; Parker et al., 1999; Ricciardi, Hoopes, Marchetti & Lockwood, 2013). The comparative functional response (CFR) has been proposed as an approach that “can unify invasion ecology” (Dick et al., 2017a), providing a trait-based methodology for measuring resource-use interactions of non-native species as a means of predicting their potential ecological impact. The CFR approach is based on the comparison of the functional response (FR) of non-native and trophically analogous native species, where the difference in the magnitude of the FR can predict and quantify the relative impacts of the non-native species (Dick et al., 2014).

A FR is defined as the relationship between consumption rate and resource density (Holling, 1959b). There are three types of FR: Type I – an increasing linear relationship between resource density and consumption until a threshold of satiation; Type II – a decelerating rate of consumption that reaches an asymptote at higher densities; and Type III – an S-shaped curve where the consumption rate first accelerates at low densities and then decelerates towards satiation (Holling, 1959b). The FR type of the predator can have direct effects on the stability of the prey population, for instance Type II FR has destabilising effects with high consumption at low densities, whereas Type III has stabilising effects with a refuge of predation at low densities (Oaten & Murdoch, 1975; Sinclair et al., 1998). As FRs measure the *per capita* effect of a non-native species, it can be combined with the abundance of the species in the field, in a metric called the Relative Impact Potential (RIP), to enhance the predictive power of this approach (Dick et al., 2017b).

A native to North America, the channel catfish (*Ictalurus punctatus* Rafinesque, 1818) has been introduced worldwide for aquaculture and sport fishing purposes (CABI, 2019). The species was first introduced in Brazil in 1971 and, having potentially escaped accidentally from aquaculture cages (Orsi & Agostinho, 1999; Welcomme, 1988), it is now widely found in Brazilian water bodies (Cruz-Spindler et al., 2012; Ota, Message, Graça & Pavanelli, 2015; Zanatta et al., 2010). As a non-native species, *I. punctatus* is a potential threat to local fish assemblages through competition and predation of native species (e.g. Matsuzaki et al., 2011; Townsend & Winterbourn, 1992; Troca & Vieira, 2012). For example, the decline and extinction of species, such as the razorback sucker *Xyrauchen texanus*, in communities invaded by *I. punctatus* has been reported (CABI, 2019). Additionally, *I. punctatus* demonstrates high rates of dispersal, which is aided by its ability to adapt to a wide range of habitats (Olden & Poff, 2005).

The aim of the present study was to apply the CFR approach and the RIP metric to assess the potential impacts of the invasive *I. punctatus* in a highly-diverse river in Brazil, the River Guaraguaçu. The abundance of the species is not constantly monitored, but there are reports about the establishment of this species in the river basin, as well as, in other basins of the Paraná State (Vitule, Umbria & Aranha, 2005; Vitule, 2008). Based on the species traits and invasion history of *I. punctatus* (Gutierre, Vitule, Freire & Prodocimo, 2014; Tatarenkov, Barreto, Winkelman & Avise, 2006; Townsend & Winterbourn, 1992; Wellborn, 1988), and considering its taxonomic distinctiveness within the recipient community (Ricciardi & Atkinson, 2004), the likely impact of *I. punctatus* on native species and ecosystems is through an elevated consumption of small stream fishes.

To address the paucity of research on non-native species impacts in South American countries (Bellard & Jeschke, 2016; Speziale, Lambertucci, Carrete & Tella, 2012; Vitule, Freire, Vazquez, Nuñez & Simberloff, 2012), the present study aims to enhance the understanding of the impacts of introduced predators in the Neotropical region, where biodiversity is both rich and threatened (Lowry et al., 2013; Myers, Mittermeier, Mittermeier, da Fonseca & Kent, 2000).

2 Methods

2.1 Species used

Ictalurus punctatus is a nocturnal forager that uses a variety of habitats with preference for shallow warm waters (Bayley & Harrison, 1948; Braun & Phelps, 2016; Jordan, Neumann & Schultz, 2004), detecting food using olfactory and visual cues (Becker, 1983). Young *I. punctatus* feed mainly on insects, but adults are usually omnivorous and prey also on fish (Bailey & Harrison, 1948; Braun & Phelps, 2016; Hill, Duffy &

Thompson, 1995; Tyus & Nikirk, 1990). The species has opportunistic behaviour and is generalist, preying on a wide range of available resources in non-native regions (Endo et al., 2015; Haubrock et al., 2018).

The FRs of the non-native channel catfish *I. punctatus* were compared to a native consumer, the South American silver catfish *Rhamdia quelen* (Quoy & Gaimard, 1824), which is a benthic species widely distributed in lakes, reservoirs and rivers of Central and South America (Gomes, Golombieski, Gomes & Baldisserotto, 2000; Silfvergrip, 1996). There are few studies on the biology of *R. quelen*, which like *I. punctatus* is representative of the Siluriformes and is generally described as nocturnal and omnivorous when mature (Bailey & Harrison, 1948; Gomes et al., 2000; Goudie, Davis & Simco, 1983). Adults of *R. quelen* are considered generalists, feeding on fish, crustaceans, insects, vegetation and organic detritus (Guedes, 1980; Meurer & Zaniboni Filho, 1997). *Ictalurus punctatus* and *R. quelen* have many similarities in their anatomic digestive systems and diet (Piedras, Pouey & Moraes, 2006) which make *R. quelen* an appropriate resident comparator for *I. punctatus* in the FR experiments described here.

Both species also spawn in the spring, with temperature an important factor in reproduction (Becker, 1983; Gomes et al., 2000). *Rhamdia quelen* has low fecundity rates and do not present parental care (Gomes et al., 2000). Females of *I. punctatus* have around 8,000 eggs per kg body weight and males guard the eggs during a period of incubation of 5 to 10 days (Becker, 1983). *Ictalurus punctatus* reaches sexual maturity at 300–375 mm in total length (TL), whereas *R. quelen* generally matures at 134–175 mm TL (Gomes et al., 2000; Jackson, 2004; Shephard & Jackson, 2005).

Blue tetra *Mimagoniates microlepis* (Steindachner, 1877), a congeneric of the locally and nationally threatened *M. lateralis* (Abilhoa & Duboc, 2004; Rosa & Lima, 2008; MMA, 2014), is abundant in natural water bodies and was used as prey in

experimental trials. Both *Mimagoniates* species are sympatric and are representatives of Glandulocaudinae, which is characterised by small stream fishes (Braga, Braga, & Vitule, 2013; Menezes & Weitzman, 2009). *Mimagoniates lateralis* is the smallest species of the genus, reaching maximum 40 mm standard length (Menezes & Weitzman, 2009; Rosa & Lima, 2008).

2.2 Experimental design

For the experimental protocol (Figure S1), which was approved by the Ethics Commission on the Use of Animals of the Biological Sciences Department of the Federal University of Paraná (CEUA/BIO UFPR – Certificate nº1027), juveniles of predator fish species (*I. punctatus* and *R. quelen*) were obtained in local aquaculture farms and stored in tanks of 300 L in the lab. Each species was kept in a different tank with constant aeration and filtration for acclimation during 30 days. Prey fish were collected in the field using fishing nets and kept in an 80 L aquarium, with constant aeration and filtration, and were also acclimated for 30 days. The temperature in the lab was kept at 22–26°C under a natural light regime. During the acclimation period predators were fed every 48 h with sausage to standardise prior experience.

Experiments were performed in 10 L opaque plastic boxes (35 × 20 × 25 cm) with constant aeration. Individuals of predator species were randomly selected one day prior to use and placed in the experimental aquariums for 24 h acclimation. Predators ($n = 9$ per species) were size-matched with respect to total length (*R. quelen*: 16.39 cm $\pm$ 1.15 and *I. punctatus*: 16.83 cm $\pm$ 2.14; Students' *t*-test, $t = -0.55$, $df = 16$, $p = 0.59$). Each predator specimen was reused in two more trials with different initial densities. After the acclimation period, they were held without food for 72 h to standardise hunger levels. Predators were then presented with the prey at five initial densities (2, 5, 10, 20, and 30 individuals), with five replicates per density. Prey were also size matched with respect to

total length by visual selection (≈ 3 cm). Experiments were initiated at 14:00 and prey consumption was recorded after 24 h. Controls were three replicates of each initial density of prey in the absence of predators.

2.3 Data analysis

The FR type was determined from a logistic regression of prey density by the proportion of consumed prey, as proposed by Juliano (2001). A negative first order term indicates a Type II FR, whereas a positive first order term followed by a negative second order term indicates a Type III FR (Pritchard, Paterson, Bovy & Barrios-O'Neill, 2017). The FR curves and the parameters attack rate (a) and handling time (h) were modelled using Maximum Likelihood Estimation (Bolker, 2008) with the Rogers' (1972) random predator equation that considers the depletion of prey without replacement along the experimental period:

$$N_e = N_0 \{1 - \exp [a (N_e h - T)]\} \quad (1)$$

where N_e is the number of prey consumed, N_0 is the initial density of prey, a is the attack rate, h is the handling time and T is the experimental period, given in days. The attack rate (a) is the instantaneous capture rate, *i.e.* the rate at which the consumer encounters resource items per unit of resource density; and the handling time (h) is the time spent capturing and consuming the prey item (Brose, 2010; Holling, 1959a; Jeschke, Kopp & Tollrian, 2002). The h was used to estimate the maximum feeding rate ($1/hT$), which represents the FR curve asymptote. To compare the FR of each species, the parameters a and h were compared between predators using the indicator variables method, as the following equation:

$$0 = N_0 - N_0 \exp \{[a + Da(j)] [h + Dh(j)] (N_e) - T\} - N_e \quad (2)$$

where j is an indicator variable that takes value 0 for *R. quelen* and 1 for *I. punctatus*. The parameters Da and Dh estimate the differences between the predators in the value of the parameters a and h , respectively. If these parameters are significantly different from zero, then the two species differ significantly in the corresponding parameters (Juliano, 2001). To visualise the uncertainty around the fitted functional responses, 95% confidence intervals were constructed by bootstrapping ($n = 2000$) data around FR curves. If confidence intervals do not overlap, then it can be stated that species FRs differ (Paterson et al., 2015). Analyses were carried in R v. 3.4.1 (R Core Team, 2015) using the ‘frair’ package (Pritchard et al., 2017) at 0.05 significance.

The Relative Impact Potential (RIP) metric was calculated using the following equation, according to proposed by Dick et al., 2017b:

$$RIP = \frac{FR_{\text{non-native}}}{FR_{\text{native}}} \times \frac{AB_{\text{non-native}}}{AB_{\text{native}}} \quad (3)$$

where FR is the estimated maximum feeding rate ($1/hT$) and AB is the field abundance/biomass of the species. When $RIP < 1$, the non-native species is predicted to have less impact than the native equivalent; when $RIP = 1$, there is no impact above that driven by native equivalents; whereas $RIP > 1$ indicates a likely non-native ecological impact (Dick et al., 2017b). Data of total abundance and biomass were taken from literature, from samples collected in River Guaraguaçu using a variety of fishing gears (Vitule, 2008) (Table S1). As data from different methods of capture and of different fishing efforts were compiled (Table S1), the capture per unit effort (CPUE) was calculated to standardise these differences. This was the number of fishes captured divided by the number of fish hooks or area of gill nets, multiplied by fishing effort in hours.

3 Results

None of the prey fish died in any of the control replicates, therefore prey mortality in the experimental trials was attributed to predation. Both species obtained a negative first-order term from logistic regressions, indicating a Type II FR (Table 1, Figure 1). Parameter estimates of the FR model were all statistically significant (Table 1). The attack rate did not differ between species ($Da = -0.646$, $z = -0.447$, $p = 0.655$), but the handling time was significantly lower for *I. punctatus* ($Dh = 0.053$, $z = 2.209$, $p < 0.05$), and thus a higher maximum feeding rate was obtained (Table 1).

The RIP calculated using field data on total abundance did not demonstrate a greater impact of the non-native species ($RIP < 1$, Table 2). Considering biomass, the impact of *I. punctatus* was found to be similar to the native species *R. quelen* ($RIP \approx 1$, Table 2). However, using CPUE data, *I. punctatus* was found to have a greater impact ($RIP > 1$, Table 2).

4 Discussion

Both predator species in the present study demonstrated a Type II FR when exposed to *M. microlepis*. Most, if not all, of the prey were consumed at low densities, which suggests that in open waters this could lead to destabilising effects on the local population (Holling, 1959a; Murdoch & Oaten, 1975; Sinclair et al., 1998). A Type II FR may therefore pose a greater concern in open waters, where endangered, such as *M. lateralis*, are often found to exist in small populations (IUCN, 2012; Sinclair et al., 1998).

The CFR approach provided evidence that *I. punctatus* is likely to exert an ecological impact to native small fish prey, and it is possible that other small-bodied endangered fish species suffer predation pressure from *I. punctatus* in a similar manner as *M. microlepis*. Despite the overlap of FR curves of *I. punctatus* and its native comparator *R. quelen* (Figure 1), the results of the present study suggest that *I. punctatus*

exerts a greater predation pressure on *M. microlepis*, with a significant lower handling time, and therefore a higher maximum feeding rate. Lower h and consequently higher $1/hT$, due to faster rates in processing prey, can explain the superior impact of invaders (Alexander, Dick, Weyl, Robinson & Richardson, 2014). The lower handling time of *I. punctatus* (Table 1) may be due to differences in the resource acquisition ability of the species (Alexander et al., 2014). *Ictalurus punctatus* has taste buds across its body and within its barbels, which aid in perception of food and trigger feeding behaviour (Caprio et al., 1993; Valentincic & Caprio, 1994). *Ictalurus punctatus* also has relatively larger eyes than other siluriform catfishes, which may facilitate visual predation in clean waters, such as of the experimental conditions (Davis, 1959). The similar FRs of *I. punctatus* and *R. quelen* towards the prey (Figure 1) also suggest that these species may have niche overlap, resulting in either resource competition or repartition in open waters. Both species seems to use the same strategy to find resources, as their barbels are sensitive to amino acids or hydrolysed proteins, which triggers feeding behaviour (Caprio, 1975; Broggi, 2014).

Ictalurus punctatus are less abundant than native *R. quelen* in the River Guaraguaçu, which suggests a lower relative predation impact than the native species. However, using standardised CPUE data, a different result emerged, with *I. punctatus* presenting a higher abundance (Table 2), and therefore a potentially greater impact than *R. quelen*. Using only the FR provides a modestly-powered predictor of non-native species impacts, because a low FR can be compensated by high abundance in the field (Lavery et al., 2017). The results obtained here, of a greater RIP using CPUE field data (Table 2), are sufficient to justify the need of management and control of *I. punctatus* population. Early interventions may be successful in avoiding non-native species impacts

(Simberloff, 2003), and in the present case, the results provide evidence to motivate the immediate action to manage *I. punctatus* at low population levels.

Propagule pressure is widely believed to play a fundamental role on the successful establishment and permanency of non-native species, contributing to its continuous impact (Copp, Templeton & Gozlan, 2007; Kolar & Lodge, 2001; Ruesink, 2005). The juvenile *I. punctatus* used in these experiments were obtained from local aquaculture, which is recognised as the major pathway of introduction of *I. punctatus* and other freshwater invasive species (CABI, 2019; Casal, 2006; Gozlan, 2008; Naylor, Williams & Strong, 2001). *Ictalurus punctatus* is among the most cultured freshwater fishes globally, and its global total production in 2016 reached 432,932 tons (FAO, 2018). There are many fish-farms in Paraná State that cultivate and sell young-of-the-year *I. punctatus* despite a legislation that classifies this species as Category I, which has its “transportation, breeding, release or translocation, cultivation, propagation, commercialization, donation or intentional acquisition prohibited by any means” (Environmental Institute of Paraná, 2015). Despite this, the production of *I. punctatus* continues to increase, and the risk of escapes from aquaculture ponds is a constant threat in Neotropical streams, contributing to a high propagule pressure (Forneck, Dutra, Zacarkim & Cunico, 2016; Lima Junior, Pelicice, Vitule & Agostinho, 2012; Vitule, Freire & Simberloff, 2009). This failed attempt to ban the introduction of *I. punctatus* in Paraná could be responsible for the declines in status of many endemic fishes (Lima Junior et al., 2018; Pelicice et al., 2017; Vitule et al., 2009). Therefore, the cultivation of the *I. punctatus* should be avoided in Paraná in order to reduce its spreading into natural water bodies.

Besides constant introductions of *I. punctatus*, the species’ reproductive behaviour (guarders, nest spawners) is more specialised than that of *R. quelen*

(nonguarders, open substratum spawners) (see Balon, 1975). The male *I. punctatus* guards the eggs until they hatch, a feature that may favour the establishment of the non-native species in the field (Marchetti, Moyle & Levine, 2004; Tatarenkov et al., 2006). Also, predation of *I. punctatus* by native fishes in the study region is unlikely due to the dorsal and pectoral spines of *I. punctatus* (Bosher, Newton & Fine, 2006).

In the present study, juveniles of similar body size were used, which contrasts natural populations. Predator body-size influences most aspects of feeding behaviour and there are allometric relationships between body size and the FR parameters, with increases in a and decreases in h as predator size increases (Brose, 2010; González-Suárez et al., 2011; Miller, Crowder, Rice & Binkowski, 1992; Vucic-Pestic, Rall, Kalinkat & Brose, 2010). In the case of the two study species, *I. punctatus* can reach larger adult sizes than *R. quelen*, as the first can be mature at size 300 mm TL and *R. quelen* is already mature at 175 mm TL (Gomes et al., 2000; Jackson, 2004; Shephard & Jackson, 2005), which may lead to an increased *per capita* effect of the non-native species that could influence the FR results.

As with many fishes, the degree of piscivory tends to increase in *I. punctatus* with increasing body length (Starostka & Nelson, 1964). As juveniles were used here, it can be expected a greater *per capita* effect of adults of in the field. Furthermore, omnivores like *I. punctatus* tend to present a Type II FR when exposed to a single type of prey, and the presence of an alternative food source may therefore affect our results (Médoc, Thuillier & Spataro, 2017; Murdoch & Oaten, 1975). Indeed, alternative food availability has been shown to change FRs of amphipods, killer shrimp *Dikerogammarus villosus* and the freshwater shrimp *Gammarus pulex*, with the superior consumption of the invasive *D. villosus* disappearing when it was presented with an alternative non-animal food (Médoc, Thuillier & Spataro, 2017).

Differences in predator biomass also likely strengthen the impact of *I. punctatus*.

The mean biomass of *I. punctatus* specimens captured in the River Guaraguaçu was more than twice of the native species and the RIP using biomass was ≈ 1 (Table 2), which suggests that the impact of *I. punctatus* in the wild is greater than the native predator despite its lower abundance. Neither predator species nor size alone can predict consistently the FR type and its associated parameters, so both predator identity and size should be considered to assess fully the potential ecological impacts of a non-native species on invaded communities (Anderson, Linares, Dodson & Semlitsch, 2016; Guo, Sheath, Amat Trigo & Britton, 2016).

It is important to consider other factors that may alter FR results, such as habitat complexity (Alexander, Dick, O'Connor, Haddaway & Farnsworth, 2012; Barrios-O'Neill, Dick, Emmerson, Ricciardi & Macisaac, 2015), which in some cases can change the FR type and parameters. This can occur if habitat complexity acts as a refuge for prey in low density (Alexander, Kaiser, Weyl & Dick, 2015) or contributes to predation of ambush predators, resulting in a higher consumption rate (Santos, García-Berthou, Hayashi & Santos, 2013). This context-dependency factor may also be an important feature to be tested in the FR of *I. punctatus*, as the River Guaraguaçu has been invaded by the African signalgrass *Urochloa arrecta*, a macrophyte that increases habitat complexity (Michelan, Thomaz, Mormul & Carvalho, 2010; Thomaz & Cunha, 2010; Vitule, Umbria & Aranha, 2006).

To quantify and predict the impacts of non-native species is of fundamental importance to conservation efforts, because such data are needed in order to create plans for management and control of damaging species. The CFR approach and the RIP metric have been shown as an effective tool for predicting the impact of existing and emerging high impact non-native species and can be performed elsewhere in a very simple fashion

(Alexander et al., 2014; Dick et al., 2017a, 2017b; Laverty et al., 2017). Therefore, the CFR approach is of particular usefulness in both developing and developed countries because data on non-native fish impacts are lacking for most species. Whereas, data on non-native species impacts are even more scarce in less developed regions, where the number of potentially threatened endemic species is often high (Frehse, Braga, Nocera & Vitule, 2016; Lövei, Lewinsohn & Network, 2012; Nuñez & Pauchard, 2010; Vitule et al., 2017).

References

- Abilhoa, V., & Duboc, L. F. (2004). Peixes. In S. B. Mikich & R. S. Bérnils (Eds), *Livro vermelho dos animais ameaçados de extinção no Estado do Paraná* (pp. 581–677). Curitiba: Mater Natura e Instituto Ambiental do Paraná.
- Alexander, M. E., Dick, J. T. A., O'Connor, N. E., Haddaway, N. R., & Farnsworth, K. D. (2012). Functional responses of the intertidal amphipod *Echinogammarus marinus*: effects of prey supply, model selection and habitat complexity. *Marine Ecology Progress Series*, 468, 191–202.
- Alexander, M. E., Dick, J. T. A., Weyl, O. L. F., Robinson, T. B., & Richardson, D. M. (2014). Existing and emerging high impact invasive species are characterized by higher functional responses than natives. *Biology Letters*, 10, 2–6.
- Alexander, M. E., Kaiser, H., Weyl, O. L. F., & Dick, J. T. A. (2015). Habitat simplification increases the impact of a freshwater invasive fish. *Environmental Biology of Fishes*, 98, 477–486.
- Anderson, T. L., Linares, C., Dodson, K. N., & Semlitsch, R. D. (2016). Variability in functional response curves among larval salamanders: comparisons across species and size classes. *Canadian Journal of Zoology*, 94, 23–30.
- Bailey, R. M., & Harrison, H. M. (1948). Food habits of the southern channel catfish (*Ictalurus lacustris punctatus*) in the Des Moines River, Iowa. *Transactions of the American*

338 *Fisheries Society*, 75, 110–138.

339 Balon, E. K. (1975). Reproductive guilds of fishes: a proposal and definition. *Journal of the*
340 *Fisheries Research Board of Canada*, 32, 821–864.

341 Barrios-O'Neill, D., Dick, J. T. A., Emmerson, M. C., Ricciardi, A., & MacIsaac, H. J. (2015).
342 Predator-free space, functional responses and biological invasions. *Functional Ecology*,
343 29, 377–384.

344 Becker, G. C. (1983). *Fishes of Wisconsin*. Madison: University of Wisconsin Press.

345 Bellard, C., Cassey, P., & Blackburn, T. M. (2016). Alien species as a driver of recent
346 extinctions. *Biology Letters*, 12, 20150623.

347 Bellard, C., & Jeschke, J. (2016). A spatial mismatch between invader impacts and research
348 publications. *Conservation Biology*, 30, 230–232.

349 Bolker, B. M. (2008). *Ecological models and data in R*. Princeton: Princeton University Press.

350 Boshier, B. T., Newton, S. H., & Fine, M. L. (2006). The spines of the channel catfish, *Ictalurus*
351 *punctatus*, as an anti-predator adaptation: an experimental study. *Ethology*, 112, 188–195.

352 Braga, R. R., Braga, M. R., & Vitule, J. R. S. (2013). Population structure and reproduction of
353 *Mimagoniates microlepis* with a new hypothesis of ontogenetic migration: implications for
354 stream fish conservation in the Neotropics. *Environmental Biology of Fishes*, 96, 21–31.

355 Braun, A. P., & Phelps, Q. E. (2016). Channel catfish habitat use and diet in the Middle
356 Mississippi River. *The American Midland Naturalist*, 175, 47–54.

357 Broggi, J. A. (2014). Hidrolisado proteico de sardinha (Clupeidae) como atrativo alimentar para
358 o jundiá (*Rhamdia quelen*). Master thesis, Universidade do Estado de Santa Catarina. 49 p.

359 Brose, U. (2010). Body-mass constraints on foraging behaviour determine population and food-
360 web dynamics. *Functional Ecology*, 24, 28–34.

361 CABI, 2019. *Ictalurus punctatus* [original text by C Engle]. In *Invasive Species Compendium*.
362 Wallingford, UK: CAB International. www.cabi.org/isc.

363 Caprio, J. (1975). High sensitivity of catfish taste receptors to amino acids. *Comparative*
364 *Biochemistry and Physiology B*, 52, 247–251.

365 Caprio, J., Brand, J. G., Teeter, J. H., Valentincic, T., Kalinoski, D. L., Kohbara, J., ... Wegert,

366 S. (1993). The taste system of the channel catfish: from biophysics to behavior. *Trends in*
367 *Neurosciences*, 16, 192–197.

368 Casal, C. M. V. (2006). Global documentation of fish introductions: the growing crisis and
369 recommendations for action. *Biological Invasions*, 8, 3–11.

370 Clavero, M., & García-Berthou, E. (2005). Invasive species are a leading cause of animal
371 extinctions. *Trends in Ecology and Evolution*, 20, 110.

372 Copp, G. H., Templeton, M., & Gozlan, R. E. (2007). Propagule pressure and the invasion risks
373 of non-native freshwater fishes in Europe: a case study of England. *Journal of Fish*
374 *Biology*, 71 (Supplement D), 148–159.

375 Cox, J. G., & Lima, S. L. (2006). Naiveté and an aquatic-terrestrial dichotomy in the effects of
376 introduced predators. *Trends in Ecology and Evolution*, 21, 674–680.

377 Cruz-Spindler, S., Leal, M. E., Lehmann, P. C., & Schulz, U. H. (2012). First record of the
378 exotic channel catfish *Ictalurus punctatus* (Rafinesque 1818) (Siluriformes: Ictaluridae) in
379 the Rio dos Sinos basin, RS, Brazil. *Biota Neotropica*, 12, 64–67.

380 Davis, J. (1959). Management of channel catfish in Kansas. *Museum of Natural History*
381 *Miscellaneous Publication*, 21, 1–56.

382 Dick, J. T. A., Alexander, M. E., Jeschke, J., Ricciardi, A., MacIsaac, H. J., Robinson, T. B., ...
383 Richardson, D. M. (2014). Advancing impact prediction and hypothesis testing in invasion
384 ecology using a comparative functional response approach. *Biological Invasions*, 16, 735–
385 753.

386 Dick, J. T. A., Alexander, M. E., Ricciardi, A., Lavery, C., Downey, P. O., Xu, M., ... Shaw,
387 R. H. (2017a). Functional responses can unify invasion ecology. *Biological Invasions*, 19,
388 1667–1672.

389 Dick, J. T. A., Lavery, C., Lennon, J. J., Barrios-O'Neill, D., Mensink, P. J., Britton, J. R., ...
390 Caffrey, J. M. (2017b). Invader Relative Impact Potential: a new metric to understand and
391 predict the ecological impacts of existing, emerging and future invasive alien species.
392 *Journal of Applied Ecology*, 54, 1259–1267.

393 Endo, T., Kaneko, S., Igari, K., Kanou, K., Nakazato, R., Kamei, R., ... Hyakunari, W. (2015).

394 Feeding characteristics of the channel catfish *Ictalurus punctatus* in the littoral zone of
 395 Lake Kitaura, Ibaraki Prefecture, Japan. *Aquaculture Science*, 63, 49–58.

396 FAO. (2018). FAO yearbook. Fishery and Aquaculture Statistics 2016. Rome: FAO.

397 Forneck, S. C., Dutra, F. M., Zacarkim, C. E., & Cunico, A. M. (2016). Invasion risks by non-
 398 native freshwater fishes due to aquaculture activity in a Neotropical stream.
 399 *Hydrobiologia*, 773, 193–205.

400 Frehse, F. A., Braga, R. R., Nocera, G. A., & Vitule, J. R. S. (2016). Non-native species and
 401 invasion biology in a megadiverse country: scientometric analysis and ecological
 402 interactions in Brazil. *Biological Invasions*, 18, 3713–3725.

403 Gallardo, B., Clavero, M., Sánchez, M. I., & Vilà, M. (2016). Global ecological impacts of
 404 invasive species in aquatic ecosystems. *Global Change Biology*, 22, 151–163.

405 Gomes, L. C., Golombieski, J. I., Gomes, A. R. C., & Baldisserotto, B. (2000). Biologia do
 406 jundiá *Rhamdia quelen* (Teleostei, Pimelodidae). *Ciência Rural*, 30, 179–185.

407 González-Suárez, M., Mugabo, M., Decencièrre, B., Perret, S., Claessen, D., & Le Galliard, J. F.
 408 (2011). Disentangling the effects of predator body size and prey density on prey
 409 consumption in a lizard. *Functional Ecology*, 25, 158–165.

410 Goudie, C., Davis, K., & Simco, B. (1983). Influence of the eyes and pineal gland on locomotor
 411 activity patterns of channel catfish *Ictalurus punctatus*. *Physiological Zoology*, 56, 10–17.

412 Gozlan, R. E. (2008). Introduction of non-native freshwater fish: is it all bad? *Fish and*
 413 *Fisheries*, 9, 106–115.

414 Guedes, D. S. (1980). *Contribuição ao estudo da sistemática e alimentação de jundiás*
 415 *(Rhamdia spp) na região central do Rio Grande do Sul (Pisces, Pimelodidae)*. Master
 416 thesis, Universidade Federal de Santa Maria. 99 p.

417 Guo, Z., Sheath, D., Amat Trigo, F., & Britton, J. R. (2016). Comparative functional responses
 418 of native and high-impacting invasive fishes: impact predictions for native prey
 419 populations. *Ecology of Freshwater Fish*, 26, 533–540.

420 Gutierrez, S. M. M., Vitule, J. R. S., Freire, C. A., & Prodocimo, V. (2014). Physiological tools
 421 to predict invasiveness and spread via estuarine bridges: tolerance of Brazilian native and

422 worldwide introduced freshwater fishes to increased salinity. *Marine and Freshwater*
423 *Research*, 65, 425–436.

424 Haubrock, P. J., Balzani, P., Johovic, I., Inghilesi, A. F., Nocita, A., & Tricarico, E. (2018). The
425 diet of the alien channel catfish *Ictalurus punctatus* in the River Arno (Central Italy).
426 *Aquatic Invasions*, 13, 575–585.

427 Hill, T. D., Duffy, W. G., & Thompson, M. R. (1995). Food-habits of channel catfish in Lake
428 Oahe, South-Dakota. *Journal of Freshwater Ecology*, 10, 319–323.

429 Holling, C. S. (1959a). Some characteristics of simple types of predation and parasitism. *The*
430 *Canadian Entomologist*, 91, 385–398.

431 Holling, C. S. (1959b). The components of predation as revealed by a study of small-mammal
432 predation of the European pine sawfly. *The Canadian Entomologist*, 91, 234–261.

433 Environmental Institute of Paraná. (2015). Portaria IAP nº 059, de 15 de abril de 2015. Curitiba:
434 Instituto Ambiental do Paraná.
435 (www.iap.pr.gov.br/arquivos/File/Lista_invasoras_PR_corrigida_set_2015.pdf)

436 IUCN. (2012). *IUCN Red List Categories and Criteria: Version 3.1*. (2nd ed.). Gland,
437 Switzerland and Cambridge, UK.

438 Jackson, D. C. (2004). Natural history and fisheries. In C. S. Tucker & J. A. Hargreaves (Eds),
439 *Biology and culture of channel catfish* (pp. 15–35). Amsterdam: Elsevier.

440 Jeschke, J., Kopp, M., & Tollrian, R. (2002). Predator functional responses: discriminating
441 between handling and digesting prey. *Ecological Monographs*, 72, 95–112.

442 Jordan, S. M., Neumann, R. M., & Schultz, E. T. (2004). Distribution, habitat use, growth, and
443 condition of a native and an introduced catfish species in the Hudson River estuary.
444 *Journal of Freshwater Ecology*, 19, 59–67.

445 Juliano, S. A. (2001). Nonlinear curve fitting. In S. M. Scheiner & J. Gurevitch (Eds), *Design*
446 *and analysis of ecological experiments* (pp. 178–196). Oxford: Oxford University Press.

447 Kolar, C. S., & Lodge, D. M. (2001). Progress in invasion biology: predicting invaders. *Trends*
448 *in Ecology and Evolution*, 16, 199–204.

449 Kumschick, S., Gaertner, M., Vilà, M., Essl, F., Jeschke, J., Pyšek, P., ... Winter, M. (2015).

450 Ecological impacts of alien species: quantification, scope, caveats, and recommendations.
 451 *BioScience*, 65, 55–63.

452 Latini, A. O., & Petrere Jr., M. (2004). Reduction of a native fish fauna by alien species: an
 453 example from Brazilian freshwater tropical lakes. *Fisheries Management and Ecology*, 11,
 454 71–79.

455 Laverty, C., Green, K. D., Dick, J. T. A., Barrios-O'Neill, D., Mensink, P. J., Médoc, V., ...
 456 Gallagher, C. (2017). Assessing the ecological impacts of invasive species based on their
 457 functional responses and abundances. *Biological Invasions*, 19, 1653–1665.

458 Lima Junior, D. P., Magalhães, A. L. B., Pelicice, F. M., Vitule, J. R. S., Azevedo-Santos, V.
 459 M., Orsi, M. L., ... Agostinho, A. A. (2018). Aquaculture expansion in Brazilian
 460 freshwaters against the Aichi Biodiversity Targets. *Ambio*, 47, 427–440.

461 Lima Junior, D. P., Pelicice, F. M., Vitule, J. R. S., & Agostinho, A. A. (2012). Aquaculture,
 462 politics and environment in Brazil: new proposals and old misconceptions. *Natureza &*
 463 *Conservação*, 10, 88–91.

464 Lövei, G. L., Lewinsohn, T. M., & the Biological Invasions in Megadiverse Regions Network
 465 (2012). Megadiverse developing countries face huge risks from invasives. *Trends in*
 466 *Ecology and Evolution*, 27(1), 2–3.

467 Lowry, E., Rollinson, E. J., Laybourn, A. J., Scott, T. E., Aiello-Lammens, M. E., Gray, S. M.,
 468 ... Gurevitch, J. (2013). Biological invasions: a field synopsis, systematic review, and
 469 database of the literature. *Ecology and Evolution*, 3, 182–196.

470 Marchetti, M. P., Moyle, P. B., & Levine, R. (2004). Invasive species profiling? Exploring the
 471 characteristics of non-native fishes across invasion stages in California. *Freshwater*
 472 *Biology*, 49, 646–661.

473 Matsuzaki, S. S., Takamura, N., Arayama, K., Tominaga, A., Iwasaki, J., & Washitani, I.
 474 (2011). Potential impacts of non-native channel catfish on commercially important species
 475 in a Japanese lake, as inferred from long-term monitoring data. *Aquatic Conservation:*
 476 *Marine and Freshwater Ecosystems*, 21, 348–358.

477 Médoc, V., Thuillier, L., & Spataro, T. (2017). Opportunistic omnivory impairs our ability to

478 predict invasive species impacts from functional response comparisons. *Biological*
 479 *Invasions*, 20, 1307–1319.

480 Menezes, N. A., & Weitzman, S. H. (2009). Systematics of the neotropical fish subfamily
 481 Glandulocaudinae (Teleostei: Characiformes: Characidae). *Neotropical Ichthyology*, 7,
 482 295–370.

483 Meurer, S., Zaniboni Filho, E. (1997). Hábito alimentar do jundiá *Rhamdia quelen* (Pisces,
 484 Siluriformes, Pimelodidae), na região do alto rio Uruguai. *Anais do XII encontro*
 485 *Brasileiro de ictiologia*.

486 Michelin, T. S., Thomaz, S. M., Mormul, R. P., & Carvalho, P. (2010). Effects of an exotic
 487 invasive macrophyte (tropical signalgrass) on native plant community composition,
 488 species richness and functional diversity. *Freshwater Biology*, 55, 1315–1326.

489 Miller, T. J., Crowder, L. B., Rice, J. A., & Binkowski, F. P. (1992). Body size and the
 490 ontogeny of the functional response in fishes. *Canadian Journal of Fisheries and Aquatic*
 491 *Sciences*, 49, 805–812.

492 MMA. (2014). Portaria nº 445, de 17 de dezembro de 2014. Peixes e Invertebrados Aquáticos
 493 Ameaçados. Ministério do Meio Ambiente.
 494 ([www.icmbio.gov.br/portal/portaldabiodiversidade/50-menu-biodiversidade/6706-](http://www.icmbio.gov.br/portal/portaldabiodiversidade/50-menu-biodiversidade/6706-portarias-fauna-ameacada)
 495 [portarias-fauna-ameacada](http://www.icmbio.gov.br/portal/portaldabiodiversidade/50-menu-biodiversidade/6706-portarias-fauna-ameacada))

496 Murdoch, W. W., & Oaten, A. (1975). Predation and population stability. *Advances in*
 497 *Ecological Research*, 9, 1–131.

498 Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. B., & Kent, J. (2000).
 499 Biodiversity hotspots for conservation priorities. *Nature*, 403, 853–858.

500 Naylor, R. L., Williams, S. L., & Strong, D. R. (2001). A gateway for exotic species. *Science*,
 501 294, 1655–1656.

502 Nuñez, M. A., & Pauchard, A. (2010). Biological invasions in developing and developed
 503 countries: does one model fit all? *Biological Invasions*, 12, 707–714.

504 Oaten, A., & Murdoch, W. W. (1975). Switching, functional response, and stability in predator-
 505 prey systems. *The American Naturalist*, 109, 299–318.

506 Olden, J. D., & Poff, N. L. (2005). Long-term trends of native and non-native fish faunas in the
507 American Southwest. *Animal Biodiversity and Conservation*, 28, 75–89.

508 Orsi, M. L., & Agostinho, A. A. (1999). Introdução de espécies de peixes por escapes acidentais
509 de tanques de cultivo em rios da Bacia do Rio Paraná, Brasil. *Revista Brasileira de*
510 *Zoologia*, 16, 557–560.

511 Ota, R. R., Message, H. J., Graça, W. J. , & Pavanelli, C. S. (2015). Neotropical siluriformes as
512 a model for insights on determining biodiversity of animal groups. *PLoS ONE*, 10(7), 1–
513 13.

514 Paolucci, E. M., MacIsaac, H. J., & Ricciardi, A. (2013). Origin matters: alien consumers inflict
515 greater damage on prey populations than do native consumers. *Diversity and Distributions*,
516 19, 988–995.

517 Parker, I. M., Simberloff, D., Lonsdale, W., Goodell, K., Wonham, M., Kareiva, P., ...
518 Goldwasser, L. (1999). Impact: toward a framework for understanding the ecological
519 effects of invaders. *Biological Invasions*, 1, 3–19.

520 Paterson, R. A., Dick, J. T. A., Pritchard, D. W., Ennis, M., Hatcher, M. J., & Dunn, A. M.
521 (2015). Predicting invasive species impacts: a community module functional response
522 approach reveals context dependencies. *Journal of Animal Ecology*, 84, 453–463.

523 Pelicice, F. M., Azevedo-Santos, V. M., Vitule, J. R. S., Orsi, M. L., Lima Junior, D. P.,
524 Magalhães, A. L. B., ... Agostinho, A. A. (2017). Neotropical freshwater fishes imperilled
525 by unsustainable policies. *Fish and Fisheries*, 18, 1119–1133.

526 Piedras, S. R. N., Pouey, J. L. O. F., & Moraes, P. R. R. (2006). Comportamento alimentar e
527 reprodutivo de peixes exóticos e nativos cultivados na zona sul do Rio Grande do Sul.
528 *Revista Brasileira de Agrociência*, 12, 341–344.

529 Pritchard, D. W., Paterson, R. A., Bovy, H. C., & Barrios-O'Neill, D. (2017). frair: an R
530 package for fitting and comparing consumer functional responses. *Methods in Ecology and*
531 *Evolution*, 8, 1528–1534.

532 R Core Team. (2015). R: a language and environment for statistical computing. Vienna: R
533 Foundation for Statistical Computing. (www.r-project.org/)

534 Ricciardi, A., & Atkinson, S. K. (2004). Distinctiveness magnifies the impact of biological
535 invaders in aquatic ecosystems. *Ecology Letters*, 7, 781–784.

536 Ricciardi, A., Hoopes, M., Marchetti, M. P., & Lockwood, J. L. (2013). Progress toward
537 understanding the ecological impacts of non-native species. *Ecological Monographs*, 83,
538 263–282.

539 Rogers, D. (1972). Random search and insect population models. *Journal of Animal Ecology*,
540 41, 369–383.

541 Rosa, R. S., & Lima, F. C. T. (2008). Os peixes brasileiros ameaçados de extinção. In A. B. M.
542 Machado, G. M. Drummond, & A. P. Paglia (Eds), *Livro vermelho da fauna Brasileira*
543 *ameaçada de extinção* (pp. 9–278). Belo Horizonte: Fundação Biodiversitas.

544 Ruesink, J. L. (2005). Global analysis of factors affecting the outcome of freshwater fish
545 introductions. *Conservation Biology*, 19, 1883–1893.

546 Salo, P., Korpimäki, E., Banks, P. B., Nordström, M., & Dickman, C. R. (2007). Alien predators
547 are more dangerous than native predators to prey populations. *Proceedings of the Royal*
548 *Society B*, 274, 1237–1243.

549 Santos, A. F. G. N., García-Berthou, E., Hayashi, C., & Santos, L. N. (2013). When habitat
550 complexity increases predation risk: experiments with invasive and neotropical native
551 fishes. *Marine and Freshwater Research*, 64, 752–760.

552 Sharpe, D. M. T., De León, L. F., González, R., & Torchin, M. E. (2017). Tropical fish
553 community does not recover 45 years after predator introduction. *Ecology*, 98, 412–424.

554 Shephard, S., & Jackson, D. C. (2005). Channel catfish maturation in Mississippi streams. *North*
555 *American Journal of Fisheries Management*, 25, 1467–1475.

556 Silfvergrip, A. M. C. (1996). *A sistematic revision of the neotropical catfish genus Rhamdia*
557 *(Teleostei, Pimelodidae)*. Doctoral dissertation, Stockholm University. 156 p.

558 Simberloff, D. (2003). How much information on population biology is needed to manage
559 introduced species? *Conservation Biology*, 17, 83–92.

560 Simberloff, D., Martin, J. L., Genovesi, P., Maris, V., Wardle, D. A., Aronson, J., ... Vilà, M.
561 (2013). Impacts of biological invasions: what's what and the way forward. *Trends in*

562 *Ecology and Evolution*, 28, 58–66.

563 Simberloff, D., & Vitule, J. R. S. (2014). A call for an end to calls for the end of invasion

564 biology. *Oikos*, 123, 408–413.

565 Sinclair, A. R. E., Pech, R. P., Dickman, C. R., Hik, D., Mahon, P., & Newsome, A. E. (1998).

566 Predicting effects of predation on conservation of endangered prey. *Conservation Biology*,

567 12, 564–575.

568 Speziale, K. L., Lambertucci, S. A., Carrete, M., & Tella, J. L. (2012). Dealing with non-native

569 species: what makes the difference in South America? *Biological Invasions*, 14, 1609–

570 1621.

571 Starostka, V. J., & Nelson, W. R. (1974). Age, growth, sexual maturity, and food of channel

572 catfish in central Lake Oahe, 1968–1969. In *Technical papers of the U.S. Fish and Wildlife*

573 *Service*. Washington D. C.: U.S. Dept. of the Interior, Fish and Wildlife Service.

574 Tatarenkov, A., Barreto, F., Winkelman, D. L., & Avise, J. C. (2006). Genetic monogamy in the

575 channel catfish, *Ictalurus punctatus*, a species with uniparental nest guarding. *Copeia*,

576 2006, 735–741.

577 Thomaz, S. M., & Cunha, E. R. (2010). The role of macrophytes in habitat structuring in aquatic

578 ecosystems: methods of measurement, causes and consequences on animal assemblages’

579 composition and biodiversity. *Acta Limnologica Brasiliensia*, 22, 218–236.

580 Townsend, C. R., & Winterbourn, M. J. (1992). Assessment of the environmental risk posed by

581 an exotic fish: the proposed introduction of channel catfish (*Ictalurus punctatus*) to New

582 Zealand. *Conservation Biology*, 6, 273–282.

583 Troca, D. F. A., & Vieira, J. P. (2012). Potencial invasor dos peixes não nativos cultivados na

584 região costeira do Rio Grande do Sul, Brasil. *Boletim do Instituto de Pesca*, 38, 109–120.

585 Trumpickas, J., Mandrak, N. E., & Ricciardi, A. (2011). Nearshore fish assemblages associated

586 with introduced predatory fishes in lakes. *Aquatic Conservation: Marine and Freshwater*

587 *Ecosystems*, 21, 338–347.

588 Tyus, H. M., & Nikirk, N. J. (1990). Abundance, growth, and diet of channel catfish, *Ictalurus*

589 *punctatus*, in the Green and Yampa rivers, Colorado and Utah. *The Southwestern*

590 *Naturalist*, 35, 188–198.

591 Valentincic, T. B., & Caprio, J. (1994). Chemical and visual control of feeding and escape
 592 behaviors in the channel catfish *Ictalurus punctatus*. *Physiology & Behavior*, 55, 845–855.

593 Vitousek, P. M., D’Antonio, C., Loope, L. L., & Westbrooks, R. (1996). Biological invasions as
 594 global environmental change. *American Scientist*, 84, 468–478.

595 Vitule, J. R. S. (2008). *Distribuição, abundância e estrutura populacional de peixes*
 596 *introduzidos no Rio Guaraguaçu, Paranaguá, Paraná, Brasil*. Doctoral dissertation,
 597 Universidade Federal do Paraná. 143 p.

598 Vitule, J. R. S., Agostinho, A. A., Azevedo-Santos, V. M., Daga, V. S., Darwall, W. R. T.,
 599 Fitzgerald, D. B., ... Winemiller, K. O. (2017). We need better understanding about
 600 functional diversity and vulnerability of tropical freshwater fishes. *Biodiversity and*
 601 *Conservation*, 26, 757–762.

602 Vitule, J. R. S., Freire, C. A., & Simberloff, D. (2009). Introduction of non-native freshwater
 603 fish can certainly be bad. *Fish and Fisheries*, 10, 98–108.

604 Vitule, J. R. S., Freire, C. A., Vazquez, D. P., Nuñez, M. A., & Simberloff, D. (2012).
 605 Revisiting the potential conservation value of non-native species. *Conservation Biology*,
 606 26, 1153–1155.

607 Vitule, J. R. S., Umbria, S. C., & Aranha, J. M. R. (2005). Registro de ocorrência do bagre-do-
 608 canal *Ictalurus punctatus* (Rafinesque, 1818) em uma importante bacia hidrográfica da
 609 Floresta Atlântica paranaense. In *I simpósio Brasileiro sobre espécies exóticas invasoras*.
 610 (www.mma.gov.br/port/sbf/invasoras/index.cfm)

611 Vitule, J. R. S., Umbria, S. C., & Aranha, J. M. R. (2006). Introduction of the African catfish
 612 *Clarias gariepinus* (Burchell, 1822) into Southern Brazil. *Biological Invasions*, 8, 677–
 613 681.

614 Vucic-Pestic, O., Rall, B. C., Kalinkat, G., & Brose, U. (2010). Allometric functional response
 615 model: body masses constrain interaction strengths. *Journal of Animal Ecology*, 79, 249–
 616 256.

617 Welcomme, R. L. (1988). *International introductions of inland aquatic species*. Rome: FAO

Fisheries Technical Paper.

Wellborn, T. L. (1988). Channel catfish: life history and biology. *Southern Regional Aquaculture Center Publication, 180*. (srac.tamu.edu/viewCategory/3)

Zanatta, A. S., Ramos, I. P., da Silva, R. J., Langeani, F., & Carvalho, E. D. (2010). Pisces, Siluriformes, Ictaluridae, *Ictalurus punctatus* (Rafinesque, 1818): first record in middle Paranapanema river reservoir, aquaculture and exotic species dispersion. *Check List*, 6, 589–591.

Figure legend

Figure 1. Type II Functional Response curves for channel catfish *Ictalurus punctatus* (dashed line) and South American silver catfish *Rhamdia quelen* (solid line) preying on blue tetra *Mimagoniates microlepis*. Points indicate mean consumption per density. Shading represents bootstrapped ($n = 2000$) 95% confidence intervals for each species.

Assessing the impacts of the channel catfish *Ictalurus punctatus* using the Comparative Functional Response approach

Table S1. Field data on total abundance (n), capture per unit effort (CPUE) and total biomass (in grams) of channel catfish *Ictalurus punctatus* and South American silver catfish *Rhamdia quelen* captured in 17 standardised samples, between 2005 and 2007, using different fishing gears. Data from Vitule (2008). Long lines were of 10 m length, and gill nets were of 30×1.70 m, with exposure times in all cases being 1500 hours.

Fishing gear (and mesh size)	Effort	n	CPUE	Biomass (g)
------------------------------	--------	-----	------	-------------

<i>Ictalurus punctatus</i>					
Longline	3 longlines with 10 fish hooks	8	0.000178	9312	
Gill net (20 mm)	Gill net	5	6.54E-05	1069	
Gill net (40 mm)	Gill net	4	5.23E-05	2075	
Gill net (60 mm)	Gill net	1	1.31E-05	740	
Fishing rod	4 anglers, $\approx$ 60 h	$\frac{1}{2}$	0.05	14496	
<i>Rhamdia quelen</i>					
Longline	3 longlines with 10 fish hooks	40	0.000889	14106	
Gill net (20 m)	Gill net	31	0.000405	8133	
Gill net (40 mm)	Gill net	33	0.000431	14768	
Fishing rod	4 anglers, $\approx$ 60 h	8	0.033333	1978	

641

642