UWS Academic Portal

A new taxonomy for post-activation potentiation in sport

Boullosa, Daniel; Beato, Marco; Dello Iacono, Antonio; Cuenca-Fernández, Francisco; Doma, Kenji; Schumann, Moritz; Zagatto, Alessandro Moura; Loturco, Irineu; Behm, David

Published in:
International Journal of Sports Physiology and Performance

DOI:
[10.1123/ijsp.2020-0350](https://doi.org/10.1123/ijsp.2020-0350)

E-pub ahead of print: 20/08/2020

Document Version
Peer reviewed version

[Link to publication on the UWS Academic Portal](#)

Citation for published version (APA):
Boullosa, D., Beato, M., Dello Iacono, A., Cuenca-Fernández, F., Doma, K., Schumann, M., Zagatto, A. M., Loturco, I., & Behm, D. (2020). A new taxonomy for post-activation potentiation in sport. *International Journal of Sports Physiology and Performance*, 15(8), 1197-1200. <https://doi.org/10.1123/ijsp.2020-0350>

General rights

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact pure@uws.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

A New Taxonomy for Post-activation Potentiation in Sport

Abstract

Post-activation potentiation (PAP) mechanisms and responses have a long scientific history. However, to this day, there is still controversy regarding the mechanisms underlying enhanced performance following a conditioning activity. More recently, the term post-activation performance enhancement (PAPE) has been proposed with differing associated mechanisms and protocols than with PAP. However, these two terms (PAP and PAPE) may not adequately describe all specific potentiation responses and mechanisms and can be also complementary in some cases. **Purpose:** this commentary presents and discuss the similarities and differences between PAP and PAPE, and subsequently elaborates on a new taxonomy for better describing performance potentiation in sport settings. **Conclusion:** the taxonomy elaborated proposes the formula “Post- [CONDITIONING ACTIVITY] [VERIFICATION TEST] potentiation in [POPULATION]”. This taxonomy would avoid erroneous identification of isolated physiological attributes and provide individualization and better applicability of conditioning protocols in sport settings.

Key Words: post-activation performance enhancement; post-activation potentiation; post-tetanic potentiation; power; strength.

Introduction

Post-activation potentiation (PAP) is a muscular phenomenon which consists of an acute increment in strength and power performances as a result of the recent voluntary contractile history.¹ Much of the past literature suggested that PAP-induced augmented power contributed to enhancement of sport-specific tasks such as explosive jumps, sprints, changes of direction, and throws.^{2,3} Many different conditioning activities have been used by coaches and researchers to induce subsequent performance enhancements, including resistance,⁴ ballistic,⁵ and flywheel⁶ exercises. The primary mechanism underlying PAP is the phosphorylation of the myosin regulatory light chain (MLC), a peripheral muscle memory mechanism, which leads to greater peak force and rate of force development (RFD)⁷. Thus, PAP has traditionally been considered one the main objectives during warm up routines.⁸

The role of PAP on sports performance has been recently debated with the proposal of an alternative term, referred to as post-activation performance enhancement (PAPE).⁹ The reasons behind this dualism (PAP vs. PAPE) refers to the association of PAP with evoked twitch verification which, in turn, would be related to MLC phosphorylation, during a very short period of time (<5 min). Conversely, PAPE would be associated with increases in voluntary performance primarily as a consequence of other potential mechanisms (e.g., temperature, water content), over longer time windows (>5 min).¹⁰ While we believe this recent proposal has merit and could shed light on current practices and further studies, we also suggest that this dualism could limit physiological interpretations. That is, the physiological mechanisms underlying an acute effect of a conditioning activity may not be always revealed by the selected test. This does not necessarily mean the inexistence of a physiological effect, but an inadequate signal-to-noise ratio to detect these changes. This fact would better explain the inconsistent results when simultaneously assessing jump performance and twitch verification,¹¹ which represents a serious limitation, since this dualism may be biasing the

search for the link between conditioning activities, physiological mechanisms, and their purported performance improvements. For instance, the deviating time course of performance enhancements (<5 min) and twitch verification (>5 min) after voluntary conditioning activities have also been reported in laboratory conditions.¹¹ Further, a very recent study found that drop jumps (DJ) performed 2-min before the twitch verification test (i.e., PAP), also enhanced supramaximal cycling performance (i.e., PAPE) and glycolytic energy contribution.¹² However, similar to other investigations, this recent study¹² did not verify the MLC phosphorylation levels with muscle biopsies, suggesting that another physiological mechanism may be involved (i.e., glycolytic energy contribution).

Therefore, the major arguments for the PAP vs. PAPE dualism are not well supported in all cases. These limitations highlight the necessity of a taxonomy of PAP to better identify the potentiation effects of conditioning activities in distinct sport settings, without biasing its potential physiological attributes. Thus, this commentary aims to briefly discuss the current knowledge, and justify and propose a new taxonomy for PAP and its possible applications in sport.

Limitations of the current evidence: the problem of linking mechanisms and verification tests

A historical overview of the studies on muscle potentiation reveals an activity-dependent terminology of potentiation studies (see Table 1), which is related to electrically evoked stimuli (i.e., staircase and post-tetanic potentiation) versus voluntary stimuli (i.e., PAP). However, the recent inclusion of the term PAPE does not follow this criterium and is related to the verification procedure, with PAP requiring twitch verification (voluntary activity → twitch verification), while PAPE would be used when the verification occurs with any type of exercise (voluntary activity → voluntary activity). The main reason for this differentiation

would be that twitch verification only evaluates peripheral changes, as it is not influenced by potential spinal and supraspinal influences.¹³ Blazeovich and Babault¹⁰ suggested that other peripheral, non-phosphorylation-dependent processes related to Ca^{2+} sensitivity (i.e., muscle temperature, [pH], and water content) would be more related to PAPE, but there is no reason to disregard that these mechanisms could also influence twitch verification results. In addition, most evidence linking MLC phosphorylation and enhanced force production capacity of fast-twitch fibers has been elaborated from animal models,⁷ with humans presenting evidence of similar MLC phosphorylation levels of both fast- and slow-twitch fibers, which could be related to the different evolutionary paths between species.¹⁴ Further, differences in laboratory (*ex vivo*, *in vivo*, and *in vitro*) and field experiments related to contraction modes (i.e., concentric, isometric, eccentric, stretch-shortening cycle), sarcomere lengths, resting $[\text{Ca}^{2+}]$, and genetic variants of both kinases and phosphatases among other factors,⁷ may also influence the potentiation effects. Furthermore, potentiation phenomena are always accompanied by varying levels of fatigue of different origins,^{15,16} complicating the relationship between potentiation mechanisms and verification tests results. Therefore, the interaction of a number of peripheral potentiation mechanisms with fatigue, makes questionable the sole proposed link between MLC phosphorylation and PAP. Meanwhile, the use of PAPE would only be valid to verify the potentiation effect in voluntary exercises, independent of the potentiation mechanisms involved in each specific case. However, there are situations whereby both PAP and PAPE could be influencing performance. For example, Low et al. (2019)¹⁷ used band-resisted squat jumps as the conditioning activity with a 5×1 km interval runs (3 min of recovery) as well as voluntary and evoked contractile properties as the subsequent performance and mechanism measures. While time to complete specific kilometer intervals (e.g., 1st and 4th km) and jump height and reactive strength index improved, the time to peak twitch decreased (higher rate of force development) following the third kilometer interval as well as

at 10 minutes post-run. The seemingly PAP-induced improvement in the rate of twitch contraction force could not conceivably be attributed to the original conditioning activity, but PAP influences could have been sustained with the kilometer run intervals. Hence, the performance testing measure may have ensured that both PAP and PAPE mechanisms were contributing concurrently. Thus, strict definitions of PAP and PAPE in these situations would not adequately describe the phenomena.

*** Place Table 1 about here ***

Another relevant confounding factor refers to the key influence of athletes' training background⁴ (i.e., strength levels and experience) and sex¹⁸ on potentiation responses, which are not always considered. For instance, speed-power athletes (e.g., sprinters and jumpers) could benefit more from brief, high-intensity conditioning activities, while endurance athletes (e.g., marathon runners and triathletes) would benefit more from submaximal prolonged conditioning activities due to an optimized PAP/fatigue balance.¹⁴ This factor possibly explains the high variability observed between athletes when applying well standardized conditioning protocols.¹⁹

Within this picture, it is inappropriate to definitively link conditioning activities, physiological mechanisms, and verification tests, since the same conditioning activity may enhance strength capacity via several physiological mechanisms, while potentiation responses would be observable with different verification tests. In addition, the role of athletes' characteristics is another important factor to be highlighted when comparing the potentiation effects of distinct conditioning protocols. Furthermore, there is conflicting evidence regarding the influence of spinal and supraspinal mechanisms on potentiation^{20,11,14,21} which may be a problem of sensitivity (i.e., signal-to-noise ratio) that warrants additional research.

123 **The New Taxonomy**

124 Given all the aforementioned limitations of the current state of the art, we propose
125 that the enhancement of any muscle performance, with simple or complex verification tests,
126 could be better taxonomized using another model, which would consider conditioning
127 stimuli, verification tests, and population of athletes, as main factors involved in these
128 relationships and subsequent classifications:

129

130 **Post- [CONDITIONING ACTIVITY] [VERIFICATION TEST] potentiation in**
131 **[POPULATION]**

132

133 Examples:

134 Post-high intensity squatting jump potentiation in resistance trained males.

135 Post-submaximal running jump potentiation in female endurance runners.

136 Post-eccentric flywheel squatting swim start potentiation in varsity trained male swimmers.

137

138 Using this model, any conditioning activity would have its own physiological
139 (potentiating and fatiguing) associated mechanism(s), which could be specifically identified in
140 each case with additional experiments. Moreover, the association of a verification test to the
141 potentiation responses, would assist to better recognize the signal-to-noise ratio after
142 identification of the error associated with the test. Further, the application of the conditioning
143 activity to a very homogenous group of athletes, would minimize the variability of potentiation
144 responses, therefore favoring the validity and applicability of the findings. Of note, considering
145 that potentiation mechanisms are mainly muscle memory mechanisms,⁷ it should be preferable
146 that athletes be evaluated with well-known exercises or, at least, be fully familiarized before
147 testing. Finally, the terms PAP and PAPE could be independently used when appropriate, with

PAP being applicable in most cases when a conditioning protocol is followed by a single exercise as verification test. However, as recently observed,¹² the simultaneous existence of PAP and PAPE should not be disregarded. In this manner, this recent study¹² would be seen as: “Post-drop jumps supramaximal cycling potentiation in recreational male cyclists (via PAP and augmented glycolytic energy contribution). Similarly, another recent study³ found that a variety of conditioning protocols resulted in enhanced change of direction performances (i.e., PAPE) with observable changes in tensiomyography parameters (i.e., PAP).

Practical Applications: The Need for Individualized Approaches

A thorough examination of the contemporary potentiation (PAP and PAPE) scientific literature highlights two main evidence-based recommendations for practitioners. Firstly, potentiation strategies can be broadly used to acutely enhance athletic performances of both individual and team sport athletes.^{6,18,22} Secondly, the high inter-individual variability and inconsistency of the potentiation responses indicate the need for individualized approaches.^{23,24} Thus, conditioning protocols looking for potentiation responses can be implemented in the following settings:

- Testing: incorporated into standardized warm-up routines of assessment procedures intended to assess maximal athletic performance at different moments of the season. This may help limiting confounding effects arising from different warm-up protocols, thus facilitating a more consistent interpretation of the performance results.^{12,18,25}
- Training: as part of advanced programs in which the conditioning activity is paired with an unloaded explosive exercise (e.g., loaded squats + vertical jumps), and performed immediately after or following a brief rest interval (i.e. complex training or contrast training).²⁶

- Competition: incorporated into warm-up strategies completed at a precise timing prior to official competitions.²⁷

Although performance potentiation mean effects are commonly observed at a group level in well standardized conditions, inconsistent findings are reported between individuals even performing the same potentiating protocols.^{18,19,24} Therefore, practitioners should be aware that the individual characteristics of athletes may lead to different responses in terms of onset and magnitude of potentiation effects. Recurring evidence suggest a more individualized approach to optimize potentiation effects by manipulating the conditioning protocol variables with the identification of intensity, volume, and recovery time, for determining optimal loads adapted to the training background and sex of athletes.^{4,18,24,28} However, contrary to frequent claims, there is no evidence supporting the need of performing biomechanically similar exercises during conditioning protocols to benefit from potentiation responses.³

For meeting these objectives, the new taxonomy would be very helpful as it avoids the inadequate use of protocols in sport settings and populations different to those in which they exhibited effectiveness. Nonetheless, after identifying customized conditioning protocols from the scientific literature, practitioners would also need to test their efficacy in specific settings, after manipulating the conditioning protocol variables, on an individual basis. Meanwhile, sport physiologists would be able to better and more precisely identify the mechanisms associated with potentiation responses for augmenting the translational value of laboratory results to the field, following previous methodological recommendations.¹⁰ In this regard, reporting negative results would also be important to improve this process, given the existing bias of publishing more positive outcomes. Finally, sport scientists are suggested to examine and report the individual responses of well-characterized athletes (e.g., sex, training experience, competitive level, period of the season) in order to better identify the factors associated with both responder and non-responder groups.

197 **Conclusions**

198 We presented a novel taxonomy for the classification of potentiation in sport. This
199 taxonomy encompasses the identification of the conditioning activity, verification test, and
200 athletic population according to this formula: Post- [CONDITIONING ACTIVITY]
201 [VERIFICATION TEST] potentiation in [POPULATION]. This proposal may potentially
202 avoid erroneous identification of physiological attributes, which should be studied separately,
203 while favoring individualization and applicability of conditioning protocols in sport settings.
204 The use of PAP and PAPE would be valid, but assuming that both definitions could be
205 complementary in some cases.

References

1. Tillin NA, Bishop D. Factors modulating post-activation potentiation and its effect on performance of subsequent explosive activities. *Sport Med.* 2009;39(2):147-66. doi:10.2165/00007256-200939020-00004
2. Seitz LB, Haff GG. Factors Modulating Post-Activation Potentiation of Jump , Sprint , Throw , and Upper-Body Ballistic Performances : A Systematic Review with Meta-Analysis. *Sport Med.* 2015;46(2):231-240. doi:10.1007/s40279-015-0415-7
3. Beato M, Madruga-parera M, Piqueras-sanchiz F, Moreno-p V, Romero-rodriguez D. Acute Effect of Eccentric Overload Exercises on Change of Direction Performance and Lower-Limb Muscle Contractile Function. *J Strength Cond Res.* 2019;(36):1-7. doi:10.1519/JSC.0000000000003359
4. Wilson JM, Duncan NM, Marin PJ, et al. Meta-analysis of postactivation potentiation and power: Effects of conditioning activity, volume, gender, rest periods, and training status. *J Strength Cond Res.* 2013;27(3):854-859. doi:10.1519/JSC.0b013e31825c2bdb
5. Maloney SJ, Turner AN, Fletcher IM. Ballistic Exercise as a Pre-Activation Stimulus: A Review of the Literature and Practical Applications. *Sport Med.* 2014;44(10):1347-59. doi:10.1007/s40279-014-0214-6
6. Beato M, McErlain-Naylor SA, Halperin I, Iacono A Dello. Current evidence and practical applications of flywheel eccentric overload exercises as postactivation potentiation protocols: A brief review. *Int J Sports Physiol Perform.* 2020;15(2):154-161. doi:10.1123/ijsp.2019-0476
7. Vandenboom R. Modulation of skeletal muscle contraction by myosin phosphorylation. *Compr Physiol.* 2017;7(1):171-212. doi:10.1002/cphy.c150044
8. McGowan CJ, Pyne DB, Thompson KG, Rattray B. Warm-Up Strategies for Sport and Exercise: Mechanisms and Applications. *Sport Med.* 2015;45(11):1523-46.

doi:10.1007/s40279-015-0376-x

9. Cuenca-Fernández F, Smith IC, Jordan MJ, et al. Nonlocalized postactivation performance enhancement (PAPE) effects in trained athletes: A pilot study. *Appl Physiol Nutr Metab*. 2017;42(10):1122-1125. doi:10.1139/apnm-2017-0217
10. Blazevich AJ, Babault N. Post-activation Potentiation (PAP) versus Post-activation Performance Enhancement (PAPE) in Humans: Historical Perspective, Underlying Mechanisms, and Current Issues. *Front Physiol*. 2019;10:1359. doi:10.3389/FPHYS.2019.01359
11. Haiko Bruno Zimmermann, Brian R. MacIntosh JDP. Does post-activation potentiation (PAP) increase voluntary performance? *Appl Physiol Nutr Metab*. 2019;1-30. doi:10.1139/apnm-2019-0406
12. de Poli R de AB, Boullosa DA, Malta E de S, et al. Cycling Performance Enhancement After Drop Jumps May Be Attributed to Postactivation Potentiation and Increased Anaerobic Capacity. *J Strength Cond Res*. 2020;ahead of p(1).
13. Behm DG. Force maintenance with submaximal fatiguing contractions. In: *Canadian Journal of Applied Physiology*. ; 2004. doi:10.1139/h04-019
14. Boullosa D, Del Rosso S, Behm DG, Foster C. Post-activation potentiation (PAP) in endurance sports: A review. *Eur J Sport Sci*. 2018. doi:10.1080/17461391.2018.1438519
15. Rassier DE, MacIntosh BR. Coexistence of potentiation and fatigue in skeletal muscle. *Brazilian J Med Biol Res*. 2000. doi:10.1590/S0100-879X2000000500003
16. Sale DG. Postactivation potentiation: Role in human performance. *Exerc Sport Sci Rev*. 2002. doi:10.1097/00003677-200207000-00008
17. Low JL, Ahmadi H, Kelly LP, Willardson J, Boullosa D, Behm DG. Prior band-resisted squat jumps improves running and neuromuscular performance in middle-

256 distance runners. *J Sport Sci Med*. 2019;18(2):301-315.
 257 <http://europepmc.org/articles/PMC6543995>.

258 18. Boullosa D, Abad CCC, Reis VP, et al. Effects of Drop Jumps on 1000-m
 259 Performance Time and Pacing in Elite Male and Female Endurance Runners. *Int J*
 260 *Sports Physiol Perform*. 2020;1-4. doi:10.1123/ijsp.2019-0585

261 19. Boullosa DA, Abreu L, Beltrame LGN, Behm DG. The acute effect of different half
 262 squat set configurations on jump potentiation. *J Strength Cond Res*. 2013.
 263 doi:10.1519/JSC.0b013e31827ddf15

264 20. Brian J. Wallace, Shapiro R, Kelly L. Wallace Mark G. Abel Atbs. Muscular and
 265 Neural Contributions to Postactivation Potentiation. *J Strength Cond Res*.
 266 2019;33(3):615-625.

267 21. Anthi Xenofondos, Dimitrios Patikas, David M Kocaja, Tahayori Behdad, Eleni Bassa
 268 1, Eleftherios Kellis CK. Post-activation potentiation : the neural effects of post —
 269 Activation depression. *Muscle Nerve*. 2015;52(2):252-259. doi:10.1002/mus.24533

270 22. Dello Iacono A, Beato M, Halperin I. The Effects of Cluster-Set and Traditional-Set
 271 Postactivation Potentiation Protocols on Vertical Jump Performance. *Int J Sports*
 272 *Physiol Perform*. 2019;1-6. doi:10.1123/ijsp.2019-0186

273 23. Chaouachi A, Poulos N, Abed F, et al. Volume , intensity , and timing of muscle
 274 power potentiation are variable. *Appl Physiol Nutr Metab*. 2011;747:736-747.
 275 doi:10.1139/H11-079

276 24. Kobal R, Pereira LA, Kitamura K, Paulo AC. Post - Activation Potentiation : Is there
 277 an Optimal Training Volume and Intensity to Induce Improvements in Vertical Jump
 278 Ability in Highly - Trained Subjects ? by. *J Hum Kinet*. 2019;69(September):239-247.
 279 doi:10.2478/hukin

280 25. Doma K, Leicht AS, Boullosa D, Woods CT. Lunge exercises with blood-flow

restriction induces post-activation potentiation and improves vertical jump performance. *Eur J Appl Physiol*. 2020;(0123456789). doi:10.1007/s00421-020-04308-6

26. Patrick Cormier, Tomás T Freitas, Jacobo Á Rubio-Arias PEA. Complex and Contrast Training: Does Strength and Power Training Sequence Affect Performance-Based Adaptations in Team Sports? A Systematic Review and Meta-analysis. *J Strength Cond Res*. 2020;Epub Ahead. doi:10.1519/JSC.0000000000003493
27. Kilduff LP, Finn C V, Baker JS, Cook CJ, West DJ. Preconditioning Strategies to Enhance Physical Performance on the Day of Competition. *Int J Sports Physiol Perform*. 2013;8(6):677-681. doi:10.1123/ijsp.8.6.677
28. Maszczyk A, Zajac A, Stastny P. Optimizing Post Activation Potentiation for Explosive Activities in Competitive Sports. 2016;52(September):95-106. doi:10.1515/hukin-2015-0197

296 Table 1. Link between terminology and activity in potentiation studies.¹⁰

Term	Definition
Staircase potentiation	A progressive increase in twitch contractile response during repeated low-frequency stimulations.
Post-tetanic potentiation (PTP)	An increase in amplitude of twitch tension after a sustained muscle tetanic stimulation, generally at a high stimulation frequency.
Post-activation potentiation (PAP)	Augmentation of evoked twitch tension induced by voluntary activation of the muscle.
Post-activation performance enhancement (PAPE)	Enhancement of subsequent voluntary, rather than electrically evoked (twitch), force production, following high-intensity voluntary conditioning contraction(s).

297