UWS Academic Portal

The use of feed and water additives for live fish transport

Vanderzwalmen, Myriam; Eaton, Lewis; Mullen, Carrie; Henriquez, Fiona ; Carey, Peter; Snellgrove, Donna; Sloman, Katherine

Published in:
Reviews in Aquaculture

DOI:
[10.1111/raq.12239](https://doi.org/10.1111/raq.12239)

Published: 14/02/2019

Document Version
Peer reviewed version

[Link to publication on the UWS Academic Portal](#)

Citation for published version (APA):

Vanderzwalmen, M., Eaton, L., Mullen, C., Henriquez, F., Carey, P., Snellgrove, D., & Sloman, K. (2019). The use of feed and water additives for live fish transport. *Reviews in Aquaculture*, 11(1), 263-278.
<https://doi.org/10.1111/raq.12239>

General rights

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact pure@uws.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

The Use of Feed and Water Additives for Live Fish Transport

M. Vanderzwalmen^{1*}, L. Eaton², C. Mullen¹, F. Henriquez¹, P. Carey³, D. Snellgrove² &
K.A. Sloman¹

¹*School of Science & Sport, University of the West of Scotland, Paisley, PA1 2BE*

²*WALTHAM Centre for Pet Nutrition, Freeby Lane, Waltham-on-the-Wolds, LE14 4RT, U.K.*

³*Pets at Home, Epsom Avenue, Stanley Green Trading Estate, Handforth, Cheshire, SK9 3RN*

*Corresponding author: myriam.vanderzwalmen@uws.ac.uk

Abstract

The transport of live fish for aquaculture, either for food or as companion animals, presents a major issue for animal welfare. The stressors associated with live transportation are well documented with a focus on maintaining water quality during transport to reduce stress. Far less considered is our ability to enhance health and welfare during transport through the use of dietary and water additives prior to and during transport. With increasing interest in the use of plant essential oils as feed additives in food fish aquaculture and the increased availability of products claiming to alleviate stress in ornamental species, there is a need for scientific investigation into these potential welfare-promoting methods. Here we summarise current knowledge on the use of food additives, water conditioners, antibiotics, antimicrobials, and probiotics to promote fish health during transport. This review aims to highlight the gaps in our knowledge surrounding promising ways of promoting fish health during transport and to stimulate new research in this area.

Key Words: fish transport, stress, water quality, welfare management

Introduction

Fish production as a food source was valued at US\$160 billion in 2014 (FAO 2016) and the estimated worth of the ornamental fish industry (including wholesale, wages, retail sales, associated materials etc.) is approximately US\$15 billion (FAO 2011). In both aquaculture enterprises, fishes will be transported at some point in the production chain although the duration and the type of transport varies greatly between the two industries. Fishes destined for food are often transported by road to various facilities in open systems where continuous monitoring of water quality and access to fish during transport is possible (Van de Sande 1974; Berka 1986; Rimmer & Franklin 1997; Lekang 2007; Espinosa-Curiel *et al.* 2016). In contrast, ornamental fishes tend to be transported over greater distances in closed systems (e.g. in plastic bags shipped by aeroplane) (Braker 1974; Berka 1986; Swann 1992; Rimmer & Franklin 1997; Cole *et al.* 1999; Marine Aquarium Council 2001; Lim *et al.* 2007). While fish transport practices have been reviewed extensively, exact information on mortality during transport is particularly limited for the ornamental fish trade with estimated values ranging from a few per cent to greater than 80% (Rubec & Cruz 2005). High mortality is also reported during recovery from transport (Froese 1988; Sadovy 2002; Rubec & Cruz 2005).

Despite the differences in transport practice between food fishes and fishes destined for the pet industry, welfare challenges for transportation are very similar. Stressors encountered during transport include handling prior to transport, deterioration of water quality during transport and increased susceptibility to metabolic shock, stress, infection and disease after transport (Pickering *et al.* 1982; Portz *et al.* 2006; Sampaio & Freire 2016). There are several excellent reviews on how water quality and husbandry practices can

improve welfare (Berka 1986; Swann 1992; Cole *et al.* 1999; Crosby *et al.* 2005a, b; Lim *et al.* 2007; Harmon 2009), but far less scientific attention has focused on the potential to enhance welfare during transport over and above good husbandry practice. For example, a range of commercial and non-commercial products have been suggested to aid fish health during transport but there is a considerable absence of studies within the scientific literature surrounding these products. This review will firstly consider whether the stress of transport can be alleviated by feed or water treatment prior to transport, and secondly consider current knowledge surrounding water additives during transport.

Can we Prepare Fishes for Transportation Stress?

Dietary supplements are commonly used in aquaculture for a wide variety of reasons including stress reduction (Peng *et al.* 2013; Vallejos-Vidal *et al.* 2016), improving specific and non-specific immune resistance (Wang *et al.* 2006; Vallejos-Vidal *et al.* 2016), enhancing colouration (Pan & Chien 2009; Kouba *et al.* 2013) and increasing growth rate (Vallejos-Vidal *et al.* 2016). Therefore, prior to transport, there is the opportunity to use dietary supplements to enhance the immune system and improve stress tolerance (Volpatti *et al.* 1998; Lim *et al.* 2002; Rollo *et al.* 2006). Dietary supplements that have been tested in relation to transport in fishes include glucan, probiotics, ascorbic acid, carotenoids and herbal supplements.

Glucan is a polysaccharide which enhances non-specific immunity in fishes and reduces susceptibility to stress and the immunosuppressive effects of stress (Volpatti *et al.* 1998; Kim *et al.* 1999; Vallejos-Vidal *et al.* 2016). Rainbow trout (*Oncorhynchus mykiss*, Walbaum 1792) fed a diet with no (0%), low (0.1%), medium (0.5%) or high (1%) levels of

glucan for four weeks showed differences in non-specific immune responses to 2 h simulated transport stress (Volpatti *et al.* 1998). Following transport stress, lymphocytes decreased in the group fed 0% glucan, whereas in all the glucan-fed groups, lymphocytes, monocytes and neutrophils, increased demonstrating that glucan increases the non-specific immune response. Phagocytosis and respiration burst activity of cells were also higher in glucan-fed groups than in the 0% group. A sub-group of fish was fed the experimental diets for a further two months following transport to determine the long-term effect of glucan on the digestive tract. Prolonged exposure to glucan resulted in slight deterioration within the epithelial cells of the stomach and gut (Volpatti *et al.*, 1998) suggesting that prolonged ingestion of glucan may not be beneficial.

Probiotics are either a mono or mixed culture of live micro-organisms that are ingested and multiply in the gut of host organisms in order to improve the indigenous microflora (Havenaar *et al.* 1992; Cross, 2002). Probiotics are increasingly being considered as a safer and more environmentally-friendly alternative to antibiotics with their use in aquaculture increasing as a response to public demand for antibiotic-free fishes (Martínez Cruz *et al.* 2012). The desired effects of probiotics include improved larval development, growth promotion, stimulation of the immune system, pathogen and disease control, stress resistance and improved water quality (Martínez Cruz *et al.* 2012). Ambas *et al.* (2015) simulated transport of marron (*Cherax cainii*, Austin 2002) reared on marron commercial feed as a control, or control feed enriched with *Bacillus mycoides* (Flugge 1886). After 48 h, survival of marron fed the control diet was $93.3 \pm 2.8\%$ S.E. vs. 100% in the probiotic group. Marron in the probiotic group also had a higher intestinal bacterial population and total haemocyte count than the controls, indicating an improved immune status when fed probiotics. Although studies on the direct benefits of probiotics in alleviating transport stress

are limited, the advantages of probiotics in response to stressors relevant to transport have also been demonstrated. Rollo *et al.* (2006) reared sea bream (*Sparus aurata*, Linnaeus 1758) larvae with a mixture of *Lactobacillus fructivorans* (Charlton *et al.* 1934) and *Lactobacillus plantarum* (Orla-Jensen 1919) for either 20 or 42 days using rotifers (*Brachionus plicatilis*, Müller 1786) and/or *Artemia salina* (Linnaeus 1758) as a vector. At the end of the rearing period, 600 fry were exposed to 6.3 pH for 1 h. The probiotic group had lower levels of mortality and whole body cortisol. Hsp70 gene expression increased after exposure to low pH with the greatest increase in Hsp70 found in sea bream reared with probiotics.

Fishes cannot synthesise ascorbic acid (vitamin C) and rely on absorption through their food (Sales & Janssens 2003; Peng *et al.* 2013). Supplementing feed with ascorbic acid has been found to reduce mortality following a stressor (Lim *et al.* 2002). Peng *et al.* (2013) fed silver pomfret (*Pampus argenteus*, Euphrasén, 1788) a diet supplemented with ascorbic acid (L-ascorbyl- 2-polyphosphate, 35% ascorbic acid equivalent at a concentration of 100, 450 or 800 mg ascorbic acid kg⁻¹ diet) for 9 weeks. Silver pomfret were then transported in darkened plastic bags for 4 h. Diet supplementation with ascorbic acid significantly reduced serum cortisol and glucose levels as well as mortality, indicating that high ascorbic acid successfully reduced stress associated with transport.

Abreu *et al.* (2014) fed wild-caught pencilfish (*Nannostomus trifasciatus*) one of four experimental diets with 0%, 0.01%, 0.1% or 0.5% beta 1,3 glucan added for 7 days. The fish were then fasted and transported for 24 h. The addition of beta 1,3 glucan reduced the net loss of K⁺ during the first 3 h of transport. The addition of 0.5% beta 1,3 glucan to the feed increased the net influx of Na⁺ between 3 and 12 hours of transport.

Combinations of ascorbic acid and glucan as supplements have been trialled in relation to transport stress. Barros *et al.* (2014) supplemented the diet of Nile tilapia (*Oreochromis niloticus*, Linnaeus 1758) with a basal diet of 125 mg kg⁻¹ of ascorbic acid (BD) followed by a diet containing 0.1% β -glucan and 600 mg kg⁻¹ ascorbic acid (GD). The fish were reared on BD for 20 days, after this period the fish were fed GD for variable durations (7, 15, 30, or 45 days) prior to undergoing transport stress. Fish from all four treatments were transported for 4 h in 100 l net cages within a 600 l fish transport tank. During recovery from the transport, fish fed 0.1% β -glucan and 600 mg kg⁻¹ ascorbic acid for 7 days had the highest cortisol concentrations and greatest increase in red blood cells and haemoglobin. However, longer exposure to β -glucan and ascorbic acid reduced plasma cortisol and after 72 h, cortisol had returned to baseline in Nile tilapia fed this diet for longer than 7 days. As β -glucan and high levels of ascorbic acid were not tested separately, it is unclear which supplement provided the observed improvements or whether it was the two combined.

Organisms undergoing stress can experience a shortage of oxygen at the cellular level causing abnormal oxidative reactions in the aerobic metabolic pathway and the generation of reactive oxygen species (ROS) (Rånby & Rabek 1978; Pan *et al.* 2010) which if not inactivated can cause oxidative damage to lipids, proteins, carbohydrates and nucleotides (Yu 1994; Chew 1995; Halliwell & Gutteridge 2015). Carotenoids are pigment molecules with known antioxidant properties that are important to animal health by inactivating free radicals. Fishes cannot synthesise carotenoids *de novo* and must acquire them through their diet. In food fish aquaculture, diets are often supplemented with carotenoids to improve fillet colouration and to reduce oxidative stress. The use of carotenoids, in particular astaxanthin, as a dietary supplement in aquaculture has been extensively reviewed and found to improve

stress resistance (Johnson *et al.* 1980; Lorenz & Cysewski 2000; Pan *et al.* 2003, 2010; Higuera-Ciapara *et al.* 2006; Pan & Chein 2009; Jagruthi *et al.* 2014; Lim *et al.* 2017). Dietary carotenoids such as astaxanthin and β -carotene have been shown to help fish cope with stressors that may act as components of transport stress (e.g. hypoxia, Pan *et al.* 2010; high ammonia, Pan *et al.* 2011). Therefore, the potential benefits of carotenoids as dietary supplements to prepare fish for transport stress warrants future study.

In aquaculture, plant extracts are mostly used as feed supplements, although some may be added directly to the water (see **Additives During Transport** below). In 2011 the total global herbal drug market was estimated at US \$62 billion and is predicted to grow to US \$5 trillion by 2050 with an annual growth rate of 5-15% (Harikrishnan *et al.* 2011). Plant-derived products can be effective as antioxidants, growth promoters, appetite stimulators, immune stimulants and stress reducers along with having anti-inflammatory and anti-carcinogenic properties (Citarasu 2010; Harikrishnan *et al.* 2011; Merlini *et al.* 2014; Bulfon *et al.* 2015). The use of plant essential oils as dietary supplements in aquaculture was recently reviewed (Sutuli *et al.* 2017) but to our knowledge only two studies have considered the effects of plant extracts on stress tolerance during transportation through administration as a dietary supplement. Turmeric (*Curcuma longa*, Linnaeus 1758) has a variety of documented medicinal properties (anti-inflammatory, Araújo & Leon 2001; immunostimulant, Chattopadhyay *et al.* 2004; antioxidant, Luthra *et al.* 2001, Saccol *et al.* 2016; anaesthetic, Saccol *et al.* 2016; anti-microbial, Luthra *et al.* 2001; anti-parasitic, Araújo & Leon 2001). Supplementing the diet of juvenile yellow tail tetra (*Astyanax aff. bimaculatu*, Linnaeus 1758) with turmeric for 60 days before a simulated 24 h transport reduced mortality, plasma lactate and plasma glucose concentrations compared to controls (Ferreira *et al.*, 2017). *Aloe vera* (Burman 1768) is commonly used in humans for a wide range of medicinal properties

184 (e.g. Vázquez *et al.* 1996; Reynolds & Dweck 1999; Vogler & Ernst 1999; Choi *et al.* 2001;
185 Choi & Chung 2003; Mahor & Ali 2016) but little is known about the benefits of using *A.*
186 *vera* as a dietary supplement in aquaculture. Zanuzzo *et al.* (2017) administered one of four
187 diets (0%, 0.5%, 1% or 2% *A. vera*) for 10 days prior to a 4 h transport of juvenile pacu
188 (*Piaractus mesopotamicus*, Holmberg 1887). Immediately following transport, the pacu were
189 divided into three sub-groups: a non-injected control group; a buffer injected group and a
190 group injected with inactivated *Aeromonas hydrophila* (Chester 1901) to stimulate their
191 immune system. This is particularly relevant to transport stress studies as fishes can become
192 highly susceptible to bacterial and viral infections during transport (Yanong 2003; Crosby *et*
193 *al.* 2005b). On arrival and after 24 h recovery, non-injected *A. vera* fed groups had
194 significantly higher cortisol concentrations than the non-injected control fish. However, in the
195 *A. hydrophila* injected fish, the cortisol levels of the fish fed 1% *A. vera* were significantly
196 lower than in the other groups. Immediately after transport, leukocyte respiratory burst was
197 higher in all (injected and non-injected) *A. vera* fed fish than in the control fish but no
198 significant difference was found between the treatments after 24 h of recovery. The
199 haemolytic activity of the complement system was significantly higher in the *A. vera* fed fish
200 on arrival and after recovery than in the control fish. The results indicate that *A. vera*
201 improved the immune system of juvenile pacu after transport and improved the stress
202 recovery of infected individuals. No explanation for why *A. vera* slowed recovery rate of the
203 non-injected juvenile pacu yet improved the recovery rate of the injected group was given.
204 Further research into the use of herbal dietary supplements in relation to transport of fishes is
205 clearly needed.

206

207 **The Use of Additives During Transport**

208

209 There are many reviews that have considered the importance of good water quality during
210 live fish transport (Berka 1986; Swann 1992; Cole *et al.* 1999; Crosby *et al.* 2005a, 2005b;
211 Lim *et al.* 2007; Harmon 2009) and a variety of products can be added to the water to
212 maintain water quality or alleviate the problems of waste products (e.g. pH buffers, zeolites,
213 AmQuel[®], nitrifying bacteria). Even when water quality is optimal for the duration of
214 transport, there are many other stressful factors (e.g. handling, high loading density and
215 crowding) (Pickering *et al.* 1982; Portz *et al.* 2006; Sampaio & Freire 2016); far less research
216 has focused on whether it is possible to add compounds to the water to alleviate the
217 physiological effects of stress. As demonstrated above, the benefits of dietary supplements
218 for improving health and welfare during transport may require many weeks of preparation;
219 the addition of compounds to the water for the period of transportation requires far less
220 advance planning. Some products such as water conditioners and salt are added directly to the
221 transport water along with fishes while others may require exposure for a short time
222 immediately prior to transport (e.g. anaesthetics). The promotion of physiological well-being
223 through addition of compounds to the water can occur *via* different physiological
224 mechanisms such as sedation, protection of mucus integrity and disease prevention. While
225 the use of traditional water additives such as salt and synthetic anaesthetics have been well
226 explored, the use of plant extracts as water conditioners and antimicrobial agents has received
227 less attention. Here we will briefly discuss the more traditional water additives but will focus
228 primarily on the use of novel and emerging water additives.

229

230 **Managing Stress by Sedation**

Anaesthetics are one of the most commonly used additives in the transport of fish (Lim *et al.* 2003; Harmon 2009; Cupp *et al.* 2017). Anaesthesia is defined as “a state caused by an applied external agent resulting in a loss of sensation through depression of the nervous system” (Akerman *et al.* 2005). Sedation can lower a fish’s metabolic rate (Ross & Ross 2008) resulting in improved water quality (Pattanasiri *et al.* 2016), lower levels of stress (Ims 2011) and often allows transportation at higher loading densities (Cupp *et al.* 2017). Frequently used synthetic anaesthetics, such as MS-222 (ethyl 3-aminobenzoate methanesulfate) and benzocaine, have been widely reviewed elsewhere (Ross *et al.* 2008; Carter *et al.* 2011; Javahery *et al.* 2012; Readman *et al.* 2013; Husen & Sharma 2014) and will not be discussed in detail here. Generally, synthetic anaesthetics initially induce stress before having a delayed stress-reduction effect (Ims 2011; Readman *et al.* 2013). An alternative to sedation through synthetic compounds is to use natural compounds such as essential oils.

The most commonly used essential oil is clove oil (*Syzygium aromaticum*, Linnaeus 1758). Clove oil has been extensively reviewed for its use as an anaesthetic (e.g. Javahery *et al.* 2012) but many other less researched essential oils have been considered for use during fish transport. Most studies have considered the use of essential oils during transport by adding them to the water at set concentrations. *Condalia buxifolia* (Reisseck 1861) methanolic extract induced sedation in silver catfish (*Rhamdia quelen*) for 6 h and during transport improved survival, water quality and reduced ion loss (Becker *et al.* 2013). Salbego *et al.* (2015) also found similar results when they sedated non-starved silver catfish with methanolic extract of *C. buxifolia* prior to transport and added *C. buxifolia* solution to the transport water. Addition of *C. buxifolia* improved water quality, reduced total ammonia

nitrogen (TAN) levels, slowed metabolism and reduced net ion efflux. Lipoperoxidation and carbonylation of proteins decreased in silver catfish transported in *C. buxifolia*, demonstrating less ROS production. Low concentrations (0.5-10 $\mu\text{l l}^{-1}$) of *C. buxifolia* induced fast sedation and higher doses did not cause harmful effects suggesting it is safe to use to induce sedation (Becker *et al.* 2013).

The use of essential oils in the water is summarised in Table 1. Interestingly, Pattanasiri *et al.* (2016) tested the release rate of low-density polyethylene (LDPE) bags coated with clove oil over a 48 h period. After the initial 2 h, the release rate of clove oil was almost constant at 12 mg l^{-1} when the bag contained 75 ml water and 14 mg l^{-1} when the bag contained 150 ml water. These levels induced sedation in Siamese fighting fish (*Betta splendens*, Regan 1910) but not anaesthesia. Survival of Siamese fighting fish transported in the clove oil-coated bags was significantly higher, with lower ammonia concentrations and higher dissolved oxygen than in control un-coated bags. Siamese fighting fish did not appear to experience any detrimental effects of prolonged clove oil exposure.

In summary, several essential oils are effective in inducing sedation in fish, however, most essential oils have only been tested at one or a few concentrations and in a single species; few have been tested for use in fish transport. The mechanism of action can vary between essential oils and in the most part is poorly understood. It is known that different species have different aversive reactions to anaesthetics, so there is a need to further explore the use of essential oils on individual species of interest (Javahery *et al.* 2012; Husen & Sharma 2014; Chambel *et al.* 2015). With the increased commercialisation of essential oils, or components of, for use in fish husbandry (e.g. AQUI-S; <http://aquatactics.com/aqui-s->

20e/), the potential for better, more refined chemicals for improving fish welfare during transport compared to more traditional synthetic chemicals seems likely.

Maintaining Mucus Integrity

The epidermal layer of fishes excretes protective mucus that serves as a barrier to the external environment (Harnish *et al.* 2011; Ottesen & Olafsen 1997; Shephard 1994). During transport and handling, fishes can lose their protective mucus, resulting in an increased risk of injuries (Harnish *et al.* 2011). Mucus loss can result in disturbed osmoregulation, loss of scales, skin damage, and bacterial, fungal and parasitic diseases (Wedemeyer 1996). When fishes are stressed or transported in large numbers within a single bag, the risks of losing mucus increases as fish are more likely to come into contact with the bag or other individuals. Protecting epidermal mucus can result in a higher quality of fish at arrival in terms of physical health, reduced risk of infection and improved aesthetic appearance.

Water conditioners usually refer to compounds added directly to the transport water of fish to reduce stress by means other than sedation. Water conditions may be either pure herbal extracts or combined with other products into a commercially available product. A review on the use of polymer-based water conditioners to reduce handling-related injuries (Harnish *et al.* 2011) identified three studies (summarised in Table 2). Wedemeyer (1996) presented findings of user surveys and husbandry procedures finding that polymer-based water conditioners reduce mortality.

Stress Coat[®] (API Aquarium Pharmaceuticals Inc., n.d.) is a commercially available water conditioner which is recommended for addition to water during transport and to tank water during other potentially stressful procedures. Several studies have used Stress Coat[®] in

husbandry procedures (Earley *et al.* 2006; Colburn *et al.* 2008; Harnish *et al.* 2011; Wong *et al.* 2015), however, much of the information on Stress Coat[®] is not in the peer-reviewed literature. Snellgrove *et al.* (2007) found that water cortisol levels excreted by goldfish exposed to Stress Coat[®] were lower following netting compared to goldfish netted with no Stress Coat[®] exposure. Edmonds (2016) found that Stress Coat[®] exposure during transport did not reduce excretion of cortisol but reduced conspecific aggression in guppies post-transport.

The main components of Stress Coat[®] (manufactured by Mars Fishcare) are *Aloe barbadensis* Mill (also known as *A. vera*) (1-10%), water (>80%), polyvinylpyrrolidone (PVP) and other non-hazardous ingredients (trade protected) (1-10%) (Mars Fishcare Inc. 2014). Most of the limited peer-reviewed research carried out directly on *A. vera* focus on it as a dietary supplement used to alleviate stressors not relevant to the transport of fish (*e.g.* Dotta *et al.* 2014; Gabriel *et al.* 2015a,b; Kim *et al.* 1999; Taiwo *et al.* 2005; Zanuzzo *et al.* 2015a,b). One study looked at *A. vera* as a dietary supplement prior to transport (Zanuzzo *et al.* 2017; see above) and an additional study by the same group used *A. vera* in the water during transport of fish. Zanuzzo *et al.* (2012) dissolved *A. vera* powder (concentrations: 0, 0.02, 0.2 and 2 mg l⁻¹) in the transport water of matrinxã (*Brycon amazonicus*, Spix and Agassiz 1829). *Aloe vera* increased the activity of the immune system, by enhancing the respiratory activity of leukocytes, in matrinxã following handling but the effects of *A. vera* were no longer apparent at the end of the 4 h transport. *Aloe vera* has potential to improve the condition of fish during transport procedures but more research is needed to gain a better understanding of the possible benefits and the best method of administration.

Sung *et al.* (2012) tested another patented product Pro-Tex[®] (Bradán Limited, a soluble variant of TEX-OE[®]) to determine whether it improved the resistance of juvenile carp

329 (*Cyprinus carpio*, Linnaeus 1758) to high ammonia levels. Pro-Tex[®] contains an extract of
330 the prickly pear cactus (*Opuntia ficus indica*, (L.) Miller 1925) which increases heat-shock
331 protein expression in humans and fishes (Wiese *et al.* 2004; Roberts *et al.* 2010; Sandilands
332 *et al.* 2010). Exposure of carp to Pro-Tex[®] (2 µl 50 l⁻¹ water) for 2 h increased survival from
333 50 to 95% and 0 to 20% when exposed to 5.92 mg l⁻¹ and 14.21 mg L⁻¹ of NH₃ respectively
334 for 1 h. Sung *et al.* (2012) also found that Pro-Tex[®] increased expression of heat shock
335 protein (Hsp70) in gill and muscle tissue of carp which may suggest potential benefits of Pro-
336 TEX[®] for fish transport. Hales *et al.* (1990) evaluated the injury-preventing capacity of a
337 water-soluble gel-coating (composed of antibiotics and non-disclosed pharmaceutical
338 components) applied to the hands of fish handlers. The unexpected results showed that spot
339 croaker (*Leiostomus xanthurus*, Lacepède 1802) handled with gel-coating had a higher
340 mortality than fish handled by collectors with non-coated hands. The authors hypothesised
341 that the composition of the gel was different from that of the mucus of the fish as mucus
342 mostly consists of protein along with lipids, carbohydrates and nucleic acids (Al-Hassan *et al.*
343 1982). Austin *et al.* (2009), published a non-peer reviewed report about the use of
344 ULTIMATE[®] (AquaScience Technologies) in the transport water of koi carp. ULTIMATE[®]
345 has two main components, the ingredients for ClorAm-X[®] (the original AmQuel[®])
346 (composed of sodium hydroxymethanesulfonate, AquaScience Technologies) and the
347 ingredients for Stress-X[®] (the original NovAqua[®]) (water, sodium thiosulfate, buffers,
348 electrolytes, proprietary synthetic polymer formulation and preservatives, Aquarium
349 Solutions). In addition, ULTIMATE[®] contains a dechloraminating agent, electrolytes
350 (including calcium, sodium and chloride ions), a polymer system and product stabilizers
351 (Austin *et al.* 2009). After 8 h of simulated sealed transport, the koi transported in
352 ULTIMATE[®] had shorter recovery times than the control fish indicated by accelerated
353 reduction in mucosal levels of haemoglobin. Unfortunately, sample size was not sufficient for

definitive results, and in addition, the methods used to detect levels of haemoglobin in the mucus were not precise enough. No published peer-reviewed paper was found that used ULTIMATE[®] in fish transport, however it is an area worth investigating based on the observations made by Austin *et al.* (2009). Despite many studies using substances such as Stress Coat[®], Novaqua[®] and Polyqua[®], little peer-reviewed information is available on the efficacy of these substances, particularly in maintaining mucus integrity.

Disruption of epidermal mucus can cause many detrimental effects including disruption to osmoregulation. The stress of transport itself can also cause changes in osmoregulation (Barton & Iwama 1991; Baldisserotto *et al.* 2007) and so during transport of freshwater species, salt (NaCl) can be added to the water to reduce the difference between the internal osmolality of the fish and that of its environment thereby reducing physiological workload required to maintain homeostasis (Nikinmaa *et al.* 1983). Table 3 gives examples of studies that have investigated the effects of NaCl addition during transport. It is clear that for some species, the addition of NaCl during transport may be beneficial but given the range of salinities that different species inhabit and differences in osmoregulatory capacity from stenohaline to euryhaline, the use of NaCl as an additive will always be very species and life-stage specific. Additionally, Tacchi *et al.* (2015) compared the skin morphology of non-transported rainbow trout to that of rainbow trout transported in fresh or salt water (5 g NaCl l⁻¹) using electron microscopy. Fish transported in salt water had a thin layer of mucus whereas fish transported in fresh water had a thick deposit of mucus. It was suggested that the addition of NaCl slowed down the release of mucus from goblet cells. A ~50-fold increase in skin-associated bacteria in fish transported in fresh water was seen compared to a ~10-fold increase in the salt water group (Tacchi *et al.* 2015). While salt water transported fish had a thinner mucus layer, the mucus layer of these fish appeared to be in better condition showing

that NaCl also has the potential to reduce subsequent stress caused by bacteria and improve skin mucus condition.

Prevention of Stress-Related Diseases

When fishes experience high levels of stress they become more susceptible to bacterial diseases, which can result in higher mortality (Yanong 2003, Crosby *et al.* 2005b). To prevent proliferation of bacteria in the transport water while the fish's immune system is weakened, antibiotics are sometimes used. Amend *et al.* (1982) tested the efficacy of several antibiotics (kanamycin, gentamicin, chloramphenicol, streptomycin, neomycin and furazolidone, each at 20 mg l⁻¹), the antiseptic acriflavine (10, 20, and 100 mg l⁻¹), the disinfectant chlorine dioxide at 20 mg l⁻¹, and the antimicrobial methylene blue (10 and 100 mg l⁻¹). The tests were done by simulating 48 h transport with southern platy (*Xiphophorus maculatus*, Günther 1866) and checking the efficacy of each substance to control bacteria levels. Kanamycin and gentamicin were found to be toxic to the fish at these concentrations. Methylene blue, chlorine dioxide, furazolidone, and acriflavine did not prevent bacteria growth. Acrafiavine at 100 mg l⁻¹, chloramphicol and streptomycin controlled bacteria levels, but caused mortality in the fish. Only neomycin was effective against bacteria and safe for the fish. Antibiotic resistance in ornamental fishes has been studied since the late 1970s and the literature shows that resistance emerges when new antibiotics become widely available (del Rio-Rodriguez & Turnbull 2002; Rose *et al.* 2013; Trust & Whitby 1976; Verner-Jeffreys *et al.* 2009). A study on ornamental fishes by Rose *et al.* (2013) found that the most effective antibiotics (cefotaxime and kanamycin) were effective against only 45% and 44% of their target bacteria respectively, and of these, 16% and 35% had developed resistance. In this study, nine bacteria were resistant to all the tested antibiotics, only one showed no

resistance at all. Dixon *et al.* (1999) found similar results in a study looking at bacterial resistance in fish imported from Singapore. Over 50% of the bacteria isolated were resistant to 7 out of 12 tested antibiotics. In addition to increased resistance, antibiotics can cause increased levels of stress in fishes. Cururu stingray (*Potamotrygon cf hystrix*, Müller and Henle 1841) transported in water containing tetracycline (200 mg l⁻¹) for 24 h had elevated corticosterone levels after 12 h compared to controls (Brinn *et al.* 2012). The administration of antibiotics during transport is also problematic because the use of antibiotics is strictly monitored and regulated in many countries (Cole *et al.* 1999; Crosby *et al.* 2005b; Brinn *et al.* 2012).

Probiotics can be added to transport water to improve water quality and reduce stress arising from low quality water. Efinol[®]L is a commercial probiotic product containing *Bacillus subtilis* (Ehrenberg 1835), *Bacillus licheniformis* (Weigmann 1898), *Lactobacillus acidophilus* (Moro 1900) and *Saccharomyces cerevisiae* (Hansen 1883); it also comprises amino acids, vitamins, minerals, free-flow and anti-caking agents (calcium carbonate and silica). Marbled hatchetfish (*Carnegiella strigata*, Günther 1864) were transported in a water solution containing 10 mg l⁻¹ probiotic Efinol[®]L. After 24 h of transport, water containing the probiotic solution had higher dissolved oxygen levels and lower ammonia concentrations (Gomes *et al.* 2008). Fish in the control group had higher body cortisol levels and higher efflux of Na⁺ and K⁺. The reduced stress levels seen in marbled hatchetfish treated with probiotics could be attributed to either a direct effect of Efinol[®]L on physiology or an indirect effect of improved water quality. Using a similar protocol, Gomes *et al.* (2009) transported cardinal tetra (*Paracheirodon axelrodi*, Schultz 1956) in water containing Efinol[®]L. The addition of Efinol[®]L resulted in a higher survival, higher water alkalinity and lower total ammonia in the water. The cortisol levels of the cardinals in the Efinol[®]L group were

significantly lower after transport compared to control fish. Efinol[®]L is not the only probiotic to have been added directly to the water. Zink *et al.* (2011) transported yellowfin tuna (*Thunnus albacares*, Bonnaterre 1788) yolk sac larvae one day post-hatching for 24 h. In the probiotic treatment, 300 ml of EcoAqua[®] (108 colony-forming units ml⁻¹ in a mix of *B. subtilis*, *B. licheniformis*, *B. megaterium* (Bary 1884), and *B. laterosporous* (Laubach 1916) EcoMicrobials LLC, Miami, Florida) was added to the water. No difference in survival was recorded between the treatments although the water quality of the bags containing probiotics was greater than in the control bags (lower pH, lower TAN and higher dissolved oxygen). Although the mechanisms of effect of these probiotic solutions are not yet fully understood, and it is unclear whether there are benefits related to reduced disease susceptibility, the positive effects seen in this limited number of studies suggest that more widespread investigation into the addition of probiotics during transport is warranted.

Concluding Remarks

There is a growing market for novel compounds which can be administered either *via* the diet or water to alleviate stress in fishes with the aim of increasing welfare. Perhaps driven in part by an increasing desire of consumers (either of food fishes or pet fishes) to purchase products which have not been exposed to synthetic chemicals, many of the emerging products are based on natural compounds or enhancing natural processes. While there is growing evidence that some of these compounds can improve welfare, much of the evidence remains anecdotal and the mechanisms of effect have been overlooked. Dietary supplements such as glucan, ascorbic acid, carotenoids, herbal supplements and probiotics may have the potential to reduce stress and mortality during transport but far more research is required to understand the capabilities of these supplements. Several commercially available water

conditioners have been considered in relation to transport stress, but there is a significant lack of peer-reviewed publications and publically available data on the testing of these products. Although, the process is not fully understood, probiotics can also be effective in reducing stress and mortality when added directly into transport water but as yet susceptibility to disease agents following such treatment remains unexplored. In order to enhance the welfare of fishes transported within aquaculture there is an urgent need to explore these emerging areas.

References

- Abasali H, Mohamad S (2010) Effects of using the *Valeriana officinalis* extract during transportation of swordtail, *Xiphophorus helleri*. *Journal of Animal and Veterinary Advances* **9**: 2377–2381.
- Abreu JS, Brinn RP, Gomes LC, McComb DM, Baldisserotto B, Zaiden SF, Urbinati EC, Marcon JL (2014) Effect of beta 1, 3 glucan in stress responses of the pencilfish (*Nannostomus trifasciatus*) during transport within the rio Negro basin. *Neotropical Ichthyology* **12**(3):623-628.
- Ackerman PA, Morgan JD, Iwama GK (2005) Anesthetics. *Supplement to the CCAC guidelines on: the care and use of fish in research, teaching and testing*. Available at: https://www.ccac.ca/Documents/Standards/Guidelines/Add_PDFs/Fish_Anesthetics.pdf

479 Al-Hassan JM, Thomson M, Criddle RS (1982) Composition of the proteinaceous gel
 480 secretion from the skin of the Arabian Gulf catfish (*Arius thalassinus*). *Marine Biology*
 481 **70**(1): 27–33. doi: 10.1007/BF00397293.

482

483 Ambas I, Fotadar R, Buller N (2015) Survival and Immunity of Marron *Cherax cainii*
 484 (Austin, 2002) Fed *Bacillus mycoides* Supplemented Diet under Simulated Transport.
 485 *Journal of Aquaculture Research & Development* **6**(390): 2. doi: 10.4172/2155-
 486 9546.1000390.

487

488 Amend DF, Croy TR, Goven, BA, Johnson KA, McCarthy DH (1982) Transportation of Fish
 489 in Closed Systems: Methods to Control Ammonia, Carbon Dioxide, pH, and Bacterial
 490 Growth. *Transactions of the American Fisheries Society* **111**(5): 603–611. doi:
 491 10.1577/1548-8659(1982)111<603:TOFICS>2.0.CO;2.

492

493 Araújo C, Leon L (2001) Biological activities of *Curcuma longa* L. *Memórias do Instituto*
 494 *Oswaldo Cruz. Fundação Oswaldo Cruz* **96**(5): 723–728. doi: 10.1590/S0074-
 495 02762001000500026.

496

497 Austin K, Janas T, Pattist K, Rice K (2009) Transporting Koi. 1–69. Available at:
 498 <file:///W:/article/artemia/Transporting Koi.pdf>

499

500 Azambuja CR, Mattiazzi J, Riffel APK, Finamor IA, Garcia LO, Heldwein CG, Heinzmann
 501 BM, Baldisserotto B, Pavanato MA, Llesuy SF (2011) Effect of the essential oil of
 502 *Lippia alba* on oxidative stress parameters in silver catfish (*Rhamdia quelen*) subjected
 503 to transport. *Aquaculture* **319**(1–2): 156–161. doi: 10.1016/j.aquaculture.2011.06.002.

504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527

Baldiasserotto B, Romero JMM, Kapoor BG eds (2007) Fish Osmoregulation. *Science Publishers*. doi: 10.1086/598309.

Barros MM, Falcon DR, De Oliveira OR, Pezzato LE, Fernandes AC, Guimarães IG, Fernandes A, Padovani CR, Sartori MMP (2014) Non-specific immune parameters and physiological response of Nile tilapia fed β -glucan and vitamin C for different periods and submitted to stress and bacterial challenge. *Fish and Shellfish Immunology* **39**(2): 188–195. doi: 10.1016/j.fsi.2014.05.004.

Barton BA, Iwama GK (1991) Physiological changes in fish from stress in aquaculture with emphasis on the response and effects of corticosteroids. *Annual Review of Fish Diseases* **1**:3–26. doi: 10.1016/0959-8030(91)90019-G.

Becker AG, da Cunha MA, de Oliveira Garcia L, Zeppenfeld CC, Parodi TV, Maldaner G, Morel AF, Baldiasserotto B (2013) Efficacy of eugenol and the methanolic extract of *Condalia buxifolia* during the transport of the silver catfish *Rhamdia quelen*. *Neotropical Ichthyology* **11**(3): 675–681. doi: 10.1590/S1679-62252013000300021.

Becker AG, Parodi TV, Zeppenfeld CC, Salbego J, Cunha MA, Heldwein CG, Loro VL, Heinzmann BM, Baldiasserotto B (2015) Pre-sedation and transport of *Rhamdia quelen* in water containing essential oil of *Lippia alba*: metabolic and physiological responses. *Fish Physiology and Biochemistry* **42**(1): 73–81. doi: 10.1007/s10695-015-0118-x.

528 Berka R (1986) The transport of live fish. A review. *Rome: Food and Agriculture*
529 *Organization of the United Nations*. Available at:
530 http://www.flyingsharks.eu/literature/fao_iata/Transport_of_live_fish_FAO.pdf.
531

532 Braker WP (1974) Collecting and transport of Caribbean marine fishes. *International Zoo*
533 *Yearbook* **14**: 40–43.
534

535 Brinn RP, Marcon JL, McComb DM, Gomes LC, Abreu JS, Baldisseroto B (2012) Stress
536 responses of the endemic freshwater cururu stingray (*Potamotrygon cf. hystrix*) during
537 transportation in the Amazon region of the Rio Negro. *Comparative Biochemistry and*
538 *Physiology - A Molecular and Integrative Physiology* **162**(2): 139–145. doi:
539 10.1016/j.cbpa.2011.07.004.
540

541 Bulfon C, Volpatti D, Galeotti M (2015) Current research on the use of plant-derived
542 products in farmed fish. *Aquaculture Research* **46**(3): 513–551. doi: 10.1111/are.12238.
543

544 Carneiro PCF, Urbinati EC (2001) Salt as a stress response mitigator of matrinxã, *Brycon*
545 *cephalus* (Günther), during transport. *Aquaculture Research* **32**(4): 297-304.
546

547 Carter KM, Woodley CM, Brown RS (2011) A review of tricaine methanesulfonate for
548 anesthesia of fish. *Reviews in Fish Biology and Fisheries* **21**(1): 51–59. doi:
549 10.1007/s11160-010-9188-0.
550

551 Chambel J, Pinho R, Sousa R, Ferreira T, Baptista T, Severiano V, Mendes S, Pedrosa R
 552 (2015) The efficacy of MS-222 as anaesthetic agent in four freshwater aquarium fish
 553 species. *Aquaculture Research* **46**(7): 1582–1589. doi: 10.1111/are.12308.
 554
 555 Chattopadhyay I, Biswas K, Bandyopadhyay U, Banerjee R (2004) Turmeric and curcumin:
 556 biological actions and medicinal applications. *Current Science* **87**(1): 44–50. Available
 557 at: <http://repository.ias.ac.in/5196/1/306.pdf> (Accessed: 21 March 2017).
 558
 559 Chew BP (1995) Antioxidant vitamins affect food animal immunity and health. *The Journal*
 560 *of nutrition* **125**(6 Suppl): 1804–1808. Available at:
 561 <http://www.ncbi.nlm.nih.gov/pubmed/7782950> (Accessed: 8 February 2017).
 562
 563 Choi S, Chung MH (2003) A review on the relationship between *Aloe vera* components and
 564 their biologic effects. *Seminars in Integrative Medicine* **1**(1): 53–62. doi:
 565 10.1016/S1543-1150(03)00005-X.
 566
 567 Choi SW, Son BW, Son YS, Park YI, Lee SK, Chung MH (2001) The wound-healing effect
 568 of a glycoprotein fraction isolated from *Aloe vera*. *British Journal of Dermatology*
 569 **145**(4): 535–545. doi: 10.1046/j.1365-2133.2001.04410.x.
 570
 571 Chow P, Chen T, Teo L (1994) Oxygen consumption rates of *Trichogaster trichopterus*
 572 (Pallas) in relation to starvation, size and sex. *The Third Asian Fisheries Forum* (ed. by
 573 LM Chou, AD).
 574

575 Citarasu T (2010) Herbal biomedicines: A new opportunity for aquaculture industry.
576 *Aquaculture International* **18**(3): 403–414. doi: 10.1007/s10499-009-9253-7.
577
578 Colburn HR, Walker AB, Berlinsky DL, Nardi GC (2008) Factors affecting survival of cobia,
579 *Rachycentron canadum*, during simulated transport. *Journal of the World Aquaculture*
580 *Society* **39**(5): 678–683. doi: 10.1111/j.1749-7345.2008.00205.x.
581
582 Cole B, Tamaru CS, Bailey R, Brown C, Ako H (1999) Shipping practices in the ornamental
583 fish industry. 1-22. Waimanalo, HI: Center for Tropical and Subtropical Aquaculture.
584
585 Cooke SJ, Schreer J, Wahl D, Philipp D (2002) Physiological impacts of catch-and-release
586 angling practices on largemouth bass and smallmouth bass. *American Fisheries Society*
587 *Symposium* **31**: 489–512.
588
589 Crosby TC, Hill JE, Martinez CV, Watson CA, Pouder DB, Roy PE (2005a) On-Farm
590 Transport of Ornamental Fish. *University of Florida Institute of Food and Agricultural*
591 *Sciences Extension Publication* **FA119**.

592

593 Crosby TC, Hill JE, Martinez CV, Watson CA, Pouder DB, Roy PE (2005b) Preparation of
594 Ornamental Fish for Shipping. *University of Florida Institute of Food and Agricultural*
595 *Sciences Extension Publication FA120*: 1–9.

596

597 Cross ML (2002) Microbes versus microbes: Immune signals generated by probiotic
598 lactobacilli and their role in protection against microbial pathogens. *FEMS Immunology*
599 *and Medical Microbiology* **34**(4): 245–253. doi: 10.1016/S0928-8244(02)00377-2.

600

601 Cupp AR, Schreier TM, Schleis SM (2017) Live Transport of Yellow Perch and Nile Tilapia
602 in AQUI-S 20E (10% Eugenol) at High Loading Densities. *North American Journal of*
603 *Aquaculture* **79**(2): 176–182. doi: 10.1080/15222055.2017.1281853.

604

605 del Rio-Rodriguez RE, Turnbull J (2002) Aerobic microflora of imported ornamental fish
606 from Singapore and South America. Part 2: Antimicrobial Resistant Profiles of motile
607 Aeromonads. *Fish veterinary Journal* **6**(1): 1–21.

608

609 Dixon BA, Yamashita J, Evelyn F (1990) Antibiotic resistance of *aeromonas* spp. isolated
610 from tropical fish imported from Singapore. *Journal of Aquatic Animal Health* **2**(4):
611 295–297. doi: 10.1577/1548-8667(1990)002<0295:AROASI>2.3.CO;2.

612

613 Dotta G, de Andrade JIA, Tavares Gonçalves EL, Brum A, Mattos JJ, Maraschin M, Martins
614 ML (2014) Leukocyte phagocytosis and lysozyme activity in Nile tilapia fed
615 supplemented diet with natural extracts of propolis and *Aloe barbadensis*. *Fish and*
616 *Shellfish Immunology* **39**(2): 280–284. doi: 10.1016/j.fsi.2014.05.020.

617

618 Earley RL, Edwards JT, Aseem O, Felton K, Blumer LS, Karom M, Grober MS (2006)
619 Social interactions tune aggression and stress responsiveness in a territorial cichlid fish
620 (*Archocentrus nigrofasciatus*). *Physiology and Behavior* **88**(4–5): 353–363. doi:
621 10.1016/j.physbeh.2006.04.002.

622

623 Edmonds EJ (2016). Non-invasive measures of stress in ornamental fish in response to
624 husbandry practices and associated chemicals. Ph.D. Thesis, University of the West of
625 Scotland.

626

627 Espinosa-Curiel I, Perez-Espinosa H, Gonzalez-Gonzalez J, Rodriguez-Jacobo J (2016) A
628 mobile platform for remote monitoring of water quality on live fish transport
629 containers: Lessons learned. *International Conference on Electronics, Communications*
630 *and Computers (CONIELECOMP)*, IEEE, Cholula, Mexico, 40–47. doi:
631 10.1109/CONIELECOMP.2016.7438550.

632

633 FAO (2011) Fisheries and Aquaculture topics. Ornamental fish. Topics Fact Sheets. Text by
 634 Devin Bartley. In: FAO Fisheries and Aquaculture Department. *Food & Agriculture*
 635 *Organization of the United States Rome*. FAO Updated 27 May 2005. [December 2011]
 636
 637 FAO (2016) The State of World Fisheries and Aquaculture 2016. Contributing to food
 638 security and nutrition for all. *Food & Agriculture Organization of the United States*
 639 *Rome*.
 640
 641 Ferreira PDMF, Rocha JS, Gomes JR, Caldas DW, Martins, MTS, Oliveira JM, Salaro AL,
 642 Zuanon, JAS (2017) *Curcuma longa* supplementation in the diet of *Astyanax aff.*
 643 *bimaculatus* in preparation for transport. *Aquaculture Research* 1–9. doi:
 644 10.1111/are.13277.
 645
 646 Froese R (1988) Relationship between body weight and loading densities in fish transport
 647 using the plastic bag method. *Aquaculture Research* **19**: 275–281. doi: 10.1111/j.1365-
 648 2109.1988.tb00430.x
 649
 650 Fromm PO (1980) A review of some physiological and toxicological responses of freshwater
 651 fish to acid stress. *Environmental Biology of Fishes* **5**(1): 79-93.

652

653 Gabriel NN, Qiang J, He J, Ma XY, Kpundeh MD, Xu P (2015a) Dietary *Aloe vera*
654 supplementation on growth performance, some haematobiochemical parameters and
655 disease resistance against *Streptococcus iniae* in tilapia (GIFT). *Fish and Shellfish*
656 *Immunology* **44**(2): 504–514. doi: 10.1016/j.fsi.2015.03.002.

657

658 Gabriel NN, Qiang J, Ma XY, He J, Xu P, Liu K (2015b) Dietary *Aloe vera* improves plasma
659 lipid profile, antioxidant, and hepatoprotective enzyme activities in GIFT-tilapia
660 (*Oreochromis niloticus*) after *Streptococcus iniae* challenge. *Fish Physiology and*
661 *Biochemistry* **41**(5): 1321–1332. doi: 10.1007/s10695-015-0088-z.

662

663 Gomes LC, Brinn RP, Marcon JL, Dantas LA, Brandão FR, Sampaio de Abreu J, Lemos
664 PEM, McComb DM, Baldisserotto B (2009) Benefits of using the probiotic Efinol[®] L
665 during transportation of cardinal tetra, *Paracheirodon axelrodi* (Schultz), in the
666 Amazon. *Aquaculture Research* **40**(2): 157–165. doi: 10.1111/j.1365-
667 2109.2008.02077.x.

668

669 Gomes LC, Brinn RP, Marcon JL, Dantas, LA, Brandão, FR, Sampaio de Abreu J, McComb
670 DM, Baldisserotto B (2008) Using Efinol[®] L during transportation of marbled
671 hatchetfish, *Carnegiella strigata* (Günther). *Aquaculture Research* **39**(12): 1292-1298.

672

673 Hales LS, Lay CC, Helfman GS (1990) Use of low-salinity water and gel-coating to
674 minimize handling mortality of spot, *Leiostomus xanthurus* (Perciformes: Sciaenidae).
675 *Aquaculture* **90**(1): 17–27. doi: 10.1016/0044-8486(90)90279-V.

676

677 Halliwell B, Gutteridge JM (1999) Free radicals in biology and medicine. *Oxford University*
678 Press, USA., 105–245. Available at:
679 [https://books.google.co.uk/books?hl=en&lr=&id=3DlKCgAAQBAJ&oi=fnd&pg=PP1](https://books.google.co.uk/books?hl=en&lr=&id=3DlKCgAAQBAJ&oi=fnd&pg=PP1&dq=Halliwell,+B.+and+Gutteridge,+J.M.,+2015.+Free+radicals+in+biology+and+medicine.+Oxford+University+Press,+USA.&ots=bnpG9VCrj_&sig=lqilAHvExnazNX3RdOOdp-0II-Q&redir_esc=y#v=onepage)
680 [&dq=Halliwell,+B.+and+Gutteridge,+J.M.,+2015.+Free+radicals+in+biology+and+me](https://books.google.co.uk/books?hl=en&lr=&id=3DlKCgAAQBAJ&oi=fnd&pg=PP1&dq=Halliwell,+B.+and+Gutteridge,+J.M.,+2015.+Free+radicals+in+biology+and+medicine.+Oxford+University+Press,+USA.&ots=bnpG9VCrj_&sig=lqilAHvExnazNX3RdOOdp-0II-Q&redir_esc=y#v=onepage)
681 [dicine.+Oxford+University+Press,+USA.&ots=bnpG9VCrj_&sig=lqilAHvExnazNX3](https://books.google.co.uk/books?hl=en&lr=&id=3DlKCgAAQBAJ&oi=fnd&pg=PP1&dq=Halliwell,+B.+and+Gutteridge,+J.M.,+2015.+Free+radicals+in+biology+and+medicine.+Oxford+University+Press,+USA.&ots=bnpG9VCrj_&sig=lqilAHvExnazNX3RdOOdp-0II-Q&redir_esc=y#v=onepage)
682 [RdOOdp-0II-Q&redir_esc=y#v=onepage](https://books.google.co.uk/books?hl=en&lr=&id=3DlKCgAAQBAJ&oi=fnd&pg=PP1&dq=Halliwell,+B.+and+Gutteridge,+J.M.,+2015.+Free+radicals+in+biology+and+medicine.+Oxford+University+Press,+USA.&ots=bnpG9VCrj_&sig=lqilAHvExnazNX3RdOOdp-0II-Q&redir_esc=y#v=onepage) (Accessed: 8 February 2017).
683
684 Harikrishnan R, Balasundaram C, Heo MS (2011) Impact of plant products on innate and
685 adaptive immune system of cultured finfish and shellfish. *Aquaculture* **317**(1): 1–15.
686 doi: 10.1016/j.aquaculture.2011.03.039.
687
688 Harmon TS (2009) Methods for reducing stressors and maintaining water quality associated
689 with live fish transport in tanks: a review of the basics. *Reviews in Aquaculture* **1**(1):
690 58–66. doi: 10.1111/j.1753-5131.2008.01003.x.
691
692 Harnish RA, Colotelo AH, Brown RS (2011) A review of polymer-based water conditioners
693 for reduction of handling-related injury. *Reviews in Fish Biology and Fisheries* **21**(1):
694 43–49. doi: 10.1007/s11160-010-9187-1.
695
696 Havenaar R, Ten Brink B, Huis JH (1992) Selection of strains for probiotic use. *Probiotics*
697 209–224. doi: 10.1007/978-94-011-2364-8_9.
698
699 Higuera-Ciapara I, Félix-Valenzuela L, Goycoolea FM (2006) Astaxanthin: A Review of its
700 Chemistry and Applications. *Critical Reviews in Food Science and Nutrition* **46**(2):
701 185–196. doi: 10.1080/10408690590957188.

702

703 Husen A, Sharma S (2014) Efficacy of Anesthetics for Reducing Stress in Fish During
704 Aquaculture Practices- a Review. *KUSET* **10**: 104–123.

705

706 Ims S (2011) The Efficacy and Stress-Reducing Capacity of MS-222, Benzoak and Aqui- S
707 for the Ornamental Cichlid Fish, *Metriaclima estherae*. *Norwegian University of*
708 *Science and Technology*.

709

710 Jagruthi C, Yogeshwari G, Anbazahan SM, Shanthi MLS, Arockiaraj J, Mariappan P,
711 Lernal Sudhakar GR, Balasundaram C, Harikrishnan R (2014) Effect of dietary
712 astaxanthin against *Aeromonas hydrophila* infection in common carp, *Cyprinus carpio*.
713 *Fish & Shellfish Immunology* **41**(2): 674–680. doi: 10.1016/j.fsi.2014.10.010.

714

715 Javahery S, Nekoubin H, Moradlu AH (2012) Effect of anaesthesia with clove oil in fish
716 (review). *Fish Physiology and Biochemistry* **38**(6): 1545–1552. doi: 10.1007/s10695-
717 012-9682-5.

718

719 Johnson DL, Metcalf MT (1982) Causes and Controls of Freshwater Drum Mortality during
720 Transportation. *Transactions of the American Fisheries Society* **111**(1): 58–62.

721

722 Johnson EA, Villa TG, Lewis MJ (1980) *Phaffia rhodozyma* as an astaxanthin source in
723 salmonid diets. *Aquaculture* **20**(2): 123–134. doi: 10.1016/0044-8486(80)90041-1.

724

725 Kim KH, Hwang YJ, Bai SC (1999) Resistance to *Vibrio alginolyticus* in juvenile rockfish
 726 (*Sebastes schlegeli*) fed diets containing different doses of aloe. *Aquaculture* **180**(1):
 727 13–21. doi: 10.1016/S0044-8486(99)00143-X.

728

729 Kouba A, Sales J, Sergejevová M, Kozák P, Masojídek J (2013) Colour intensity in angelfish
 730 (*Pterophyllum scalare*) as influenced by dietary microalgae addition. *Journal of*
 731 *Applied Ichthyology* **29**(1): 193–199. doi: 10.1111/jai.12010.

732

733 Lekang OI (2007) Transport of live fish. In: Lekang O.I. *Aquaculture Engineering*: 256-265.
 734 John Wiley & Sons, Chichester.

735

736 Lim LC, Dhert P, Chew WY, Dermaux V, Nelis H, Sorgeloos P. (2002) Enhancement of
 737 stress resistance of the guppy *Poecilia reticulata* through feeding with vitamin C
 738 supplement. *Journal of the World Aquaculture Society* **33**(1): 32–40. doi:
 739 10.1111/j.1749-7345.2002.tb00475.x.

740

741 Lim LC, Dhert P, Sorgeloos P (2003) Recent developments and improvements in ornamental
 742 fish packaging systems for air transport. *Aquaculture Research* **34**(11): 923–935. doi:
 743 10.1046/j.1365-2109.2003.00946.x.

744

745 Lim LC, Dhert P, Sorgeloos P (2007) Preparation for transport: Fish packaging technology.
 746 *Ornamental Fish International* **2**: 25–45.

747

748 Lim KC, Yusoff FM, Shariff M, Kamarudin MS (2017) Astaxanthin as feed supplement in
 749 aquatic animals. *Reviews in Aquaculture*.

750

751 Lorenz RT, Cysewski GR (2000) Commercial potential for *Haematococcus* microalgae as a
752 natural source of astaxanthin. *Trends in Biotechnology* **18**(4): 160–167. doi:
753 10.1016/S0167-7799(00)01433-5.

754

755 Luthra PM, Singh R, Chandra R (2001) Therapeutic uses of *Curcuma longa* (turmeric).
756 *Indian journal of clinical biochemistry* **16**(2): 153–60. doi: 10.1007/BF02864854.

757

758 Mahor G, Ali SA (2016) Recent update on the medicinal properties and use of *Aloe vera* in
759 the treatment of various ailments. *Biosci. Biotech. Res. Comm.* **9**(2): 273–288.

760

761 Marine Aquarium Council (2001) Core Handling, Husbandry, and Transport International
762 Performance Standard for the Marine Aquarium Trade.
763 Available at: http://pdf.usaid.gov/pdf_docs/PNACU671.pdf

764

765 Mars Fishcare (2014) Stress Coat NG.
766 [http://www.apifishcare.com/pdf/RM000184_85_Stress_Coat_Documents_MSDSVend](http://www.apifishcare.com/pdf/RM000184_85_Stress_Coat_Documents_MSDSVendors_2014_February_19_08-15-01-105_AM.pdf)
767 [ors_2014_February_19_08-15-01-105_AM.pdf](http://www.apifishcare.com/pdf/RM000184_85_Stress_Coat_Documents_MSDSVendors_2014_February_19_08-15-01-105_AM.pdf).

768

769 Martínez Cruz P, Ibáñez AL, Monroy Hermosillo OA, Ramírez Saad HC (2012) Use of
770 probiotics in aquaculture. *ISRN microbiology* **2012**:1-13. doi: 10.5402/2012/916845.

771

772 Mazik PM, Simco BA, Parker NC (1991) Influence of water hardness and salts on survival
773 and physiological characteristics of Striped Bass during and after transport.
774 *Transactions of the American Fisheries Society* **120**(July 2013): 121–126. doi:
775 10.1577/1548-8659(1991)120<0121:IOWHAS>2.3.CO;2.
776

777 Merlini LS, Vargas L, Piau R, Ribeiro RP, Merlini NB (2014) Effects of a homeopathic
778 complex on the performance and cortisol levels in Nile tilapia (*Oreochromis niloticus*).
779 *Homeopathy* **103**(2): 139–142. doi: 10.1016/j.homp.2013.08.005.
780

781 Nikinmaa M, Soivio A, Nakari T, Lindgren S (1983) Hauling stress in brown trout (*Salmo*
782 *trutta*): Physiological responses to transport in fresh water or salt water, and recovery in
783 natural brackish water. *Aquaculture* **34**(1–2): 93–99. doi: 10.1016/0044-
784 8486(83)90294-6.
785

786 Ottesen OH, Olafsen JA (1997) Ontogenetic development and composition of the mucous
787 cells and the occurrence of saccular cells in the epidermis of Atlantic halibut. *Journal*
788 *of Fish Biology* **50**(3): 620–633. doi: 10.1111/j.1095-8649.1997.tb01954.x.
789

790 Pan CH, Chien YH (2009) Effects of dietary supplementation of alga *Haematococcus*
791 *pluvialis* (Flotow), synthetic astaxanthin and β -carotene on survival, growth, and
792 pigment distribution of red devil, *Cichlasoma citrinellum* (Günther). *Aquaculture*
793 *Research* **40**(8): 871–879. doi: 10.1111/j.1365-2109.2008.02153.x.
794

795 Pan CH, Chien YH, Hunter B (2003) The resistance to ammonia stress of *Penaeus monodon*
 796 Fabricius juvenile fed diets supplemented with astaxanthin. *Journal of Experimental*
 797 *Marine Biology and Ecology* **297**(1): 107–118. doi: 10.1016/j.jembe.2003.07.002.
 798

799 Pan CH, Chien, YH, Wang YJ (2010) The antioxidant capacity response to hypoxia stress
 800 during transportation of characins (*Hyphessobrycon callistus* Boulenger) fed diets
 801 supplemented with carotenoids. *Aquaculture Research* **41**(7): 973–981. doi:
 802 10.1111/j.1365-2109.2009.02380.x.
 803

804 Pan CH, Chien YH, Wang YJ (2011) Antioxidant defence to ammonia stress of caracins
 805 (*Hyphessobrycon eques* Steindachner) fed diets supplemented with carotenoids.
 806 *Aquaculture Nutrition* **17**(3): 258–266. doi: 10.1111/j.1365-2095.2009.00747.x.
 807

808 Pattanasiri T, Taparhudee W, Suppakul P (2016) Anaesthetic efficacy of clove oil-coated
 809 LDPE bag on improving water quality and survival in the Siamese fighting fish, *Betta*
 810 *splendens*, during transportation. *Aquaculture International* **25**(1):1–13. doi:
 811 10.1007/s10499-016-0020-2.
 812

813 Peng SM, Shi ZH, Fei Y, Gao QX, Sun P, Wang JG (2013) Effect of high-dose vitamin C
 814 supplementation on growth, tissue ascorbic acid concentrations and physiological
 815 response to transportation stress in juvenile silver pomfret, *Pampus argenteus*. *Journal*
 816 *of Applied Ichthyology* **29**(6): 1337–1341. doi: 10.1111/jai.12250.
 817

818 Pickering A, Pottinger T, Christie P (1982) Recovery of the brown trout, *Salmo trutta* L.,
 819 from acute handling stress: a time course study. *Journal of Fish Biology* **20**(2): 229–
 820 244.
 821
 822 Pikulkaew S, Khumpirapang N, Chaisri W Okonogi S (2017) Effects of *Alpinia galanga* oil
 823 on anesthesia and stress reduction in *Oreochromis niloticus*. *Drug Discoveries &*
 824 *Therapeutics* **11**(4):186-192.
 825
 826 Plumb JA, Grizzle JM, Rogers WA (1988) Survival of Caught and Released Largemouth
 827 Bass after Containment in Live Wells. *North American Journal of Fisheries*
 828 *Management* **8**(3): 325–328. doi: 10.1577/1548-
 829 8675(1988)008<0325:SOCARL>2.3.CO;2.
 830
 831 Portz DE, Woodley CM, Cech JJ (2006) Stress-associated impacts of short-term holding on
 832 fishes. *Reviews in Fish Biology and Fisheries* **16**(2): 125–170. doi: 10.1007/s11160-
 833 006-9012-z.
 834
 835 Ramírez-Duarte WF, Pineda-Quiroga C, Martínez N, Eslava-Mocha PR (2011) Use of
 836 sodium chloride and zeolite during shipment of *Ancistrus triradiatus* under high
 837 temperature. *Neotropical Ichthyology* **9**(4): 909–914. doi: 10.1590/S1679-
 838 62252011005000036.
 839
 840 Ramírez-Duarte WF, Pineda-quiroga C, Rueda NM (2013) Evaluation of the use of sodium
 841 chloride, eugenol, and zeolite in confinement of *Ancistrus triradiatus*. *Orinoquia* **17**(1):
 842 84–95.

843

844 Rånby BG, Rabek JF (1978) Singlet oxygen reactions with organic compounds and
 845 polymers. *John Wiley & Sons*.

846

847 Readman GD, Owen SF, Murrell JC, Knowles (2013) Do Fish Perceive Anaesthetics as
 848 Aversive?. *PLoS ONE* **8**(9): e73773. doi: 10.1371/journal.pone.0073773.

849

850 Reynolds T, Dweck AC (1999) *Aloe vera* leaf gel: A review update. *Journal of*
 851 *Ethnopharmacology* **68**(1–3): 3–37. doi: 10.1016/S0378-8741(99)00085-9.

852

853 Rimmer MA, Franklin B (1997) Development of live fish transport techniques. QDPI.
 854 Available at: http://frdc.com.au/research/Final_Reports/1993-184-DLD.pdf

855

856 Roberts RJ, Agius C, Saliba C, Bossier P, Sung YY (2010) Heat shock proteins (chaperones)
 857 in fish and shellfish and their potential role in relation to fish health: a review. *Journal*
 858 *of Fish Diseases* **33**(10): 789–801. doi: 10.1111/j.1365-2761.2010.01183.x.

859

860 Rollo A, Sulpizio R, Nardi M, Silvi S, Orpianesi C, Caggiano M, Cresci A, Carnevali O.
 861 (2006) Live microbial feed supplement in aquaculture for improvement of stress
 862 tolerance. *Fish Physiology and Biochemistry* **32**(2): 167–177. doi: 10.1007/s10695-
 863 006-0009-2.

864

865 Rose S, Hill R, Bermudez LE, Miller-Morgan T (2013) Imported ornamental fish are
 866 colonized with antibiotic-resistant bacteria. *Journal of Fish Diseases* **36**(6): 533–542.
 867 doi: 10.1111/jfd.12044.

868

869 Ross LG, Ross B, Ross B (2008) Anaesthetic and sedative techniques for aquatic animals/
870 Oxford: *Blackwell Publishing*.
871

872 Rubec PJ, Cruz FP (2005) Monitoring the chain of custody to reduce delayed mortality of
873 net-caught fish in the aquarium trade. *SPC Live Reef Fish Information Bulletin* **13**: 13–
874 23.
875

876 Saccol EM, Jerez-Cepa I, Ourique GM, Pes TS, Gressler LT, Mourao RH, Martinez-
877 Rodriguez G, Mancera JM, Baldisserotto B, Pavanato MA, Martos-Sitcha JA (in press)
878 *Myrcia sylvatica* essential oil mitigates molecular, biochemical and physiological
879 alterations in *Rhamdia quelen* under different stress events associated to transport.
880 *Research in Veterinary Science*.
881

882 Saccol EMH, Toni C, Pês TS, Ourique GM, Gressler LT, Silva LVF, Mourão RHV, Oliveira
883 RB, Baldisserotto B, Pavanato MA (2016) Anaesthetic and antioxidant effects of
884 *Myrcia sylvatica* (G. Mey.) DC. and *Curcuma longa* L. essential oils on tambaqui
885 (*Colossoma macropomum*). *Aquaculture Research* **48**(5): 1–20. doi:
886 10.1111/are.13034.
887

888 Sadovy Y (2002) Death in the live reef fish trades. *SPC Live Reef Fish Information Bulletin*,
889 **10**: 1–32.
890

891 Salaro AL, Camplelo DAV, Tavares MM, Braga LGT, Pontes MD, Zuanon JAS (2015)
892 Transport of *Astyanax altiparanae* Garutti and Britski, 2000 in saline water. *Acta*

893 *Scientiarum - Biological Sciences* **37**(2): 137–142. doi:
894 10.4025/actascibiolsci.v37i2.26884.
895
896 Salbego J, Becker AG, Gonçalves JF, Menezes CC, Heldwein CG, Spanevello RM, Loro VL,
897 Schetinger MRC, Morsch VM, Heinzmann BM, Baldisserotto B (2014) The essential
898 oil from *Lippia alba* induces biochemical stress in the silver catfish (*Rhamdia quelen*)
899 after transportation. *Neotropical Ichthyology* **12**(4): 811–818. doi: 10.1590/1982-0224-
900 20130178.
901
902 Salbego J, Becker AG, Parodi TV, Zeppenfeld CC, Gonçalves JF, Loro VL, Morsch VMM,
903 Schetinger MRC, Maldaner G, Morel AF, Baldisserotto B (2015) Methanolic extract of
904 *Condalia buxifolia* added to transport water alters biochemical parameters of the silver
905 catfish *Rhamdia quelen*. *Aquaculture* **437**: 46–50. doi:
906 10.1016/j.aquaculture.2014.11.022.
907
908 Sales J, Janssens GPJ (2003) Nutrient requirements of ornamental fish. *Aquatic Living*
909 *Resources* **16**(6): 533–540. doi: 10.1016/j.aquiliv.2003.06.001.
910
911 Sampaio FDF, Freire CA (2016) An overview of stress physiology of fish transport: Changes
912 in water quality as a function of transport duration. *Fish and Fisheries* **17**(4)1–18. doi:
913 10.1111/faf.12158.
914
915 Sandilands J, Drynan K, Roberts RJ (2010) Preliminary studies on the enhancement of
916 storage time of chilled milt of Atlantic salmon, *Salmo salar* L., using an extender

917 containing the TEX-OE[®] heat shock-stimulating factor. *Aquaculture Research* **41**(4):
918 568–571. doi: 10.1111/j.1365-2109.2009.02354.x.

919

920 Sepulchro LCOR, Carvalho MAG, Gomes LC (2016) Salinity does not alter the effectiveness
921 of menthol as an anesthetic and sedative during the handling and transport of juvenile
922 fat snook (*Centropomus parallelus*). *Brazilian Journal of Biology*. Instituto
923 Internacional de Ecologia **76**(3): 757–763. doi: 10.1590/1519-6984.04115.

924

925 Shephard KL (1994) Functions for fish mucus. *Reviews in Fish Biology and Fisheries* **4**(4):
926 401–429. doi: 10.1007/BF00042888.

927

928 Snellgrove DL, Alexander LG, Stevenson AE (2007) The effect of Stress Coat[®] on the
929 excretory loss of cortisol from the common goldfish *Carassius auratus* following a
930 netting stressor. Aquaculture Conference abstract, [Feb. 26 – Mar. 2, San Antonio,](#)
931 [Texas, USA.](#)

932

933 Sung YY, Roberts RJ, Bossier P (2012) Enhancement of Hsp70 synthesis protects common
934 carp, *Cyprinus carpio* L., against lethal ammonia toxicity. *Journal of Fish Diseases*
935 **35**(8): 563–568. doi: 10.1111/j.1365-2761.2012.01397.x.

936

937 Sutili FJ, Gatlin DM, Heinzmann BM, Baldisserotto B (2017) Plant essential oils as fish diet
938 additives: benefits on fish health and stability in feed. *Reviews in Aquaculture*.

939

940 Swann L (1992) Transport of Fish in Bags. Extention publications (MU), Illinois-Indiana Sea
941 Grant Program.

942

943 Swanson C, Mager RC, Doroshov SI, Cech JJ (1996) Use of Salts, Anesthetics, and Polymers
944 to Minimize Handling and Transport Mortality in Delta Smelt. *Transactions of the*
945 *American Fisheries Society* **125**(2): 633–644. doi: 10.1577/1548-8659(1996)125.

946

947 Tacchi L, Lowrey L, Musharrafieh R, Crossey K, Larragoite ET, Salinas I (2015) Effects of
948 transportation stress and addition of salt to transport water on the skin mucosal
949 homeostasis of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* **435**: 120–127. doi:
950 10.1016/j.aquaculture.2014.09.027.

951

952 Taiwo V, Olukunle OA, Ozor IC, Oyejobi AT (2005) Consumption of aqueous extract of raw
953 *Aloe Vera* leaves: histopathological and biochemical studies in rat and tilapia. *African*
954 *Journal of Biomedical Research* **8**(3): 169–178.

955

956 Tondolo JSM, De Amaral LP, Simões LN, Garlet QI, Schindler B, Oliveira TM, Da Silva
957 BF, De Gomes LC, Baldisserotto B, Mallmann CA, Heinzmann BM (2013) Anesthesia
958 and transport of fat snook *Centropomus parallelus* with the essential oil of *Nectandra*
959 *megapotamica* (Spreng.) Mez. *Neotropical Ichthyology* **11**(3): 667–674. doi:
960 10.1590/S1679-62252013000300020.

961

962 Trust TJ, Whitby JL (1976) Antibiotic resistance of bacteria in water containing ornamental
963 fishes. *ANTIMICROBIAL AGENTS AND CHEMOTHERAPY* **10**(4): 598–603.

964

965 Urbinati EC, Carneiro PCF (2006) Sodium chloride added to transport water and
 966 physiological responses of Matrinxã *Brycon amazonicus* (Teleost: Characidae). *Acta*
 967 *Amazonica* **36**(4): 569–572. doi: 10.1590/S0044-59672006000400020.
 968
 969 Vallejos-Vidal E, Reyes-López F, Teles M, MacKenzie S (2016) The response of fish to
 970 immunostimulant diets. *Fish and Shellfish Immunology* **56**: 34–69. doi:
 971 10.1016/j.fsi.2016.06.028.
 972
 973 Van Den Sande AP (1974) The long-distance transport of fishes and aquatic invertebrates.
 974 *International Zoo Yearbook* **14**: 43–47. doi: 10.1111/j.1748-1090.1974.tb00759.x
 975
 976 Vázquez B, Avila G, Segura D, Escalante B (1996) Antiinflammatory activity of extracts
 977 from *Aloe vera* gel. *Journal of ethnopharmacology* **55**(1): 69–75. doi: 10.1016/S0378-
 978 8741(96)01476-6.
 979
 980 Verner-jeffreys DW, Welch TJ, Schwarz T, Pond MJ, Woodward MJ, Haig SJ, Rimmer
 981 GSE, Roberts E, Morrison V, Baker-Austin C (2009) High prevalence of multidrug-
 982 tolerant bacteria and associated antimicrobial resistance genes isolated from ornamental
 983 fish and their carriage water. *PLoS ONE* **4**(12): 8388. doi:
 984 10.1371/journal.pone.0008388.
 985
 986 Vogler BK, Ernst E (1999) *Aloe vera*: a systematic review of its clinical effectiveness. *The*
 987 *British journal of general practice: the journal of the Royal College of General*
 988 *Practitioners* **49**(447): 823–828.
 989

- Volpatti D, D'Angelo L, Jeney G, Jeney Z, Anderson DP, Galeotti, M (1998) Nonspecific immune response in fish fed glucan diets prior to induced transportation stress. *Journal of Applied Ichthyology* **14**(3–4): 201–206. doi: 10.1111/j.1439-0426.1998.tb00642.x.
- Wang YJ, Chien YH, Pan CH (2006) Effects of dietary supplementation of carotenoids on survival, growth, pigmentation, and antioxidant capacity of characins, *Hyphessobrycon callistus*. *Aquaculture* **261**(2): 641–648. doi: 10.1016/j.aquaculture.2006.08.040.
- Wedemeyer G (1996) *Physiology of Fish in Intensive Culture Systems*. New York: Chapman & Hall, *International Thompson Publishing*. Available at: [https://books.google.co.uk/books?hl=en&lr=&id=KU_s_aHoW86oC&oi=fnd&pg=PR11&dq=Wedemeyer+GA+\(1996\)+Physiology+of+fish+in+intensive+culture+systems&ots=iDglR5skaO&sig=ad1mTDGpVbSGy2vWkO_1-hV2Svk#v=onepage&q=](https://books.google.co.uk/books?hl=en&lr=&id=KU_s_aHoW86oC&oi=fnd&pg=PR11&dq=Wedemeyer+GA+(1996)+Physiology+of+fish+in+intensive+culture+systems&ots=iDglR5skaO&sig=ad1mTDGpVbSGy2vWkO_1-hV2Svk#v=onepage&q=) (Accessed: 24 February 2017).
- Weirich CR, Tomasso JR, Smith TIJ (1992) Confinement and transport-induced stress in white bass *Morone chrysops* x striped bass *M. saxatilis* hybrids: Effect of calcium and salinity. *Journal of the World Aquaculture Society* **23**(1): 49–57. doi: 10.1111/j.1749-7345.1992.tb00750.x.
- Wiese J, McPherson S, Odden M (2004) Effect of *Opuntia ficus indica* on symptoms of the alcohol hangover. *Archives of Internal Medicine* **164**(12): 1334–1340. Available at: <http://jamanetwork.com/journals/jamainternalmedicine/fullarticle/217133> (Accessed: 21 March 2017).

1015 Wong SC, Dykstra M, Campbell JM, Earley RL, Dykstra M, Wong SC, Earley RL (2015)
 1016 Measuring water-borne cortisol in convict cichlids (*Amatitlania nigrofasciata*): is the
 1017 procedure a stressor? *Behaviour* **145**(10): 1283–1305.
 1018

1019 Yanong RPE (2003) Use of Antibiotics in Ornamental Fish Aquaculture 1. *University of*
 1020 *Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences,*
 1021 *EDIS.*
 1022

1023 Yu BP (1994) Cellular defenses against damage from reactive oxygen species. *Physiological*
 1024 *reviews* **74**(1): 139–162.
 1025

1026 Zanuzzo FS, Biller-Takahashi, JD, Urbinati, EC (2012) Effect of *Aloe vera* extract on the
 1027 improvement of the respiratory activity of leukocytes of matrinxã during the transport
 1028 stress. *Revista Brasileira de Zootecnia* **41**(10): 2299–2302. doi: 10.1590/S1516-
 1029 35982012001000023.
 1030

1031 Zanuzzo FS, Sabioni RE, Montoya NF, Favero G, Urbinati EC (2017) *Aloe vera* enhances the
 1032 innate immune response of pacu (*Piaractus mesopotamicus*) after transport stress and
 1033 combined heat killed *Aeromonas hydrophila* infection. *Fish and Shellfish Immunology*
 1034 **65**: 198–205. doi: 10.1016/j.fsi.2017.04.013.
 1035

1036 Zanuzzo FS, Urbinati EC, Nash GW, Gamperl AK (2015a) Steelhead trout *Oncorhynchus*
 1037 *mykiss* metabolic rate is affected by dietary *Aloe vera* inclusion but not by mounting an
 1038 immune response against formalin-killed *Aeromonas salmonicida*. *Journal of Fish*
 1039 *Biology* **87**(1): 43–53. doi: 10.1111/jfb.12690.

1040

1041 Zanutto FS, Urbinati, EC, Rise ML, Hall JR, Nash GW, Gamperl AK (2015b) *Aeromonas*

1042 *salmonicida* induced immune gene expression in *Aloe vera* fed steelhead trout,

1043 *Oncorhynchus mykiss* (Walbaum). *Aquaculture* **435**: 1–9. doi:

1044 10.1016/j.aquaculture.2014.09.010.

1045

1046 Zeppenfeld CC, Toni C, Becker AG, Miron DS, Parodi TV, Heinzmann BM, Barcellos LJG,

1047 Koakoski G, Rosa JGS, Loro VL, Cunha MA, Baldissierotto B (2014) Physiological and

1048 biochemical responses of silver catfish, *Rhamdia quelen*, after transport in water with

1049 essential oil of *Aloysia triphylla* (L’Herit) Britton. *Aquaculture* **418**: 101–107. doi:

1050 10.1016/j.aquaculture.2013.10.013.

1051

1052 Zink IC, Benetti DD, Douillet PA, Margulies D, Scholey VP (2011) Improvement of water

1053 chemistry with *Bacillus* probiotics inclusion during simulated transport of yellowfin

1054 tuna yolk sac larvae. *North American Journal of Aquaculture* **73**(1): 42-48

1055

1056 Table 1. Research into the use of essential oils during the transport of fishes.

Species	Life stage/Size	Essential oil	Concentration	Stressor	Effect	Reference
Silver catfish (<i>Rhamdia quelen</i>)	Juvenile (mean: 64.5 ± 6.1 g and 18.85 ± 0.57 cm)	<i>Lippia alba</i>	0, 10 µl l ⁻¹	5, 6 or 7 h transport	Reduction of lipoperoxidation (LPO), catalase, superoxide dismutase and glutathione-S-transferase in the liver in fish transported for 5 h. Reduction of LPO in the gill of fish transported for 5 h and 7 h.	Azambuja <i>et al.</i> , 2011
Silver catfish (<i>Rhamdia quelen</i>)	mean: 262.0 ± 73.5 g, 38.5 ± 1.1 cm	<i>Aloysia triphylla</i>	0, 30, 40 µl l ⁻¹	6 h transport	Lower plasma cortisol and ion loss compared to control, and higher plasma Na ⁺ and Cl ⁻ concentrations. Lower hepatic glycogen and glucose	Zeppenfeld <i>et al.</i> , 2014

					concentrations in the liver.	
					Lower muscle lactate and higher muscle glucose levels.	
Fat snook (<i>Centropomus parallelus</i>)	Juvenile (mean $\pm$ SEM: 1.6 $\pm$ 0.4 g and 4.8 $\pm$ 0.4 cm)	Menthol (5-Methyl-2-(propan-2-yl)cyclohexan-1-ol)	0, 3.7 or 7.4 mg l ⁻¹	10 h transport	Effective anaesthetic for short term handling. No effects on mortality, ammonia, dissolved oxygen, nitrite levels after transport.	Sepulchro <i>et al.</i> , 2016
Fat snook (<i>Centropomus parallelus</i>)	Juvenile	<i>Nectandra megapotamica</i>	0, 15, 30 μ l l ⁻¹	10 h transport	Higher post-transport mortality in the 30 μ l l ⁻¹ than the 15 μ l l ⁻¹ and the control group.	Tondolo <i>et al.</i> , 2013
Silver catfish	mean $\pm$	<i>Condalia buxifolia</i>	0, 5, 10 μ l l ⁻¹	6 h transport +	Lower water TAN in both groups	Salbego <i>et al.</i> 2015

(Rhamdia quelen)	SEM:		pre-sedation	transported in <i>C. buxifolia</i> ..
	420.1 ±			
	8.8 g and			Lower net efflux of Na ⁺ , Cl ⁻ and K ⁺ in
	21.2 ± 2.3			both groups transported in <i>C. buxifolia</i> .
	cm			Higher PvO ₂ , PvCO ₂ and HCO ₃ ⁻ in 5 µl
				l ⁻¹ group.
				Lower hepatic lactate in the 10 µl l ⁻¹
				group.
				Lower muscle lactate in both 5 and 10 µl
				l ⁻¹ vs. control group.
				Improved antioxidant status.

Silver catfish	mean ±	<i>L. alba</i>	0, 30, 40 µl l ⁻¹	6 h transport +	Lower water TAN.	Becker <i>et al.</i> 2015
----------------	--------	----------------	------------------------------	-----------------	------------------	---------------------------

(<i>Rhamdia quelen</i>)	SEM:			pre-sedation		
	420.1 ±				Lower net efflux of Na ⁺ , Cl ⁻ and K ⁺ .	
	8.8 g and					
	21.2 ± 2.3				Higher PvO ₂ , PvCO ₂ and HCO ₃ ⁻ in the	
	cm				40 µl l ⁻¹ group.	
					Higher plasma cortisol in the 30 µl l ⁻¹	
					compared to the control group.	
Silver catfish	mean ±	<i>L. alba</i>	0, 30, 40 µl l ⁻¹	6 h transport +	Lower net efflux of Na ⁺ , Cl ⁻ and K ⁺ .	Salbego <i>et al.</i> 2014
(<i>Rhamdia quelen</i>)	SEM:			pre-sedation		
	420.1 ±				An exposure of 30-40 µl l ⁻¹ induced	
	8.8 g and				oxidative stress and elevated cortisol.	
	21.2 ± 2.3					
	cm					
Silver catfish	mean ±	<i>C. buxifolia</i>	0, 25, 50 µl l ⁻¹	12 h transport	Lower non-ionized ammonia levels.	Becker <i>et al.</i> 2013
(<i>Rhamdia quelen</i>)	SEM:					

	1.50 ±				Lower net efflux of Na ⁺ , Cl ⁻ and K ⁺ .	
	0.02 g					
	and 165.7					
	± 22.5 g					
Swordtail fish (<i>Xiphophorus hellerii</i>)	mean: 2.49 ± 0.62 g	Valerian root (<i>Valeriana officinalis</i>)	1 g l ⁻¹	24 h simulated transport	Lower mortality and whole body cortisol.	Abasali & Mohamad (2010)
Nile tilapia (<i>Oreochromis niloticus</i>)	Juvenile (mean: 1.34 ± 0.07 g and 4.25 ± 0.22 cm)	<i>Alpinia galanga</i>	150 mg l ⁻¹	4 h simulated transport at three loading densities (100, 200, and 300 fish/plastic bag)	No mortality. Slowed movement, higher dissolved oxygen and lower TAN in all loading densities. Lower NH ₃ in the 100 and 200 fish/plastic bag densities.	Pikulkaew <i>et al.</i> (2017)
Silver catfish (<i>Rhamdia quelen</i>)	Juveniles (mean:	<i>Myrcia sylvatica</i>	25, 35 µl l ⁻¹	6 h transport	Lower plasma cortisol and lactate levels, increased Na ⁺ /K ⁺ -ATPase gill activity.	Saccol <i>et al.</i> (in press)

	8.9 ± 2.7	Lower gene expression of corticotropin-
	g and	releasing hormone,
	12.4 ± 1.3	proopiomelanocortins, prolactin and
	cm)	somatolactin indicating lower stress
		pathways activation
1057		
1058		
1059		

1060 Table 2. Studies investigating the effects of polymer-based water conditioners (adapted from Harnish *et al.* 2011).

Species	Life stage/Size	Water conditioner	Concentration	Stressor	Effect	Reference
Smallmouth bass (<i>Micropterus dolomieu</i>)	Adult	Catch'n'Rel-ease Formula [®]	5 g l ⁻¹	Live release angling tournaments	Cardiac disturbances recovered within ~60min for control fish. Cardiac disturbances in fish exposed to Catch'n'Release lasted for ~180min	Cooke <i>et al.</i> (2002)
Delta smelt (<i>Hypomesus transpacificus</i>)	mean: 4.7 (August) 5.1 cm (November)	NovAqua [®] in 8 g l ⁻¹ NaCl	0.5 ml l ⁻¹	Holding and transport post-capture	NovAqua in 8‰ NaCl increased 72 h survival (54.8%) when compared to the control with 8‰ NaCl (27.9%).	Swanson <i>et al.</i> (1996)
Largemouth bass (<i>Micropterus salmoides</i>)	Unreported	Unspecified commercial product [§]	1 mg 75 l ⁻¹ water	Live release angling tournaments	Survival for fish held in water with conditioner for 3-9 h was higher (96.5%) than fish held in unconditioned water (90.8%).	Plumb <i>et al.</i> (1988)

1061 ^sThe water conditioner contained unspecified quantities of sodium chloride, potassium chloride, sodium thiosulfate, pyrogenic silica, dimethylketone, alpha-
1062 methylquinoline, methylene blue, nitromersol, ethylenediaminetetraacetate, triethyleneglycol, and acriflavine (Plumb *et al.* 1988).
1063

1064 Table 3. Research into the effects of adding NaCl to the water of fishes during transport.

Species	Life stage/Size	Concentration	Stocking density	Stressor	Effect	Reference
Brown trout (<i>Salmo trutta</i>)	mean: 76.2 ± 1.7 g, 20.4 ± 0.1 cm.	0.6 g l ⁻¹ NaCl	100 g l ⁻¹	14 h transport	Smaller increase of blood oxygen carrying capacity than fish in the control group. No reduction of plasma osmolality and higher plasma levels of Cl ⁻ and Na ⁺ . Higher liver glycogen and muscle lipid contents when compared to control fish.	Nikinmaa <i>et al.</i> 1983

Freshwater drum (<i>Aplodinotus grunniens</i>)	mean: 36.5 cm	5 g l ⁻¹ NaCl	60 g l ⁻¹ 120 g l ⁻¹	6 h transport	Reduction in immediate and delayed mortality after transport.	Johnson & Metcalf 1982
Striped bass (<i>Morone saxatilis</i>)	mean ± SE: 72 ± 2.5 g	0 and 1 g l ⁻¹ NaCl	180 g l ⁻¹	5 h transport	Reduction in delayed mortality over a 4-week period following transport.	Mazik <i>et al.</i> 1991
Lower plasma cortisol, glucose and potassium concentrations, and higher plasma sodium and chloride concentrations during recovery.						
White bass (<i>M. chrysops</i>) x striped bass hybrids	mean ± SE: 38.7 ± 1.1 g and 58.7 ± 1.1 g	Fresh water (5, 10, 20, 40, 80 mg l ⁻¹ Ca ²⁺) Salt water (1,	36.4 g l ⁻¹ 55.2 g l ⁻¹	6.5 h confinement	Highest survival for fish in fresh water was at 80 mg l ⁻¹ Ca ²⁺ at both stocking densities (with higher survival at higher	Weirich <i>et al.</i> 1992

8, 16, 24 g l⁻¹

NaCl)

stocking).

Highest survival in salt water

was at 8 g l⁻¹ NaCl, with similar

survival at both densities.

Plasma osmolality decreased

during confinement in fresh

water.

In sea water, fish in 8 g l⁻¹

maintained plasma osmolality.

Plasma osmolality at 16 and 24

g l⁻¹ increased.

mean group	Fresh water (5, 60 g l ⁻¹	12 h	Fish in all treatments had <5%
weights = 3.7	10, 20, 40, 80	simulated	mortality with no variation in
± 0.3 to 55.8	mg l ⁻¹ Ca ²⁺)	transport	mortality levels between the
± 1.7 g	Salt water (1, 8, 16, 24 g l ⁻¹		treatments.
	NaCl)		

Xenocara	mean: 10.4 ±	0.5, 1 g l ⁻¹	61.75 g l ⁻¹	48 h transport	Lower blood glucose levels	Ramírez-
(<i>Ancistrus triradiatus</i>)	4.6 g	NaCl	¹		after transport in both 0.5 and 1	Duarte <i>et</i>
					g l ⁻¹ NaCl groups compared to	<i>al.</i> 2011
					levels in fish transported in	
					fresh water or with zeolites.	
					No significant difference	
					between 0.5 and 1 g l ⁻¹ NaCl	

groups.

Lower mortality immediately
post transport and 7 days post
transport in both 0.5 and 1 g l⁻¹
NaCl groups compared to levels
in fresh water or with zeolites.

No significant difference
between 0.5 and 1 g l⁻¹ NaCl.

Xenocara	mean: 9.0 ±	1 or 2 g l ⁻¹	137.5 g l ⁻¹	12 h	Lower blood glucose	Ramírez-
(<i>Ancistrus triradiatus</i>)	6.4 g and 7.0	NaCl	¹	simulated	concentrations.	Duarte <i>et</i>
	± 1.4 cm			transport		<i>al.</i> 2013
					Reduction in mortality	
					following transport.	

1 or 2 g l⁻¹ 82.3 g l⁻¹ 48 h Lower blood glucose
NaCl simulated concentrations but higher
transport mortality during the 7 day post
transport recovery period in the
2 g l⁻¹ NaCl treatment.

<i>Astyanax altiparanae</i>	Fingerlings	0, 3, 6, 9, 12, (mean: 0.37 ± 0.05 g)	0.37 g ± 0.05 l ⁻¹	96 h salinity exposure with no food.	All fish survived for up to 6 h in 0, 3, 6, and 9 g L ⁻¹ NaCl. After 96h, mortality was 75% on the 9 g L ⁻¹ NaCl and 100% in the 12, and 15 g L ⁻¹ NaCl.	Salaro <i>et</i> <i>al.</i> 2015
		0, 3, 6, and 9 g l ⁻¹ NaCl	22, 30 and 37 g l ⁻¹	8 h simulated transport	Blood glucose levels were significantly lower in the 30 and 37 g l ⁻¹ fish transported in the 3, 6, 9 g l ⁻¹ NaCl compared to the 0 g l ⁻¹ NaCl.	

Rainbow trout (<i>Oncorhynchus mykiss</i>)	mean: 200 g	0.5 g l ⁻¹ NaCl	Unknown	5 h transport	No increase in plasma glucose levels, compared to fish transported in fresh water where transport elevated plasma glucose.	Tacchi <i>et al.</i> 2015
					Fish transported in 0.5 NaCl g l ⁻¹ had a thinner mucus layer than fish transported in fresh water (see Water Conditioners)	

1065

1066

1067

1068

1069