UWS Academic Portal

An exploratory study of the relationship between psychosocial hazard and ambulatory physiological response in higher education employees

Campbell, Thomas George; Westbury, Tony; Davison, Richard; Florida-James, Geraint

Published in:
International Journal of Workplace Health Management

DOI:
[10.1108/IJWHM-11-2015-0068](https://doi.org/10.1108/IJWHM-11-2015-0068)

Published: 12/09/2016

Document Version
Peer reviewed version

[Link to publication on the UWS Academic Portal](#)

Citation for published version (APA):
Campbell, T. G., Westbury, T., Davison, R., & Florida-James, G. (2016). An exploratory study of the relationship between psychosocial hazard and ambulatory physiological response in higher education employees. *International Journal of Workplace Health Management*, 9(3), 360-374. <https://doi.org/10.1108/IJWHM-11-2015-0068>

General rights

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact pure@uws.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Emerald

International Journal
of Workplace
Health Management

**An Exploratory Study of the Relationship between
Psychosocial Hazard and Ambulatory Physiological
Response in Higher Education Employees.**

Journal:	<i>International Journal of Workplace Health Management</i>
Manuscript ID	IJWHM-11-2015-0068.R2
Manuscript Type:	Research Paper
Keywords:	Workplace Health, Workplace Wellness, Stress, Psychosocial hazard, work-related demand, Management Standards Indicator Tool

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

An Exploratory Study of the Relationship between Psychosocial Hazard and Ambulatory
Physiological Response in Higher Education Employees.

Abstract

Purpose: As exposure to psychosocial hazard at work represents a substantial risk factor for employee health in many modern occupations, being able to accurately assess how employees cope with their working environment is crucial. As the workplace is generally accepted as being a dynamic environment consideration should be given to the interaction between employees and the acute environmental characteristics of their workplace. The aim of this study was to investigate the effects of both acute demand and chronic work-related psychosocial hazard upon employees through ambulatory assessment of heart rate variability and blood pressure.

Design: A within-subjects repeated measures design was used to investigate the relationship between exposure to work-related psychosocial hazard and ambulatory heart rate variability and blood pressure in a cohort of higher education employees. Additionally the effect of acute variation in perceived work-related demand was investigated.

Results: Two dimensions of the Management Standards were found to demonstrate an association with heart rate variability; more hazardous levels of “demand” and “relationships” were associated with decreased SDNN. Significant changes in blood pressure and indices of heart rate variability were observed with increased acute demand.

Originality: This is the first attempt to combine the Health and Safety Management Standards Indicator Tool with physiological assessment of employees. The results provide evidence of associations between scores on the indicator tool and ambulatory heart rate variability as well as demonstrating that variation in acute perceived work-related demand is associated with alterations to autonomic and cardiovascular function. This has implications not only for

employee health and workplace design but also for future studies employing ambulatory physiological monitoring.

Introduction

The ‘ivory towers’ of academia have traditionally afforded relative sanctuary from exposure to occupational stress, primarily through high levels of autonomy and intellectual freedom. The role of the academic was once clearly delineated, with teaching and research constituting the majority of workload, whilst administration accounted for relatively little work time (Houston *et al.*, 2006). However, in the UK, universities have been forced to prioritise fiscal performance following reductions in public funding in the wake of the Education Reform Act (1988). UK Government policy now dictates that universities must contribute to the economy (Lam, 2010) with research funding being largely dependent upon this contribution (Etkowitz *et al.*, 2000). As a result, despite no reduction in teaching or research responsibilities, academics must devote significantly more time to administrative work (Kinman and Jones, 2003; Tight, 2010) and are increasingly being tasked with securing research funding through entrepreneurial activities. Academia is therefore no longer immune from the sources of occupational stress associated with globalisation and market forces.

Recent reports of academics suffering from stress as a result of overload are ample, with work overload, task overload, and role overload, as well as the difficulty of balancing multiple roles and lack of role clarity, being commonly cited factors (Winter *et al.*, 2000; Gillespie *et al.*, 2001; Kinman and Jones, 2003; Barret and Barret, 2007; Devenport, 2008). Stress has been identified as a key predictor of academics’ intent to move institution (Ryan *et al.*, 2012), and is also associated with intention of leaving the profession entirely (Kinman and Jones, 2003). The deleterious effects of exposure to stress upon health, particularly the incidence of hypertension and cardiac disease, are widely known, and a study of UK academics found that one quarter had suffered from a stress related illness in the previous

year (Kinman and Jones, 2003). Higher education employees have also been shown to be at greater risk of psychological illness than the general population (Winefield *et al.*, 2003); and UK lecturers report poorer than average levels of psychological wellbeing (Johnson *et al.*, 2005).

Although the exact mechanisms are still to be determined, the autonomic nervous system is a likely pathway linking exposure to psychosocial strain and disease (Thayer and Lane, 2007). Heart rate variability, an independent predictor of cardiovascular mortality and cardiac events (Tsuji *et al.*, 1996; Kikuya *et al.*, 2000), provides a non-invasive insight into the functioning of the autonomic nervous system. Time domain measures of heart rate variability such as the standard deviation of the normal-to-normal interval (SDNN), which represents the total variability occurring across all of the components that contribute to HRV, are calculated from the intervals between successive normal heart beats, while spectral analysis allows for quantification of the variability of the signal occurring within distinct frequency components (Task Force of the European Society of Cardiology and The North American Society of Pacing and Electrophysiology., 1996). Low frequency (LF: 0.025 to 0.15 Hz), high frequency (HF; 0.15 to 0.4 Hz), and the ratio between the two (LFHF ratio) are the most widely reported components, with high frequency heart rate variability reflecting vagal parasympathetic activity. Although often claimed to provide a measure of sympathetic activity, low frequency heart rate variability actually reflects baroreflex function (Goldstein, 2011; Rahman *et al.*, 2011; Reyes del Paso *et al.*, 2013). Reduced vagal tone has been found to represent a risk factor for cardiovascular disease (Liao 1997; Curtis and O'Keefe, 2002) and work-related psychosocial strain has repeatedly been associated with heart rate variability (Van Amelsvoort *et al.*, 2000; Vrijkotte *et al.*, 2000; Hjortskov *et al.*, 2004; Lucini *et al.*, 2007 Chandola *et al.*, 2008 Loerbroks *et al.*, 2010) and increased ambulatory blood pressure (Van Egeren, 1992; Fauvel *et al.*, 2001; Brown *et al.*, 2006; Guimont *et al.*, 2006).

In the UK, The Health and Safety Executive, who act as the national independent watchdog for work-related health, safety and illness, currently advocate the use of a risk assessment approach to identify environments believed to invoke work stress, through the application of their management standards and associated indicator tool (Health and Safety Executive, 2005). The Indicator Tool is a 35-item self-report questionnaire which measures exposure to various dimensions of work design that, if not properly managed, are associated with poor health and well-being, lower productivity and increased sickness absence. The Indicator Tool not only differs from conventional models of occupational stress, such as the job-demand-control model (Karasek, 1979) and its subsequent adaptations (Johnson & Hall, 1988; Demerouti *et al.*, 2001) or the effort-reward imbalance model (Siegrist *et al.*, 1986), by assessing a greater number of dimensions, but also in the belief that each of the seven dimensions represent a potential risk to employees health and wellbeing in isolation. Despite being firmly grounded in occupational stress theory, the overarching premise of this approach is appealing in its simplicity, in that minimising exposure to factors known to represent a hazard for the experience of stress reduces the incidence of stress-related problems.

Although there is no shortage of evidence to demonstrate that exposure to work-related psychosocial strain is associated with both blood pressure and heart rate variability, to date no attempt has been made to specifically investigate the effects of exposure to psychosocial hazard using the Management Standards Indicator Tool. As a recent nationwide survey of UK higher education employees reported lower than average scores on all but one of the management standards (Kinman and Court, 2010) this cohort of employees are potentially at significant risk of experiencing unfavourable health outcomes. The main aim of the present study was therefore to investigate whether exposure to psychosocial hazard at work is associated with autonomic function in higher education employees. Additionally, although rarely attended to during ambulatory workplace assessment, there is evidence to

suggest that heart rate variability is affected by acute work-related factors (Hjortskov *et al.*, 2004; Filaire, 2010). Therefore, a further aim of the study was to investigate whether work-time ambulatory assessments of heart rate variability and blood pressure are influenced by acute demand. The three hypotheses of the study were: i) academics would be exposed to greater psychosocial hazard and would demonstrate less favourable work-time cardiovascular and autonomic responses than general staff; ii) scores on the management standards indicator tool would be positively associated with physiological stress responses i.e. greater perceived exposure to psychosocial hazard would result in greater blood pressure and reduced heart rate variability; and iii) blood pressure and measures of heart rate variability will differ according to the acute work-related demand of the measurement day.

Method

Participants

Participants were recruited via an advertisement placed on Edinburgh Napier University's internal staff intranet and email. Inclusion criteria required participants to be employed by the university on a permanent full-time contracted basis and to have been working in that role for a minimum of 6 months. Participants were excluded if they smoked, exceeded the UK governments recommended safe drinking limits, reported having cardiovascular disease or mental disorders, or were taking medications which might affect cardiovascular function. This resulted in twenty participants (13 male, 7 female) volunteering to participate in the study. Given the observational nature of the study, a case-control approach was adopted whereby the case comprised academic employees (n=10, 5 male: 5 female) with general employees forming the control group (n=10, 8 male; 2 female). All academic staff had teaching responsibilities and the job title of "lecturer" whilst general staff exclusively worked in a support capacity, as administrators or technicians. The academics

had a mean age of 40.6 ± 8.7 yrs and non-academics a mean age of 32.7 ± 5.8 yrs. All participants gave written informed consent, and approval for the study was obtained by the Research Ethics Committees of Edinburgh Napier University's Faculty of Health, Life and Social Sciences. All data collection occurred between the months of August and December 2012.

Measures

The Health and Safety Executives Management Standards Indicator and Analysis Tools

Full details of the development of the management standards indicator tool are available elsewhere (Cousins *et al.*, 2004). However, the indicator tool quantifies the following dimensions of work strain: Demands (workload, work pattern work environment); Control (autonomy over working practices); Managerial Support (encouragement and resources); Peer Support (colleague encouragement and support); Relationships (positive working, avoidance of conflict); Role (understanding of role or clarity, non-conflicting roles); and Change (how effectively is change managed and communicated). Participants indicate the extent to which various statements reflect their experiences at work over the preceding six month period, for example "I have a choice in deciding how I do my work" and "I have to work very intensively". Responses are provided on a 5-point scale: 1 (never), 2 (seldom), 3 (sometimes), 4 (often) and 5 (always). The Indicator Tool has a high level of reliability, with a goodness of fit index of 0.92 (Edwards, Webster, Van Laar, and Easton, 2008), and Chronbach's alpha values ranging from .78 to .87 for individual scales (Cousins, Mackay, Clarke, Kelly, Kelly, and McCaig, 2004). Responses were analysed using the management standards analysis tool (Health and Safety Executive, 2007), in accordance with the methodology previously reported by Houdmont *et al.* (2012). Scores range from 1 (poor) to 5 (desirable) for each of the seven dimensions, with lower scores representing greater risk exposure. The management standards analysis tool also compares the data with the UK

Health and Safety Executive benchmark data, from 136 organisations. This latter comparison provided a measure of risk for each dimension (excellent, good, poor, very poor).

Acute Psychosocial Demand

A visual analogue scale was used to measure the perceived strength of acute psychosocial demand as this method has previously been shown to provide a meaningful and useful assessment of occupational stress (Lesage and Berjot, 2011). The scale had a range of 100mm and was anchored at the midpoint by the term 'average demand', whilst 0mm and 100mm were labelled as representing "not at all" and "very" demanding days respectively. Scores obtained from the scale were then used to differentiate between the demands of the two days at an intra-individual level.

Physiological Measures

Ambulatory blood pressure and heart rate variability were measured using a combined ambulatory blood pressure monitor and electrocardiogram with a sampling rate of 200Hz (Cardiotens, Meditech, Budapest, Hungary). The five electrocardiogram leads were attached using Ambu Blue VLC long term monitoring electrodes (Ambu Ltd, St Ives, UK) at the following locations: left anterior axillary line, intercostals space 5 (x2), sternum, manubrium sterni, right anterior axillary line, intercostals space 5. The device was programmed to automatically obtain readings of diastolic and systolic blood pressure at 30 minute intervals. Participants also wore an Actiheart (Camntech, Cambridge) to measure physical activity over the assessment period. This is a one dimensional accelerometer which was programmed to measure activity levels at one minute intervals.

Procedures

Participants completed the UK Health and Safety Executive’s Management Standards Indicator tool once, prior to undergoing an ambulatory assessment of heart rate variability and blood pressure on two non-consecutive work days. The participants selected these days based upon the expectation that the two days would contain different levels of work-related demand. The distance between repeated measures was between 3 and 14 days. Upon arrival at the laboratory, participants were fitted with the monitoring device (Figure 1). The signal integrity was checked in real time and a manual blood pressure reading was performed in the laboratory to ensure integrity of the cuff prior to commencing the ambulatory recording. Participants were instructed to go about their working day as normal until their return to the laboratory, when the instrumentation was removed and data uploaded. To check whether the predicted differences in acute characteristics between days were present, at the end of each study day participants provided a rating of how demanding they perceived the day to have been on the visual analogue scale.

Analysis

Analysis of heart rate variability was performed using Cardiovisions software (Meditech, Hungary) which utilises a Fast Fourier Transformation. The following indices of heart rate variability were obtained: SDNN, low frequency (LF: 0.04-0.15Hz), high frequency (HF: 0.15-0.4Hz) and the ratio between these (LFHF ratio). Low frequency and high frequency values are expressed in both absolute terms and also as normalised units i.e. LFnu is equivalent to $LF / (LF+HF) \times 100$. The raw data were visually inspected and any periods of recording where a normal QRS complex could not be identified were marked accordingly and not included in any subsequent analysis. The period of ambulatory recording was manually selected and spectral analysis of heart rate variability was performed for both

frequency bands. Half hourly blood pressure readings were also obtained from the same software. Scores on the visual analogue scale were used to differentiate between the less and more demanding days at the individual level according to the participant's subjective appraisal.

Statistical analysis was performed using SPSS Version 20.0.0. As the Shapiro-Wilk test of normality revealed the heart rate variability data to be non-normally distributed, a logarithmic transformation was applied prior to further analyses. The effect of day upon perceived acute demand and physical activity were investigated with mixed ANOVAs, with job type entered as the between subjects factor. The effects of day upon blood pressure and parameters of heart rate variability were investigated by means of mixed ANOVA, with job type the "between" and day the "within" subject factors, controlling for the effects of age and gender. Partial Pearson's correlations were performed to determine the relationship between scores on the Management Standards Indicator Tool and physiological parameters (SBP, DBP, LF, HF, LFnu, HFnu, LFHF ratio and SDNN) controlling for the effects of age.

Results

Mean scores for the cohort as whole revealed varied levels of perceived exposure to psychosocial hazard for different dimensions of the management standards, according to the categorical scores provided by the UK Health and Safety Executive's analysis tool. Scores for the dimensions of, Demand, Control, Management Support and Peer Support fell within the "excellent" category (being at, above or close to the 80th percentile), relationships and control were categorised as being "good", whilst "role" received a "poor" score (below average but

above the 20th percentile) (Table 2). There was a clear group effect however, with general staff reporting “excellent” scores for all 7 dimensions of the indicator tool, whilst the academic group only achieved “excellent” scores for the Control and Peer Support. The academics also reported “poor” scores for both Management Support and Relationships, and fell in the “very poor” category (below the 20th percentile) for both Role and Change.

The average duration of ambulatory recording was 421 ± 39 minutes and all mean blood pressure readings were within normal ambulatory ranges (Mancia *et al.*, 1995). In all cases the participants correctly anticipated the respective demands of the two study days, as all individual scores on the visual analogue scale were greater on the more demanding day than on the less demanding day. The results of the ANOVA revealed there to be a significant main effect of day upon perceived acute demand $F(1, 18) = 4.59, p=.046$. There was also a significant main effect of day upon the following physiological measures: [SBP, $F(1,15) = 5.207, p=.038$] [HR, $F(1,15) = 5.749, p=.030$] [SDNN, $F(1,15) = 9.967, p=.007$] [LFnu, $F(1,15) = 18.339, p=.001$] [HFnu, $F(1,15) = 21.231, p=.001$] [LFHF ratio, $F(1,15) = 28.006, p=.001$] but not upon DBP, $F(1,15) = 3.214, p=.093$, or activity level, $F(1,15) = 2.265, p=.153$. There were no main effects of gender, age, or job type upon any measures of blood pressure, heart rate variability or activity. There was a significant interaction between the effects of day, gender and job type upon LFnu, $F(1, 15) = 5.555, p=.032$ and LFHF ratio, $F(1, 15) = 4.761, p=.045$. Post hoc tests revealed LFnu to be lower among male academics compared to female academics on the less demanding day $F(1, 15) = 5.029, p=.040$ but there was no difference between these groups on the more demanding day $F(1, 15) = 1.705, p=.211$. Similarly the LFHF ratio was reduced among male academics compared to female academics on the less demanding day, $F(1, 15) = 4.913, p=.043$ but there was no difference on the more demanding day $F(1, 15) = 1.097, p=.312$.

The Health and Safety Executive's Management Standards was found to be associated with SDNN: the relationships standard was positively associated with SDNN on both assessment days ($r=.467$, $p=.04$ and $r=.493$, $p=.03$ for the less and more demanding days respectively) while the demand standard was associated with SDNN on the less demanding day ($r=.632$, $p=.004$) (Table 5).

Discussion

The aim of the present study was to investigate exposure to psychosocial hazard in higher education employees and the physiological consequences of this exposure. It was hypothesised that: i) academics would be exposed to greater psychosocial hazard and would demonstrate less favourable work-time cardiovascular and autonomic responses than general staff; ii) scores on the management standards indicator tool would be positively associated with physiological stress responses i.e. greater perceived exposure to psychosocial hazard would result in greater blood pressure and reduced heart rate variability; and iii) blood pressure and measures of heart rate variability will differ according to the acute work-related demand of the measurement day. The study found that academics reported poorer scores than the general staff, but this was not reflected in work-time physiological functioning, as no differences were found in either blood pressure or heart rate variability between occupational groups. Therefore, the findings do not completely support the first hypothesis. As SDNN was associated with both the demand and relationships standards, the second hypothesis is at least partially supported. Additionally, daily work-related demands were shown to influence work-time ambulatory heart rate variability and blood pressure which supports the third hypothesis of the study.

This appears to be the first study to report upon the disparity in perceived exposure to psychosocial hazard according to role type amongst higher education employees, using the

Indicator Tool. The unfavourable scores reported by the academics for all dimensions of the management standards, with the exception of control and peer support, suggests that this group of employees may be subjected to potentially harmful levels of work-related psychosocial hazard. The dimension of Role received a particularly poor score, lending weight to the contention that academics are now expected to balance multiple roles and supporting previous findings regarding role as a source of stress amongst academics (Winter, Taylor and Sarros, 2000; Gillespie *et al.*, 2001; Kinman and Jones, 2003; Barret and Barret, 2007; Devenport, 2008). The “excellent” level of Control reported by both occupational groups is arguably indicative of the high level of autonomy that has historically been considered to characterise academic work, suggesting this favourable aspect of the job remains prominent, and also appears to extend to non-academic roles, possibly as a result of the institutional culture. Certainly, high levels of control have previously been reported amongst academics (Winter *et al.*, 2000; Winefield and Jarrett, 2001). Equally, the “excellent” level of Peer Support reported by academics and general staff alike may reflect the collegiate culture of academic institutions.

Despite reporting different levels of exposure to psychosocial hazard, academic and general staff did not differ in terms of their physiological response to work on either day. Although this may be artefactual, arising from the small sample size and subsequent lack of statistical power, there are also a number of possible theoretical explanations for this, the most simplistic being that different exposures to psychosocial hazard, as quantified by the Indicator Tool, do not significantly influence work-time autonomic functioning. Alternatively there may be a discrepancy between actual, or perceived, and reported psychosocial hazard, with academics reporting inflated exposures. However, neither of these explanations satisfactorily accommodates previous claims that the Indicator Tool has been associated with stress (Gyllensten and Palmer, 2005; Bevan *et al.*, 2010) and stress related health outcomes

(Kerr *et al.*, 2009). A final explanation, provided by the inter-individuality of the physiology underlying the stress response, may therefore be more plausible. It has previously been demonstrated that measures of heart rate variability can differ substantially between individuals (Thayer and Lane, 2007) and the present study certainly supports this inter-individuality in autonomic function. Additionally, Ilies *et al.* (2010) recently reported that a between-individuals analysis failed to find an association between workload and blood pressure, whilst the within-individual approach revealed positive associations between the two variables. Therefore, traditional cross-sectional analysis may not provide the optimal means of investigating the physiological response to work-related psychosocial hazard unless consideration is given to individual baseline values. However, this raises its own methodological challenges and it has yet to be established whether reactivity to acute laboratory stressors bears any correlation to reactivity during exposure to chronic, naturally occurring, stressors (Ho *et al.*, 2010).

Notwithstanding the somewhat rudimentary method of quantifying the acute demand of the two study days in relative terms, the present study demonstrated that perceived levels of daily work-related demand influence the physiological response to work amongst higher education employees. Kamark *et al.* (2005) have previously shown various dimensions of psychosocial stress, including “task demand”, to be associated with ambulatory blood pressure and cardiovascular risk while Ilies *et al.* (2010) found daily levels of negative effect to be associated with blood pressure over a ten day period. The present findings demonstrate that amongst higher education employees acute work-related demand influences both ambulatory blood pressure and heart rate variability in the expected direction. Although ambulatory blood pressure did not fall out with the normal range, even on the more demanding day, this should not necessarily be interpreted as evidence that increases in acute work-related demand are unlikely to represent a risk to long-term cardiovascular health. The levels of perceived

demand reported on the more demanding day were actually relatively moderate and it is certainly possible that exposure to greater levels of acute demand would result in greater elevations in blood pressure. Additionally, there is evidence that the relationship between cardiovascular outcome and blood pressure is continuous (Lewington *et al.*, 2002), with any elevation in blood pressure being potentially harmful. Similarly, despite the absence of clinical guidelines for heart rate variability it has been adequately demonstrated that decreased heart rate variability is associated with long term health risks. Fundamentally, diminished heart rate variability is indicative of a reduction in the flexibility and adaptability occurring within an individual's regulatory systems which are normal characteristics of healthy functioning (Shaffer *et al.*, 2014). As SDNN provides a measure of the fluctuation occurring across all of the factors contributing towards HRV, a reduction in this parameter can be interpreted as representing an unfavourable physiological response. This, coupled with an increased LFHF ratio on the more demanding day, suggests that exposure to greater demand may have long-term implications for the health of higher education employees.

Irrespective of potential long-term consequences, the variation in physiological function in response to acute demand is an important finding with significant methodological implications for future research. Such a finding suggests that in order to meaningfully interpret ambulatory physiological data, during work time at least, consideration must be given to the acute characteristics of the assessment day and how representative they are of the norm. Whilst this may seem somewhat obvious, given the main purpose of ambulatory monitoring is to obtain an assessment within the environment of interest, workplace investigations seldom attempt to quantify the acute psychosocial characteristics of the environment, beyond that which is typical to the specific occupation. However, there is a growing acknowledgement of the dynamic nature of the work environment (Ilies and Judge, 2002; Ilies and Judge 2004; Beal and Weiss, 2003; Ilies *et al.*, 2010) which should not be

overlooked for the sake of simplicity, and assessments should therefore be conducted on multiple work days of varying demand. Additionally, adopting such an approach would simultaneously go some way to addressing the issue of inter-individuality by enabling analysis to be performed at the within-individual level. Differences in individual physiological response across days of varying demand may potentially provide a more meaningful insight into the extent to which employees are coping with the demands of work than attempting to incorporate baseline values obtained in the laboratory.

According to the conceptual basis of the management standards, the dimensions which obtain the lowest categorical score could be considered to represent the greatest threat to employee health and wellbeing. In the present study role received the lowest categorical score, followed by relationships and change. Perhaps surprisingly then, role was found not to be associated with heart rate variability, suggesting that amongst higher education employees, high levels of role uncertainty may not directly influence autonomic function. The association between relationships and SDNN appears to be more in keeping with the underlying premise of the management standards, although the association between demands, which received an excellent score, and SDNN points to a more complex relationship. The demands dimension of the Indicator Tool has previously been shown to be a significant predictor of the subjective experience of stress (Gyllenstein and Palmer, 2005) and the present findings offer support for exposure to demand being implicated in the relationship between workplace strain and stress related ill-health. This may have potential implications for the interpretation of the management standards indicator tool or for prioritising workplace interventions, which has been identified as a consideration where respondents report poor scores across several dimensions of the Indicator Tool (Bevan *et al.*, 2010).

The possible interaction between chronic and acute exposure to psychosocial demand certainly lends further weight to the argument that the single-shot approach to

investigating the autonomic and cardiovascular response to the psychosocial work environment is limited by its inability to account for the dynamism that is inherent in many occupations. In conclusion this exploratory study provides initial evidence of a relationship between the management standards indicator tool and ambulatory heart rate variability and adds to the existing body of literature demonstrating that intra-individual variation in acute work-related demand is associated with altered blood pressure and heart rate variability. Further investigations should attempt to more accurately establish the interactions between the management standards, acute psychosocial demand and autonomic and cardiovascular functioning.

Limitations

Given the small sample size, and selective nature of the sample the results presented are of limited external validity. Additionally, the possibility of self-selection bias cannot be ruled out and, despite the protocol being designed to be as minimally invasive as possible, employees exposed to very high levels of psychosocial work-related hazard may be less likely to participate in research which places additional demands upon them during the working day. Additionally, as the participants selected the study days, it is reasonable to assume they may have deliberately precluded participation on days they anticipated being unusually high in acute demands, given the time required to have the instrumentation attached and removed. Certainly, variation in the acute demands of the two study days was relatively small, so the full extent of variation in autonomic function in response to acute demands may not have been captured by the present study.

References

- Barrett, L. and Barrett, P. (2007) Current practice in the allocation of academic workloads. *Higher Education Quarterly*, 61(4), 461-478.
- Beal, D. J. and Weiss, H. M. (2003) Methods of Ecological Momentary Assessment in Organizational Research. *Organizational Research Methods*, 6(4), 440-464.
- Bevan, A., Houdmont, J. and Menear, N. (2010) The management standards indicator tool and the estimation of risk. *Occupational Medicine*, 60, 525-531.
- Brown D. E., James, G. D. and Mills, P. S. (2006) Occupational differences in job strain and physiological stress: Female nurses and school teachers in Hawaii. *Psychosomatic Medicine*, 68, 524-530.
- Chandola, T., Britton, A., Brunner, E., Hemingway, H., Malik, M., Kumari, M., Badrick, E., Kivimaki, M. and Marmot, M. (2008) Work stress and coronary heart disease: what are the mechanisms? *European Heart Journal*, **29**, 640-648.
- Cousins, R., Mackay, C. J., Clarke, S. D., Kelly, C., Kelly, P. J., and McCaig, R. H. (2004) 'Management Standards' and work-related stress in the UK: Practical development. *Work and Stress*, 18(2), 113-136.

- Curtis, B. M. and O'Keefe, J. H. (2002) Autonomic tone as a cardiovascular risk factor: the dangers of chronic fight or flight. *Mayo Clinic Proceedings*, 77(1), 45-54.
- Demerouti, E., Bakker, A. B., Nachreiner, F. and Schaufeli, W. B. (2001) The job demands-resources model of burnout. *Journal of Applied Psychology*, 86(3), 499-512.
- Devenport, T. J., Biscombe, K. and Lane, A. M. (2008) Sources of stress and the use of anticipatory, preventative and proactive coping strategies by higher education lecturers. *Journal of Hospitality, Leisure, Sport and Tourism Education*, 7(1), 70-81.
- Education Reform Act (1998) London: HMSO
- Edwards, J. A., Webster, S., Van Laar, D. and Easton, S. (2008) Psychometric analysis of the UK Health and Safety Executive's Management standards work-related stress indicator tool. *Work and Stress: An International Journal of Work, Health and Organisations*, 22(2), 96-107.
- Etzkowitz, H., Webster, A., Gebhardt, C. and Terra, B. R. (2000) The Future of the University and the University of the Future: Evolution of Ivory Tower to 'Entrepreneurial Paradigm'. *Research Policy*, 29(2), 313-330.
- Fauvel, J. P., Quelin, P., Ducher, M., Rakotomalala, H. and Laville, M. (2001) Perceived Job Stress but not Individual Cardiovascular Reactivity to Stress Is Related to Higher Blood Pressure at Work. *Hypertension*, 38, 71-75.
- Filaire, E., Portier, H., Massart, A., Ramat, L. and Teixeira, A. (2010) Effect of lecturing to 200 students on heart rate variability and alpha-amylase activity. *European Journal of Applied Physiology*, 108, 1035-1043.

- Gillespie, N. A., Walsh, M., Winefield, A. H., Dua, J. and Stough, C. (2001) Occupational stress in universities: Staff perceptions of the causes, consequences and moderators of stress. *Work and Stress: An International Journal of Work, Health and Organisations*, 15(1), 53-72.
- Goldstein, D. S., Benth, O., Park, M. Y., Sharabi, Y. (2011) Low-frequency power of heart rate variability is not a measure of cardiac sympathetic tone but may be a measure of modulation of cardiac autonomic outflows by baroreflexes. *Experimental Physiology*, 96(12) 1255-1261.
- Guimont, C., Brisson, C., Dagenais, G. R., Milot, A., Vézina, M., Mâsse, B., Moisan, J., Laflamme, N. and Blanchette, C. (2006) Effects of job strain on blood pressure: A prospective study of male and female white collar-workers. *American Journal of Public Health*, 96(8), 1436-1443.
- Gyllenstein, K. and Palmer, S. (2005) The relationship between coaching and workplace stress: A correlational study. *International Journal of Health Promotion and Education*, 43(3), 97-103.
- Health and Safety Executive (2005). *HSE Management Standards Indicator Tool*. Available at:
<http://www.hse.gov.uk/stress/standards/pdfs/indicatortool.pdf>. [accessed 10th January 2013].
- Health and Safety Executive (2007) *HSE Management Standards Analysis Tool*. Available at:
<http://www.hse.gov.uk/stress/standards/downloads.html> [accessed 10th January 2013].

- Hjortskov, N., Rissén, D., Blangsted, A. K., Fallentin, N., Lundberg, U. and Søgaard, K. (2004) The effect of mental stress on heart rate variability and blood pressure during computer work. *European Journal of Applied Physiology*, 92, 84-89.
- Ho, R. C. M., Neo, L. F., Chua, A. N. C., Cheak, A. A. A. and Mak, A. (2010) Research on psychoneuroimmunology: does stress influence immunity and cause coronary artery disease? *Annals Academy of Medicine Singapore*, 39, 191-196.
- Houdmont, J., Kerr and Randall, R. (2012) Organisational psychosocial hazard exposures in UK policing: Management Standards Indicator Tool reference values. *Policing: An International Journal of Police Strategies and Management*, 35, 182-197.
- Houston, D., Meyer, L. H. and Paewai, S. (2006) Academic staff workloads and job satisfaction: expectations and values in academe. *Journal of Higher Education Policy and Management*, 28(1), 17-30.
- Huikuri, H. V. and Stein, P. K. (2012) Clinical application of heart rate variability after acutemyocardial infarction. *Frontiers in Physiology*, 3(41), 1-5.
- Ilies, R. and Judge, T. A. (2002) Understanding the dynamic relationships among personality, mood, and job satisfaction: a field experience sampling study. *Organizational Behaviour and Human Decision Processes*, 89, 1119-1139.
- Ilies, R. and Judge, T. A. (2004) An experience-sampling measure of job satisfaction and its relationships with affectivity, mood at work, job beliefs, and general job satisfaction. *European Journal of Work and Organizational Psychology*, 13(3), 367-389.
- Ilies, R., Dimotakis, N. and De Pater, I. E. (2010) Psychological and Physiological reactions to high workloads: implications for well-being. *Personnel Psychology*, 63, 407-436.

- Johnson, S., Cooper, C., Cartwright, S., Donald, I., Taylor, P. and Millet, C. (2005) The experience of work-related stress across occupations. *Journal of Managerial Psychology*, 20(2), 178-187.
- Johnson, J. V. and Hall, E. M. (1988) Job strain, work place social support, and cardiovascular disease: A cross-sectional study of a random sample of the Swedish working population. *American Journal of Public Health*, 78(10), 1336-1342.
- Kamarck, T. W., Schwartz, J. E., Shiffman, S., Muldoon, M. F., Sutton-Tyrell, K. and Janicki, D. L. (2005) Psychosocial stress and cardiovascular risk: what is the role of daily experience? *Journal of Personality*, 73(6), 1749-1774.
- Karasek, R. A. (1979) Job demands, job decision latitude, and mental strain: Implications for job redesign. *Administrative Science Quarterly*, 24, 285-308.
- Kerr, R., McHugh, M. and McCrory, M. (2009) HSE management standards and stress-related work outcomes. *Occupational Medicine*, 29, 574-579.
- Kikuya, M., Hozawa, A., Ohokubo, T., Ichiro, T., Michimata, M., Matsubara, M., Ota, M., Nagai, K., Araki, T., Satoh, H., Ito, S., Hisamichi, S. and Imai, Y. (2000) Prognostic significance of blood pressure and heart rate variabilities: The Ohasama study, *Hypertension*, 36, 901-906.
- Kinman, G. and Jones, F. (2003) 'Running up the down escalator': stressors and strains in UK academics. *Quality in Higher Education*, 9(1), 21-38.
- Kinman, G. and Court, S. (2010) Psychosocial hazards in UK universities: adopting a risk assessment approach. *Higher Education Quarterly*, 64(4), 413-428.
- Lam, A. (2010) From 'Ivory Tower Traditionalists' to entrepreneurial Scientists. Academic scientists in fuzzy university boundaries. *Social Studies of Science*, 40(2), 307-340.

Lesage, F. X. and Berjot, S. (2011) Validity of occupational stress assessment using a visual analogue scale. *Occupational Medicine*, 61(6), 434-436.

Lewington, S., Clarke, R., Qizilbash, N., Peto, R., Collins, R. (2002) Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet*. 14(360), 1903-1913.

Loerbroks, A., Schilling, O., Haxsen, V., Jarczok, M. N., Thayer, J. F. and Fischer, J. E. (2010) The fruits of ones labour: Effort-reward imbalance but not job strain is related to heart rate variability across the day in 35-44 year old workers. *Journal of Psychosomatic Research*, 69(2), 151-159.

Liao, D., Cai, J., Rosamond, W. D., Barnes, R. W., Hutchinson, R. G., Whitset, E. A., Rautaharju, P. and Heiss, G. (1997) Cardiac Autonomic Function and Incident Coronary Heart Disease: A Population-based Case-Cohort Study: The ARIC Study. American *Journal of Epidemiology*, 145(8), 696-706.

Lucini, D., Riva, S., Pizzinelli, P. and Pagani, M. (2007) Stress Management at the Worksite: Reversal of Symptoms Profile and Cardiovascular Dysfunction. *Hypertension*, 49, 291-297.

Mancia, G., Sega, R., Bravi, C., DeVito, G., Valagussa F., Cesana, G., and Zanchetti, A. (1995) Ambulatory blood pressure normality: results from the PAMELA study. *Journal of Hypertension*, 13(12), 1377-1390.

Rahman, F., Pechnik, S., Gross, D., Sewell, L., Goldstein, D. S. (2011) Low frequency power of heart rate variability reflects baroreflex function, not cardiac sympathetic innervation. *Clinical autonomic research*, 21(3), 133-134.

- 1
2
3 Reyes del Paso, G. A., Langewitz, W., Mulder, L. J., van Roon, A., Duschek, S. (2013) The
4 utility of low frequency heart rate variability as an index of sympathetic cardiac tone:
5 a review with emphasis on a reanalysis of previous studies. *Psychophysiology*, 50(5),
6 477-487.
7
8
9
10
11
12 Ryan, J. F., Healy, R. and Sullivan, J. (2012) Oh, won't you stay? Predictors of faculty intent
13 to leave a public research university. *Higher Education*, 63(4), 421-437.
14
15
16
17
18 Shaffer, F., McCraty, R. and Zerr, C. L. (2014) A healthy heart is not a metronome: an
19 integrative review of the heart's anatomy and heart rate variability. *Frontiers in*
20 *Psychology*, 5, 1040.
21
22
23
24
25 Siegrist, J., Siegrist, K. and Weber, I. (1986) Sociological concepts in the etiology of chronic
26 disease: the case of ischemic heart disease. *Social Science and Medicine*, 22(2), 247-
27 253.
28
29
30
31
32 Singh, J. P., Larson, M. G., Tsuiji, H., Evans, J. C., O'Donnell, C. J. and Levy, D. (1998)
33 Reduced heart rate variability and new-onset hypertension: insights into pathogenesis
34 of hypertension: The Framington Heart Study. *Hypertension*, 32, 293-297.
35
36
37
38
39
40 Task Force of The European Society of Cardiology and The North American Society of
41 Pacing and Electrophysiology. (1996) Heart rate variability. Standards of
42 measurement, physiological interpretation, and clinical use. *European Heart Journal*,
43 17, 354-381.
44
45
46
47
48
49 Thayer, J. F. and Lane, R. D. (2007) The role of vagal function in the risk for cardiovascular
50 disease and mortality. *Biological Psychology*, 74(2), 224-242.
51
52
53
54
55
56
57
58
59
60

- Thayer, J. F., Yamamoto, S. S. and Brosschot, J. F. (2010) The relationship between autonomic imbalance, heart rate variability and cardiovascular disease risk factors. *International Journal of Cardiology*, 2(28), 122-131.
- Tight, M. (2010) Are academic workloads increasing? The post-war survey evidence in the UK. *Higher Education Quarterly*, 64(2), 200-215.
- Tsuji, H., Larson, M. G., Venditti, F. S., Manders, E. S., Evans, J. C., Feldman, C. L. and Levy, D. (1996) Impact of reduced heart rate variability on risk for cardiac events. The Framington Heart Study. *Circulation*, 94, 2850-2855.
- Van Amselsvoort, L. G. P. M., Schouten, E. G., Maan, A. C., Sweenne, C. A. and Kok, F. J. (2000) Occupational determinants of heart rate variability. *International Archives of Occupational Health*, 73, 255-262.
- Van Egeren, L. F. (1992) The relationship between job strain and blood pressure at work, at home, and during sleep. *Psychosomatic Medicine*, 54(3), 337-343.
- Vrijkotte, T. G. M., van Doornen, L. J. P., and de Geus, E. J. C. (2000) Effects of work stress on ambulatory Blood Pressure, Heart Rate, and Heart Rate Variability. *Hypertension*, 35, 880-886.
- Winefield, A. H. and Jarrett, R. J. (2001). Occupational stress in university staff. *International Journal of Stress Management*, 8, 285-298.
- Winefield, A. H., Gillespie, N., Stough, C., Dua, J., Hapuarachchi, J. and Boyd, C. (2003) Occupational stress in Australian university staff: Results from a national survey. *International Journal of Stress Management*, 10(1), 51-63.

1
2
3 Winter, R., Taylor, T. and Sarros, J. (2000) Trouble at Mill: quality of academic worklife
4
5 issues within a comprehensive Australian university. *Studies in Higher Education*, 25(3),
6
7 279- 294.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Participant characteristics grouped by job type.

	Academics (n=10)	Non-Academics (n=10)
Age (years ± SD)	40.6 ± 8.7	32.7 ± 5.8*
Gender (M/F)	5/5	8/2
Physical Activity (METs ± SD)†	3055 ± 2281	6159 ± 6030
BMI (kg/m ² ± SD)	24.99 ± 4.06	22.48 ± 1.57
Work Ability ††	41.3 ± 3.64	44 ± 3.5

† Weekly METs calculated from the International Physical Activity Questionnaire. †† Self-reported Work Ability from the Work Ability Index (7-27= poor; 28-36=moderate, 37-43=good, 44-49=excellent). * Independent t-test revealed significant difference between groups (p<.05).

170x68mm (96 x 96 DPI)

Table 2 A comparison of mean Management Standards scores with benchmark data by occupational groups

Dimension	Academics (n=10)		General Staff (n=10)		All Staff (n=20)	
	Score	Category†	Score	Category	Score	Category
Demands	2.99	Poor	3.68	Excellent	3.33	Excellent
Control	3.95	Excellent	4.00	Excellent	3.98	Excellent
Management Support	3.46	Poor	4.06	Excellent	3.76	Excellent
Peer Support	3.95	Excellent	3.93	Excellent	3.94	Excellent
Relationships	3.65	Poor	4.15	Excellent	3.9	Good
Role	3.84	Very Poor	4.48	Excellent	4.16	Poor
Change	2.77	Very Poor	3.37	Excellent	3.07	Good
Global	3.51	††	3.96	††	3.73	††

† Category: derived from comparison with benchmark data from 136 organisations. Excellent: at, above or close to the 80th percentile. Good: Better than average but not yet at, above or close to the 80th percentile. Poor: Below average but above 20th percentile. Very Poor: below the 20th percentile - Urgent action needed. †† UK HSE do not provide categorisation of a global score.

176x101mm (96 x 96 DPI)

Table 3. Self-reported acute demand by occupation and self-reported chronic psychosocial demand.

Group	Less Demanding Day	More Demanding Day
All Staff (n=20)	3.64 ± 1.97	4.93 ± 1.93*
Academics (n=10)	3.27 ± 1.54	5.60 ± 2.56*
General Staff (n=10)	3.72 ± 1.99	4.57 ± 1.22*

* Mixed ANOVA revealed significant difference from less demanding day (p<0.05)

175x56mm (96 x 96 DPI)

Table 4. Ambulatory data for less demanding day and more demanding day presented by occupational group

	Academics (n=10)		General Staff (n=10)		All Staff (n=20)	
	Less Demanding	More Demanding	Less Demanding	More Demanding	Less Demanding	More Demanding
SBP	120.72 ± 7.67	125.08 ± 5.59	122.52 ± 6.28	127.66 ± 9.55	121.62 ± 6.89	126.37 ± 7.73*
DBP	75.18 ± 5.85	77.95 ± 5.63	76.48 ± 4.62	79.25 ± 4.51	75.83 ± 5.18	78.60 ± 5.01
HR	76.94 ± 10.61	84.81 ± 13.58	67.24 ± 6.67	68.34 ± 6.47	72.09 ± 9.96	76.58 ± 13.36*
SDNN	90.38 ± 33.21	77.35 ± 24.93	111.49 ± 25.06	98.58 ± 28.29	100.94 ± 30.61	87.97 ± 28.15**
LF	1497.9 ± 674.00	1857.5 ± 1150.0	2069.2 ± 1078.58	2096.8 ± 1036.3	1783.5 ± 923.10	1977.15 ± 72.45
LF _{nu}	74.80 ± 13.85	81.5 ± 10.15	74.0 ± 9.98	77.5 ± 7.28	74.4 ± 11.76	79.50 ± 8.83**
HF	558.50 ± 620.19	457.3 ± 489.07	757.0 ± 807.42	586.0 ± 333.71	657.75 ± 708.08	521.65 ± 412.81
HF _{nu}	24.10 ± 12.99	18.10 ± 9.10	24.6 ± 9.43	21.60 ± 6.88	24.35 ± 11.05	19.85 ± 8.00**
LFHF ratio	4.15 ± 2.32	6.25 ± 4.47	3.6 ± 1.89	4.33 ± 2.55	3.87 ± 2.08	5.29 ± 3.67**
Activity†	34.29 ± 9.45	49.03 ± 18.59	44.50 ± 40.17	47.96 ± 32.44	39.40 ± 28.88	49.49 ± 25.74

* Significantly different from less demanding day (Mixed ANOVA controlling for age and gender, $p < 0.05$). SBP: systolic blood pressure, DBP: diastolic blood pressure, HR: heart rate, LF: low frequency power, LF_{nu}: low frequency power in normalized units, HF: high frequency power, HF_{nu}: high frequency power in normalized units, LFHF ratio: low to high frequency ratio. † Mean activity counts calculated over period of ambulatory assessment from.

178x132mm (96 x 96 DPI)

Table 5. Partial Correlations between HSEMS and Heart Rate Variability adjusted for age (n=20)

	Time	Global	Demands	Control	M Support	P Support	Relationships	Role	Change
LF (log10)	LD	.092	.104	.075	-.091	.062	.272	-.048	.176
	MD	-.174	-.044	-.180	-.346	-.017	-.014	-.213	.071
HF (log10)	LD	.125	.310	.193	-.089	.067	.309	.010	-.123
	MD	-.074	.152	-.119	-.268	.013	.153	-.140	-.069
LFHF ratio (log10)	LD	-.071	-.431	-.263	-.077	-.058	-.286	-.148	.266
	MD	-.114	-.447	-.078	-.050	-.162	-.449	-.003	.115
SDNN (log10)	LD	.295	.632**	.171	.020	.098	.467*	.045	-.029
	MD	.210	.436	.120	-.036	.140	.493*	-.045	.068

. LD = Less Demanding Day. MD = More Demanding Day. Δ = Change from LD to MD. *Significant at p < 0.05, ** significant at P < 0.01, Partial Pearson's correlation controlling for effect of age.

180x99mm (96 x 96 DPI)

Figure 1. Participant instrumented with both a Cardiotens combined ABPM and ECG monitor and an Actiheart monitor.

76x111mm (96 x 96 DPI)