



UWS Academic Portal

The importance of behaviour in improving the production of shrimp in aquaculture

Bardera, Guillermo; Usman, Nafiha; Owen, Matthew; Pountney, Daniel ; Sloman, Katherine A.; Alexander, Mhairi E.

Published in:
Reviews in Aquaculture

DOI:
[10.1111/raq.12282](https://doi.org/10.1111/raq.12282)

E-pub ahead of print: 22/08/2018

Document Version
Peer reviewed version

[Link to publication on the UWS Academic Portal](#)

Citation for published version (APA):

Bardera, G., Usman, N., Owen, M., Pountney, D., Sloman, K. A., & Alexander, M. E. (2018). The importance of behaviour in improving the production of shrimp in aquaculture. *Reviews in Aquaculture*, 11(4), [12282]. <https://doi.org/10.1111/raq.12282>

General rights

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact pure@uws.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

"This is the peer reviewed version of the following article: Bardera, Guillermo et al. (2018) *The importance of behaviour in improving the production of shrimp in aquaculture*. Reviews in Aquaculture, Vol.X. pp. xx. which has been published in final form at <https://doi.org/10.1111/raq.12282>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions."

<https://authorservices.wiley.com/author-resources/Journal-Authors/licensing/self-archiving.html>

To be completed on final publication.

1 **The importance of behaviour in improving the production of shrimp in aquaculture**

2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

Guillermo Bardera^{1*}, Nafiha Usman², Matthew Owen², Daniel Pountney², Katherine A. Sloman¹, Mhairi E. Alexander¹

¹Institute of Biomedical and Environmental Health Research, School of Health & Life Sciences, University of the West of Scotland, Paisley, Scotland, PA1 2BE, U.K.

²Skretting Aquaculture Research Centre (ARC), Stavanger, 4016, Norway.

*Corresponding author: guillermo.bardera@uws.ac.uk
+44 7563743625

Running title: Shrimp behaviour in aquaculture

35 **ABSTRACT**

36

37 There is increasing recognition within the aquaculture industry that understanding the
38 behaviour of farmed animals can help provide solutions to feeding problems. However, most
39 studies have focused on finfish production, with fewer behavioural studies on feeding
40 processes in commercially-produced crustaceans. More than 60% of crustacean aquaculture
41 is attributed to the production of penaeids, particularly the Pacific white-leg shrimp
42 (*Litopenaeus vannamei* Boone). The profitability of the Pacific white-leg shrimp for
43 aquaculture stems from its ability to survive in a wide range of environments and its fast
44 growth at high densities. However, there are significant setbacks within their farming. In
45 particular, while they can move rapidly to take food pellets, they can be slow to consume
46 them leading to food wastage and subsequent economic losses for the industry.
47 Understanding shrimp behaviour provides a starting point for refinements to feeding
48 practices. Here we review the different influences on shrimp behaviour which are likely to
49 influence productivity such as individual-level effects (e.g. moulting, sex), environmental
50 influences (e.g. photoperiod, conspecific presence) and water quality (e.g. salinity,
51 temperature). Although work on feed management has been conducted, providing
52 information on nutrition, feeding frequency and schedules, here we demonstrate that such
53 advances must be accompanied by behavioural approaches to allow development of optimal
54 feeding efficiencies and to support the continued growth of the crustacean aquaculture
55 industry.

56

57

58 **Key words:** Aquaculture, Chemoattraction, Crustacean, Feeding behaviour, Feeding
59 effectors, *Litopenaeus vannamei*.

60 INTRODUCTION

61

62 The production of crustaceans in aquaculture is a global industry with large commercial and
63 economic importance (Bondad-Reantaso *et al.* 2012). Marine shrimp in particular dominate
64 crustacean aquaculture (Briggs *et al.* 2004) and of the 7 million tonnes produced annually,
65 more than 60% is attributed to the production of penaeid shrimp (FAO 2016). Many countries
66 rely heavily on such shrimp production and as a result it has been one of the fastest growing
67 aquaculture sectors in Asia and America (Briggs *et al.* 2004; Funge-Smith & Briggs 2005;
68 FAO 2016).

69

70 Modern day shrimp aquaculture originated in Japan in the 1930s (Rosenberry 2001;
71 Chamberlain 2010), however technologies developed in the 1970s expanded production in
72 other Asian countries, such as Indonesia, the Philippines and Thailand as well as the
73 Americas (Rosenberry 2001; Chamberlain 2010; Nash 2011). Initially the main focus of this
74 development was the culture of the tiger shrimp (*Penaeus monodon* Fabricius) (Yi *et al.*
75 2016), for which there was a large demand in Asian and US markets (Nash 2011; Rimmer *et*
76 *al.* 2013). However, the intensification and densification of *P. monodon* farming led to
77 outbreaks of disease in the 1990s that severely depleted aquaculture stocks over a decade
78 (Flegel 2009). As a result, the culture of Pacific white-leg shrimp (*Litopenaeus vannamei*
79 Boone) was introduced (Hall 2004; Flegel 2009; Chamberlain 2010), and it is now the most
80 heavily farmed shrimp species representing around 80% of total marine penaeid shrimp
81 production (FIGIS 2015).

82

83 The focus on *L. vannamei* production was partly due to advantages over *P. monodon* with
84 regards to disease resistance (Liao & Chien 2011; Bondad-Reantaso *et al.* 2012).

85 Additionally, the species has an increased tolerance to high stocking densities (typically 60-
86 150/m², but up to 400/m²) (Briggs *et al.* 2004; Funge-Smith & Briggs 2005; Jory & Cabrera
87 2012), improved feed conversion rates and low protein requirements (of around 20-35%
88 crude protein) (Briggs *et al.* 2004; Funge-Smith & Briggs 2005; Bondad-Reantaso *et al.*
89 2012; Jory & Cabrera 2012), higher average daily growth rates (up to 3 g/week) (Briggs *et al.*
90 2004; Funge-Smith & Briggs 2005; Bondad-Reantaso *et al.* 2012; Jory & Cabrera 2012),
91 high tolerance to a large range of water parameters including salinity and temperature (Briggs
92 *et al.* 2004; Funge-Smith & Briggs 2005; Bondad-Reantaso *et al.* 2012; Jory & Cabrera
93 2012), and higher survival rates during larval rearing (50 – 60%) (Funge-Smith & Briggs
94 2005). Moreover, in marketing terms, *L. vannamei* has been shown to be preferred by
95 consumers over *P. monodon* (Funge-Smith & Briggs 2005).

96

97 As a result of this intensification of production, *L. vannamei* has been the subject of a large
98 number of scientific studies that has resulted in substantial knowledge on the biology (e.g.
99 Chamberlain & Lawrence 1981; Dall *et al.* 1990; Vega-Villasante *et al.* 2000), genetics (e.g.
100 Garcia *et al.* 1994; Gitterle *et al.* 2005) and zootechnics (e.g. Briggs *et al.* 2004; Cuéllar-
101 Anjel *et al.* 2010) of the species. Feeding efficiency of farmed stock has been an important
102 focus, however there is a general lack in information of individual feeding behaviour and the
103 aspects that may be important for determining optimal feeding efficiency in *L. vannamei*, as
104 well as in other commercially important species. This paper therefore, with a specific focus
105 on *L. vannamei*, reviews the current problems associated with the feeding of farmed shrimp
106 and how understanding their behaviours may promote greater feeding efficiency and
107 subsequent increases in production. By drawing on behavioural information from
108 neurophysiological and nutritional studies which have been developed in *L. vannamei* as well
109 as other commercial and non-commercial crustacean species, we aim to present

110 considerations that we believe to be important for detecting and developing optimal feeding
111 practices.

112

113

114 **FEEDING PROBLEMS IN SHRIMP AQUACULTURE**

115

116 Crustacean aquaculture continues to experience global growth (Bondad-Reantaso *et al.*
117 2012). However, there are a range of factors that contribute to inefficiencies in farming
118 practices leading to reduced production as well as significant economic losses (Neiland *et al.*
119 2001; Engle *et al.* 2017). Common problems include poor larvae production across a range of
120 farmed species (Juinio-Menez & Gotanco 2004; Thuy & Ngoc 2004; Vijayakumaran *et al.*
121 2009; Jeffs 2010), the incidence of disease (Rodríguez *et al.* 2003; Flegel 2009), pollution
122 effects due to waste products (Briggs & Funge-Smith 1994; Martin *et al.* 1998; Boyd *et al.*
123 2007), and the escape of individuals (Cuéllar-Anjel *et al.* 2010; Felix 2013). Of particular
124 interest here, however, are the studies that have documented the feeding inefficiencies
125 associated with crustacean production (e.g. Chanratchakool *et al.* 1998; Davis *et al.* 2006;
126 Boyd *et al.* 2007; Smith & Tabrett 2013), which are predominantly in relation to expensive
127 feed production costs, feeding-management practices as well as poor feeding efficiency by
128 individuals.

129

130 The manufacture of feed is one of the largest costs in aquaculture production (Lawrence &
131 Lee 1997; Boyd *et al.* 2007; Smith & Tabrett 2013; Zhou *et al.* 2017), and depending on the
132 system in question, may represent as much as 60-80% of total production costs
133 (Chanratchakool *et al.* 1998; Hertrampf & Piedad-Pascual 2000; Smith *et al.* 2002; Cuzon *et*
134 *al.* 2004; Rola & Hasan 2007; Hasan & New 2013). Commercial shrimp feed formulations

135 commonly include fish meal, often representing the primary and most expensive protein
136 ingredient (Lim & Dominy 1990; Tacon & Barg 1998; Dersjant-Li 2002; Cruz-Suárez *et al.*
137 2007). There are clear beneficial effects of marine protein sources for crustacean nutrition,
138 such as the presence of essential fatty acids, vitamins, cholesterol and minerals, in addition to
139 the enhancement of feeding activity, palatability and growing processes through attractants
140 and growth factors (Molina-Poveda *et al.* 2002; Cruz-Suárez *et al.* 2007; Sookying *et al.*
141 2013). However, the current state of global fish stocks has triggered a costlier production of
142 fish meal (Hardy & Tacon 2002; Sookying *et al.* 2013), and as a result, less expensive plant
143 and terrestrial animal proteins are being increasingly used in aqua-feed formulations since
144 they are readily available, economical, and sustainable (Lim & Dominy 1990; Watanabe
145 2002; Samocha *et al.* 2004a; Amaya *et al.* 2007; Gatlin *et al.* 2007; Naylor *et al.* 2009;
146 Sookying *et al.* 2013). Moreover, studies specifically on *L. vannamei* suggest that it is
147 possible to replace around 75-100% of the total amount of crude protein with proteins of non-
148 marine origin, without compromising shrimp performance (Davis & Arnold 2000; Fox *et al.*
149 2004; Hernández *et al.* 2004; Samocha *et al.* 2004a; Amaya *et al.* 2007).

150

151 A major restriction in shrimp aquaculture is often an inability to observe stock during its
152 development, causing difficulties for determining biomass and survival (Davis *et al.* 2006;
153 Smith & Tabrett 2013). Feed management is therefore a complex task during shrimp
154 production, often resulting in overfeeding and the generation of large amounts of waste
155 (Martinez-Cordova *et al.* 1998; Nunes & Parsons 1999; Davis *et al.* 2006; Sun *et al.* 2016).
156 Overfeeding can lead to high levels of non-beneficial microorganisms in addition to hypoxic
157 conditions at certain times within the day, therefore shrimp feeding protocols must be well
158 organised in time as well as space and in accordance with population size (Cuéllar-Anjel *et al.*
159 *al.* 2010). In addition, feed waste may be compounded by poor physical quality of feed and/or

160 environmental conditions (see environmental and water quality effects sections), which
161 negatively influence crustacean sensorial capacity and food detection. Poor physical quality
162 of feed can lead to premature breakdown of food (Davis *et al.* 2006) and reduce food
163 digestibility through nutrient leaching (Sanchez *et al.* 2005; Cruz-Suárez *et al.* 2007).
164 Spoilage of food combined with poor environmental conditions could also result in
165 oversaturation of chemoreceptors (Lee & Meyers 1996; De la Haye *et al.* 2012), and
166 disruption of chemical cues within the water (Lee & Meyers 1996; Nunes *et al.* 2006).
167 Studies exist on the relationships between shrimp species, stock mass and size, and the
168 correct amount of feed required for optimal production (Jory 1995; Jory & Cabrera 2012).
169 However, the difficulties in checking development of the whole stock can make dispensing
170 the optimal quantity of feed difficult. The use of feeding trays or automatic feeders in shrimp
171 farming is one solution to this problem as they are useful for the control and monitoring of
172 the amount of feed consumed by shrimp, and thus reduce the incidence of overfeeding
173 (Martinez-Cordova *et al.* 1998; Bador 2000; Pontes & Arruda 2005b; Molina & Espinoza
174 2018). In addition, automatic feed dispensers using passive acoustic systems (hydrophones)
175 can measure the quantity of feed consumed by shrimp which provides a more accurate
176 estimate of the current feeding demand in ponds when compared to feeding trays (Smith &
177 Tabrett 2013). This in turn can improve the environmental conditions of the production ponds
178 (Seiffert & Andreatta 2004; Cuéllar-Anjer *et al.* 2010). However, for this to be a truly
179 effective solution there needs to be high feeding efficiency of individuals, which can be
180 confirmed by the collection of behavioural data (Zhou *et al.* 2017).

181

182

183 **CRUSTACEAN FEEDING BEHAVIOUR**

184

185 We suggest here that in order to address the problems associated with feeding efficiencies, a
186 focus on individual feeding behaviours in crustaceans is necessary. Therefore, an important
187 step is to outline the mechanisms involved in food detection and to link shrimp physiology to
188 behaviour. Crustaceans are considered to be generalist feeders (Smith *et al.* 2005; Hay 2011).
189 They must identify chemical cues of potential food, distinguishing relevant cues against
190 background chemicals and determining their qualities (i.e., molecular structures), quantities
191 (i.e., concentrations), and locations (i.e., spatiotemporal dynamics) (Lee & Meyers 1996;
192 Moore & Grills 1999; Derby *et al.* 2001). They respond to primary metabolites (e.g. sugars,
193 proteins, amino acids, products of respiration, metabolic wastes) because these indicate the
194 presence of food resources in general (Carr 1978; Zimmer-Faust 1987; Hay 2011).

195

196 In the case of decapod crustaceans, such as *L. vannamei*, there are three different detection
197 paths for locating a food source that include the visual, mechanoreceptor and chemoreceptor
198 systems, with associated receptors which are present on most shrimp appendages. However,
199 decapods perceive food mainly via chemical stimuli rather than mechanoreception or vision
200 as water is an excellent medium to transmit these types of cues (Hindley 1975; Atema 1995;
201 You *et al.* 2006). Crustaceans use chemoreception to intercept food across longer distances
202 and once in closer proximity to the food source they can apply visual and mechanoreception
203 detection. As such, these two pathways of detection act to obtain additional information in
204 relation to feeding processes (Derby & Sorensen 2008).

205

206 **Chemosensory receptors**

207 Crustaceans possess a large number of chemoreceptors that are packaged into sensilla that are
208 simple sensory receptors consisting of modified hair-shaped cells (Derby & Sorensen 2008).
209 Sensilla are extremely diverse in structure and are distributed over the body on antennules,

210 but also on the second antennae, legs, mouthparts, cephalothorax, abdomen and telson
211 (Gleeson *et al.* 1993; Cate & Derby 2001; Horner *et al.* 2004; Derby & Sorensen 2008). This
212 wide distribution allows crustaceans to maximise their ability to locate and discriminate
213 environmental chemical cues and physical information (Steullet *et al.* 1999; Horner *et al.*
214 2000; Derby & Steullet 2001).

215

216 Each antennule has hundreds of thousands of chemoreceptor neurons (Grünert & Ache 1988;
217 Cate & Derby 2001) and is sensitive to many different chemicals, in particular small, water-
218 soluble molecules such as amino acids, amines, nucleotides, and sometimes sugars and
219 peptides (Ache 1982; Carr *et al.* 1984; Carr & Derby 1986). Antennular chemoreception is
220 specialized for detection of chemical cues (Rittschof 1992; Koehl 2011) and plays an
221 important role in search initiation and orientation toward the source of a distant chemical
222 stimulus (Reeder & Ache 1980; Devine & Atema 1982; Derby 2000; Derby *et al.* 2001). The
223 majority of chemoreceptors are found on the non-aesthetasc/non-olfactory regions on the
224 medial and lateral flagella of the antennules (Schmidt *et al.* 1992; Cate & Derby 2001;
225 Steullet *et al.* 2002; Horner *et al.* 2004; Schmidt & Derby 2005). In addition, crustaceans
226 possess a tuft on the distal part of the lateral antennular flagella which is composed of
227 chemoreceptor sensilla, denominated as the aesthetasc/olfactory region (Grünert & Ache
228 1988; Derby *et al.* 2001; Steullet *et al.* 2002; Horner *et al.* 2004), and is the most intensively
229 studied part of crustacean chemosensory pathways (Grasso & Basil 2002; Derby & Sorensen
230 2008). Studies have demonstrated that aesthetasc and non-aesthetasc regions have some
231 functional redundancy. For example, work with spiny lobster (*Panulirus argus* Latreille)
232 (Reeder & Ache 1980; Derby *et al.* 2001; Steullet *et al.* 2001; Schmidt & Derby 2005),
233 American lobster (*Homarus americanus* H. Milne-Edwards) (Devine & Atema 1982) and
234 crayfish (*Cambarus bartonii* Fabricius) (Dunham *et al.* 1997) has revealed that after selective

235 removal of their aesthetasc sensilla, these animals can compensate for the loss if their non-
236 aesthetasc chemoreceptors are intact. Therefore, both kinds of receptors assess the quality of
237 food, mediate olfactory learning and enable orientation to distant food sources (Derby *et al.*
238 2001; Steullet *et al.* 2001, 2002; Horner *et al.* 2004).

239

240 Chemical receptors located on other parts of the body are also important. Pereopod dactyl
241 chemoreceptors control local grasping reflexes and food recognition (Fuzessery & Childress
242 1975; Zimmer-Faust & Case 1983; Dunham *et al.* 1997). Walking legs may play a leading
243 role in food detection if the crustacean has lost the antennules (Hazlett 1971). Maxillipeds
244 and mouthpart chemoreceptors mediate the decision to ingest food (Derby & Atema 1982;
245 Derby *et al.* 2001; Aggio *et al.* 2012), whereas pleopod beating produces a water current
246 around the shrimp carrying organic molecules. This activity may provide additional
247 information via chemoreception on the presence of prey, conspecifics or predators (Devine &
248 Atema 1982; Costero & Meyers 1993; Atema 1995; Klages *et al.* 2002), however, the precise
249 role of these chemoreceptors on the parts of the body detailed here is poorly understood.

250

251 In addition to all chemical receptors which mediate food recognition and ingestion, a recent
252 study has identified a novel pathway of branchial nutrient absorption in the green shore crab
253 (*Carcinus maenas* Linnaeus). This phenomena is the first finding in marine arthropods
254 (Blewett & Goss 2017) and may provide a new level of complexity to crustacean nutrition.

255

256 **Behavioural responses**

257 Chemosensory stimuli mediate all phases of feeding in crustaceans (Holland & Borski 1993;
258 Grey *et al.* 2009), and these can be categorised into a series of behavioural responses such as
259 detection and orientation towards a food source (Ache 1982; Kurmaly *et al.* 1990; Lee &

260 Meyers 1996; Moore & Grills 1999) and handling of food (Steiner & Harpaz 1987; Lee &
261 Meyers 1996; Derby *et al.* 2016). In general, when feed is offered, crustaceans change their
262 behavioural profile and there are observed increases in the frequency of behaviours related to
263 searching for feed, such as exploration and crawling, and feed grabbing (Da Costa *et al.*
264 2016). This is coupled with a decrease in the frequency of those behaviours not associated
265 with feeding, such as cleaning and burying (Pontes & Arruda 2005a; Silva *et al.* 2012).
266 Crustaceans have also been noted to express habituation behaviours, including learning to
267 search in a particular area or type of habitat (Shuranova *et al.* 2005), to capture and handle
268 food more efficiently (Derby & Atema 1981), and developing a change in acceptability of
269 food (Derby & Atema 1981; Steiner & Harpaz 1987; Daniel & Derby 1988). These
270 habituation behaviours suggest a potential application in conditioning farmed shrimp to the
271 use of automatic feed dispensers as has been observed in finfish aquaculture (e.g. Bratland *et*
272 *al.* 2010; Zion *et al.* 2010, 2011a,b; Folkedal *et al.* 2018).

273

274 In crustaceans, during the first feeding behaviour phases of detection and orientation towards
275 a food source, chemoreceptors on the antennules have been shown to have important roles
276 (Ache 1982; Lee & Meyers 1996). This has been evidenced through the observations of
277 antennule flicking behaviours, where there is rapid movement of the antennules through the
278 surrounding fluid, allowing chemicals to bind repeatedly to receptor sites (e.g. Schmitt &
279 Ache 1979; Derby & Atema 1982; Carr & Derby 1986; Zimmer-Faust 1991; Koehl 2006;
280 Thiel & Breithaupt 2011). A number of studies have observed the greatest antennule flicking
281 frequency at the onset of a stimulus, and is therefore an important feeding behaviour as it
282 indicates chemical perception (e.g. Price & Ache 1977; Schmitt & Ache 1979; Reeder &
283 Ache 1980; Devine & Atema 1982; Allison *et al.* 1992; Gleeson *et al.* 1993). Changes to
284 flicking frequency have been shown to improve the exchange of chemical substances, as

285 flicking movements enhance the discharge of the chemoreceptors, allowing new chemical
286 substances to attach again to the chemoreceptor binding sites (Snow 1973; Schmitt & Ache
287 1979; Atema 1985; Gleeson *et al.* 1993). Decreased flicking may therefore denote an
288 impaired or reduced ability to detect the chemical stimulus and therefore the presence of food
289 (Allison *et al.* 1992; Krång & Rosenqvist 2006; De la Haye *et al.* 2011).

290

291 Antennular flicking in response to chemostimulation is a wide-spread behaviour among
292 crustacean species. For example, increased rates in flicking have been recorded in the giant
293 freshwater prawn (*Macrobrachium rosenbergii* De Man) (to a maximum of 40 flicks/min)
294 when presented with feed with high betaine concentrations (Harpaz & Steiner 1990). Devine
295 and Atema (1982) found that 96% of lobsters (*H. americanus*) maintaining antennular
296 flicking walked in a straight line towards a food source. Reeder and Ache (1980) observed
297 circular movements in the spiny lobster (*P. argus*) when one antennule was ablated.
298 Antennular flicking frequencies of the Dungeness crab (*Metacarcinus magister* Dana) were
299 observed to treble from 20-40 flicks/min to 60-120 flicks/min when individuals were
300 stimulated with different concentrations of clam samples added to the water (Pearson *et al.*
301 1979). Schmitt and Ache (1979) concluded from the same species that an increase in
302 antennular flicking upon detection of a stimulus improved the temporal resolution of
303 chemical sources in the environment. Antennular flicking however varies among crustacean
304 species. Work on the spiny lobster (*P. argus*) observed antennular flicking activity rates
305 between 25 – 90 flicks/min (Daniel & Derby 1991; Goldman & Koehl 2001), whereas
306 Allison *et al.* (1992) observed a maximum rate of 20 flicks/min in the crayfish (*C. bartoni*).
307 The hermit crab (*Pagurus bernhardus* Linnaeus) has been shown to perform flicking rates as
308 high as 140-160 flicks/min (De la Haye *et al.* 2012).

309

310 An increase in flicking activity, however, is not always necessary and when water moves in a
311 steady flow, crustacean species have been observed to reduce the movement of their
312 antennules (Snow 1975; Goldman & Koehl 2001). Flow type of water can be important and
313 studies with American lobster (*Homarus gammarus* Linnaeus) (Moore *et al.* 1991; Mjos *et al.*
314 1999) and spiny lobster (*P. argus*) (Wilkins *et al.* 1996; Horner *et al.* 2004) under different
315 turbulent flow regimes observed that antennules change position accordingly. In addition to
316 changes in water flow, possible interferences with chemoreception by substances such as
317 manganese (Engdahl 1997; Krång & Rosenqvist 2006) or sub-optimal culture parameters (i.e.
318 low pH conditions) (Allison *et al.* 1992; De la Haye *et al.* 2011, 2012) can cause changes in
319 antennular flicking activity in crustaceans. Moreover, work with stomatopods, such as mantis
320 shrimp, has shown an increase of antennule flicking activity under low light conditions, when
321 there would be less reliance on visual cues (Cheroske *et al.* 2009). Therefore, such variations
322 in flicking behaviours can produce confusion for researchers identifying behavioural
323 responses towards experimental feeds or additives, and whether affinity between feed and
324 antennule flicking activity is directly related to feeding attraction or if it is moderated by
325 other reasons.

326

327 Variation in antennular flicking activities may be a result of different water sampling
328 techniques made by crustaceans. For example, in lobsters, rapid downward motions of
329 antennules may be due to the removal of water previously sampled from the aesthetasc
330 receptors to allow detection of new chemical-bearing water (Goldman & Koehl 2001;
331 Schmidt & Derby 2005). Another characteristic of flicking activity is that, in some cases, it is
332 dependent on crustacean size. Studies on stomatopods found that larger individuals have
333 more rapid chemical diffusion on aesthetasc surfaces than smaller animals (Mead *et al.*
334 1999). However, this is likely to be species-specific (Goldman & Koehl 2001).

335 In penaeid shrimp, a limitation in observing the use of antennular flicking is the small size of
336 the shrimp's antennules compared with those of other species, such as lobsters (Pittet *et al.*
337 1996). However, studies with small hermit crabs (*P. bernhardus*) (3 – 4 cm carapace length)
338 found antennular flicking to be a tractable measure of chemo-responsiveness (De la Haye *et*
339 *al.* 2012) which demonstrates the potential for observations of *L. vannamei* in similar kinds of
340 studies. The structure and function of the olfactory organs of crustaceans, as a group, is
341 highly congruent (Hallberg *et al.* 1992), therefore, any observed effect in other species such
342 as *H. americanus*, *P. argus* or *P. bernhardus* is likely to occur in other marine crustaceans in
343 a similar way. Nevertheless, there is little information about antennular flicking activity in
344 penaeid shrimp. Moreover, shrimp exhibit more movements of their mouthparts and less of
345 their antennules when compared with both lobsters and crabs (Pittet *et al.* 1996).

346

347 Recognition of food implies its discrimination from non-food items (Hindley 1975), and the
348 elicitation of food-acceptance or food-rejection behaviour is seemingly dependent on a set of
349 chemoreceptors in the dactyl receptors and in the mandibular-oral area (Hazlett 1971;
350 Fuzessery & Childress 1975; Hindley 1975, Robertson *et al.* 1981). Once the crustacean is
351 close to the food source, it makes contact with the food item with the dactyl of the walking
352 leg and moves it towards the maxillipeds and mouthparts. The dactyls of the first two pairs of
353 walking legs bear chemotactic organs which aid with the acceptance process (Ameyaw-
354 Akumfi 1977). Oral handling of feed via the maxillipeds is generally indicated by the
355 disappearance of the food item from the mouthparts and its appearance in the stomach, which
356 can often be observed (Steiner & Harpaz 1987). However, the onset of a feed-handling
357 behavioural pattern does not guarantee that the shrimp will actually go on to consume the
358 feed (Harpaz & Steiner 1987). After feeding, crustaceans present a stereotypic behaviour
359 referred to as antennular grooming (Zimmer-Faust *et al.* 1984; Barbato & Daniel 1997;

360 Wroblewska *et al.* 2002; Schmidt & Derby 2005). This behaviour has been attributed to the
361 activation of aesthetascs regions through the antennules wiping against maxillipeds and the
362 rubbing movements of maxillipeds against themselves (“auto-grooming”) (Barbato & Daniel
363 1997; Wroblewska *et al.* 2002).

364

365 In many species feed rejection behaviours follow a well-defined behavioural pattern. For
366 example, *M. rosenbergii* individuals that go on to reject feed have been observed to make
367 regular movements around food items, with little interaction. Also, any individuals that
368 handled feed in their maxillipeds, made no movements of the mouthparts (Steiner & Harpaz
369 1987). The hermit crab (*Pagurus granosimanus* Stimpson) has been observed in general to
370 flick the second antennae back and away after contacting the chemical source when food
371 rejection behaviour follows (Wight *et al.* 1990). Food is then pushed away energetically with
372 the chelipeds and the crab backs away from it. In other instances, *P. granosimanus* has been
373 observed to grasp feed hesitantly with the chelipeds before rejecting it.


374

375

376 **INFLUENCES ON SHRIMP FEEDING BEHAVIOUR**

377

378 Understanding shrimp behaviour provides a starting point for refinements to feeding
379 practices, however, there are many influences on shrimp behaviour which are likely to be
380 species and context-specific (Fig. 1). These can be broadly divided into individual level
381 effects, environmental effects and water quality effects.



382

383 **Figure 1.** The influences on shrimp behaviour that are considered likely to affect feeding efficiencies under
 384 farmed conditions.

385

386 **Individual Level Effects**

387

388 **Moulting** A physiological process whereby crustaceans periodically shed their exoskeleton as
 389 a result of their natural growth (Molina-Poveda *et al.* 2002; De Oliveira-Cesar *et al.* 2006;
 390 Rusaini & Owens 2011), moulting affects feeding, growth and reproduction (Robertson *et al.*
 391 1987; Vega-Villasante *et al.* 2000; Molina-Poveda *et al.* 2002). The moulting process in
 392 crustaceans is driven mainly by hormones, but also it is influenced by environmental
 393 conditions, developmental stages and physiological states (Chan *et al.* 1988; Cheng *et al.*
 394 2002). Penaeids have been observed to suppress feeding activity in the preliminary stages
 395 before and during moulting (Dall 1986; Harpaz *et al.* 1987; Chan *et al.* 1988; Dall *et al.* 1990;
 396 Vega-Villasante *et al.* 2000; Moss & Moss 2006; Table 1). This cyclic behaviour of non-
 397 feeding is termed “physiologic fast” and may be caused by non-functionality of some
 398 structures such as the mouth, oesophagus or stomach when an individual sheds the

399 exoskeleton (Vega-Villasante *et al.* 2000). During these periods, penaeids rely on reserves of
400 lipids and carbohydrates to successfully achieve the next moulting phases (Chan *et al.* 1988;
401 Fernández *et al.* 1997). The hepatopancreas is generally considered as the major storage
402 organ in decapod crustaceans, presenting an important role in relation to the accumulation
403 and mobilization of reserves during this time (Allen 1971, Comoglio *et al.* 2004, 2005).
404 Vega-Villasante *et al.* (2000) found that it is possible to calculate the quantity of feed
405 required for shrimp culture according to their moult status and it has been shown in *L.*
406 *vannamei* that the best feed balance and protein efficiency is when shrimp are fed according
407 to their moult cycle (Molina *et al.* 2000).

408

409 There are a range of factors that may influence moulting. Studies in *L. vannamei* have
410 demonstrated the importance of temperature and age. For example, at ranges of 26-30 °C the
411 average moult cycle length was reported as 4-5 days in juveniles aged 1 month, 10-13 days in
412 2 month-old shrimp and around 15-20 days in 6 month-old sub-adults (De Oliveira-Cesar *et*
413 *al.* 2006). However, Chan *et al.* (1988) performed the same study at 20-22° C with adult *L.*
414 *vannamei* and found moulting cycle length was between 28-40 days. In addition to
415 temperature, environmental conditions such as lunar cycle can modulate and coordinate
416 crustacean physiology (Dall *et al.* 1990; Griffith & Wigglesworth 1993). Molina *et al.* (2000)
417 found that moulting in *L. vannamei* occurred mainly during the new moon cycle in around
418 80% of the population. Similar results were observed in the Northern pink shrimp (*Penaeus*
419 *duorarum* Burkenroad), with a high moulting occurrence close to the full moon event (Fuss
420 & Ogren 1966). The presence of weak levels of dissolved oxygen (DO) or extreme water
421 salinity can also reduce moulting frequency in shrimp through changes in shrimp metabolism
422 (Aquacop *et al.* 1988; Allan & Maguire 1991; Staples & Heales 1991). Culture parameters

423 (i.e. temperature, salinity, dissolved oxygen) are therefore very important in modulating
 424 shrimp physiology and behaviour (see section below).

425

426 In terms of behaviour, moulting processes can have substantial effects on overall activity
 427 (such as locomotor or swimming activity), but are particularly important in feeding activity
 428 (Chan *et al.* 1988). The greatest activity levels are generally observed during the inter-moult
 429 phase (Table 1; C). Later, feeding activity is reduced when shrimp are closer to moult in the
 430 Pre-moult phase (D) until an inhibition status is reached, however general activity continues
 431 to be high. During the moulting process, feeding and general activity levels reach the lowest
 432 levels. In the post-moult phase (A-B) shrimp refrain from feeding activities and general
 433 activity is restored in accordance to the exoskeleton hardening. Importantly, depending on
 434 their moult stage stress can affect shrimp differently; pre-moult (D₂) and post-moult (A)
 435 (Table 1) are the phases most sensitive to stress (Robertson *et al.* 1987; Wajsbrot *et al.* 1990),
 436 and this can have an additional influence on their overall behaviour.

437

438 **Table 1.** Moulting cycle of penaeid shrimp (*L. vannamei*), indicating changes in overall and feeding activity
 439 depending of the phase. Data obtained from Chan *et al.* (1988) and Vega-Villasante *et al.* (2000).

440

441

Phase	Proportion of cycle length (%)	Moulting state	Description	Feeding /Overall
A	2%	Post-Moult	New exoskeleton is very soft, with a parchment texture.	None/Weak
B	4%	Post-Moult	New exoskeleton becomes harder.	None/Restored
C ₁	5%	Inter-Moult	New exoskeleton completely formed and resistant.	Restored/Maximal
C ₂₋₃	35%	Inter-Moult	Exoskeleton achieves maximum rigidity.	Maximal/Maximal
D ₀₋₁	43%	Pre-Moult	Detachment of epithelial cuticle. Secretion of new cuticular layers started.	Decreasing/Maximal

D₂₋₃	11%	Pre-Moult	Colouration of the new cuticular layers. Old exoskeleton soft. Opening of the dehiscence groove. Water is absorbed by shrimp.	None/Maximal
E	-	Moult	Tegument expulsion, shrimp leave exoskeleton. Shrimp body expands.	None/None

442

443

444 **Starvation period** It is likely that under high stocking densities, depending on the feeding
445 method, some individuals may have reduced access to feed. Crustaceans, however, are
446 generally able to withstand and recover from prolonged periods of starvation (Stuck *et al.*
447 1996; Hervant *et al.* 1997). Under starvation conditions, *L. vannamei* can reduce their
448 metabolic rate and moulting frequency to save energy (Stuck *et al.* 1996; Comoglio *et al.*
449 2004). Juvenile *L. vannamei* have been noted to survive up to 15 days under starvation,
450 although an increased mortality rate of 35-45% was recorded during days 9 to 15 (Comoglio
451 *et al.* 2004). As seen during moulting, shrimp reserves are mainly limited to lipids stored in
452 the hepatopancreas (Allen 1971; Comoglio *et al.* 2004, 2005). During starvation, *L. vannamei*
453 uses triacylglycerol (ester lipid) for short periods, but use protein during prolonged starvation
454 periods (Cuzon *et al.* 1980; Stuck *et al.* 1996). During a four-week starvation period the
455 kuruma shrimp (*Penaeus japonicus* Spence Bate) utilised carbohydrates first and then lipids
456 to meet their energy requirements. However, proteins were not utilised until after two weeks
457 of starvation (Cuzon *et al.* 1980). Overall body mass may not be affected due to significant
458 increases in body water content during starvation that occurs so as to maintain body volume
459 and internal turgidity (Wilcox & Jeffries 1976; Cuzon *et al.* 1980; Stuck *et al.* 1996).
460 Recovery after very long starvation periods sometimes cannot be achieved if damage to the
461 hepatopancreas and loss of enzyme synthesis has occurred (Comoglio *et al.* 2004). Starvation
462 level can also influence the ability of shrimp to respond to additional stressors such as

463 hypoxia and salinity changes (Hochachka & Lutz 2001; Palacios *et al.* 2004; Zhang *et al.*
464 2006).

465

466 There is limited information on the impact of starvation period on feeding behaviour in
467 relation to aquaculture. Normally when nutritional studies are carried out, specimens are
468 starved for a period of 18-24 hours (e.g. Holland & Borski 1993; Sanchez *et al.* 2005; Nunes
469 *et al.* 2006; Derby *et al.* 2016). It has been shown that feeding activity and behavioural
470 responses in crustaceans can increase across these time periods (Lee & Meyers 1997). In
471 addition, Sanchez *et al.* (2005) discovered that shorter starvation periods of 18 hours did not
472 constitute enough time to observe an effect on feeding rate and response. This means that
473 variation in starvation periods can influence feeding threshold and subsequent behaviours
474 (Costero & Meyers 1993), with relevance in nutritional studies to test novel attractants for
475 shrimp aquaculture.

476

477 In relation to shrimp farming, high starvation periods are commonly linked to the idea of a
478 decrease in shrimp growth and survival (Stuck *et al.* 1996). However, some studies suggest
479 that starvation followed by re-feeding may improve productivity. For example, Wu and Dong
480 (2002), working with the Chinese white shrimp (*Penaeus chinensis* Osbeck) tested different
481 periods of starvation followed by feeding *ad libitum* (e.g. 1 day starvation:4 days *ad libitum*,
482 2:8 and 4:16). While shrimp subjected to the different combinations showed a faster mass
483 gain during the re-feeding period compared to a control group fed *ad libitum* throughout,
484 after 32 days none of the starved and re-fed reached the same body mass as the controls. Lin
485 *et al.* (2008) reported compensatory growth in *L. vannamei* 3 – 9 days after short periods of
486 starvation (1 – 3 days). Moreover, Maciel *et al.* (2018) suggest restricted feed programs in
487 shrimp aquaculture can reduce operating costs (i.e. labour and feeding costs) and improve the

488 pond water quality. These strategies, however, have to be properly controlled as starvation
489 can affect negatively shrimp productivity (i.e. lower growth and survival) and induce
490 negative behaviours such as cannibalism (Martin *et al.* 1998; Kautsky *et al.* 2000; Arnold *et*
491 *al.* 2006; Zhang *et al.* 2010).

492

493 **Sex differences** In penaeid shrimp, females are larger than males (Moss *et al.* 2002; Campos-
494 Ramos *et al.* 2006; Moss & Moss 2006; Gopal *et al.* 2010) due to a greater gain in mass per
495 moult cycle (Hansford & Hewitt 1994). This is most likely caused by different biological
496 requirements at the onset of sexual maturation (Pérez-Rostro & Ibarra 2003). Penaeids
497 possess a stable genetic sex determination system, not determined by environmental
498 conditions as in some crustacean taxa such as copepods and amphipods (Legrand *et al.* 1987).
499 In *L. vannamei*, sexual size dimorphism begins at approximately 10 g (Chow & Sandifer
500 1991) and becomes significant around 17 g (Pérez-Rostro *et al.* 1999; Pérez-Rostro & Ibarra
501 2003). Females are generally heavier than males with a larger cephalothorax and a wider first
502 abdominal segment (Chow & Sandifer 1991; Pérez-Rostro *et al.* 1999, Pérez-Rostro & Ibarra
503 2003), and a study by Accioly *et al.* (2013) determined the possibility of recognising sexual
504 dimorphism in *L. vannamei* using geometric morphometrics in these structures.

505

506 Studies on *P. monodon* (Hansford & Hewitt 1994; Gopal *et al.* 2010) and *L. vannamei* (Moss
507 *et al.* 2002; Moss & Moss 2006) suggest that female monosex cultures grow faster than
508 mixed-sex cultures or male monosex cultures. However, there are very few studies in
509 penaeids that relate variation in behaviour to sexual differences. Moss and Moss (2006)
510 observed that *L. vannamei* males were more active swimmers than females, which normally
511 spent most time on the bottom of tanks. Moreover, males presented more aggressive
512 behaviours than females when competing for food. Males tended to monopolise the food

513 source for more time than females, even though they were smaller. It is therefore clear that
514 gender is a more important factor than size for explaining variations in feeding activity in this
515 species. It appears that *L. vannamei* females do not have a competitive advantage over food,
516 one of the possible causes of sexual growth dimorphism in this species is physiological
517 advantage (Moss & Moss 2006). For example, Hansford and Hewitt (1994) reported that *P.*
518 *monodon* females had a lower feed conversion ratio and a more efficient digestibility of food
519 energy compared to males. At the same time, males could have a greater metabolic rate
520 and/or stress levels than females (Taylor *et al.* 2004; Moss & Moss 2006). This could explain
521 the higher aggressiveness and feeding activity observed in males, behaviour that has also
522 been recorded in other crustaceans such as rock shrimp (*Rhynchocinetes typus* H. Milne
523 Edwards) (Dennenmoser & Thiel 2007) and American lobster (*H. americanus*) (Karavanich
524 & Atema 1998) as well as other finfish aquaculture species such as Atlantic salmon (*Salmo*
525 *salar* Linnaeus) (Cutts *et al.* 1998) and rainbow trout (*Oncorhynchus mykiss* Walbaum)
526 (Metcalf 1986). However, the paucity of information makes it difficult to make definitive
527 conclusions here and more research is required.

528 **Personality** Behaviour is perhaps the most flexible phenotypic feature, offering the quickest
529 response of animals to unexpected events in a variable environment (Hazlett 1995). This
530 phenotypic plasticity allows appropriate responses to changing conditions or situations (Sih *et*
531 *al.* 2004; Dingemanse *et al.* 2010), having major impacts on individual fitness (Sih *et al.*
532 2004; Gherardi *et al.* 2012; Briffa *et al.* 2015). However, it is also evident that phenotypic
533 plasticity might present some constraints with regards to sensory capabilities, morphological
534 limitations or limits on learning abilities (Hazlett 1995; DeWitt *et al.* 1998; Briffa *et al.* 2015;
535 Briffa & Sneddon 2016). Such limits on behavioural plasticity appear to vary between and
536 within populations (Wilson 1998), leading in some cases to variation in behaviour between
537 individuals (Briffa *et al.* 2008; Briffa *et al.* 2015). Such variations in behaviours may be

538 considered as ‘personality’ traits when they are observed to be consistent and repeatable in
539 individuals across time in different contexts (e.g. feeding, mating) and situations (e.g.
540 presence of food or predators) (Briffa & Weiss 2010; Briffa & Twyman 2011; Gherardi *et al.*
541 2012). In many studies, individuals are distinguished based on the ‘shyness-boldness axis’
542 defined by Wilson *et al.* (1993). Combinations of behaviours that occur across different
543 situations and contexts with behavioural consistency within individuals of a population may
544 also be observed and are described as ‘behavioural syndromes’ (Gherardi *et al.* 2012; Briffa
545 *et al.* 2015).

546 Reliable, repeatable individual variation in behaviour, or animal personality, has been
547 reported in a wide group of taxa including mammals, birds and fish. Recent studies have also
548 uncovered repeatable behavioural differences in invertebrates such as arachnids, cnidarians
549 and crustaceans (see Reviews of Dingemanse *et al.* 2010 and Gherardi *et al.* 2012). In
550 crustaceans, evidence for bold and shy individuals has been found in the hermit crab *P.*
551 *bernhardus* where bold individuals showed high levels of exploratory behaviour in a new
552 environment, investigated novel objects and when disturbed showed a sudden startle response
553 of short duration (Briffa *et al.* 2008). Similarly, Vainikka *et al.* (2011) observed personality
554 traits in crayfish (*Astacus astacus* Linnaeus) and proposed that boldness in this species
555 influences predatory risk and intraspecific competition.

556 The occurrence of personality traits in crustacean aquaculture, particularly in penaeid shrimp,
557 may be important in the context of feeding where bolder individuals might be more likely to
558 interact with feed and consume it at a faster rate. Currently there is no evidence to support
559 this within shrimp aquaculture, however, in finfish aquaculture some studies highlight the
560 presence of personalities and coping styles (stress responses) which can provide advantages
561 for culture, for example in establishing selection-based breeding programmes or improving

562 domestication (Castanheira *et al.* 2013; Ibarra-Zatarain *et al.* 2016). It may also be possible to
563 link proactive (bold) individuals with important aquaculture outputs, such as growth, disease
564 resistance, food conversion, or survival (Ibarra-Zatarain *et al.* 2016). Therefore,
565 understanding the role of animal personality in commercial settings may be of great
566 importance in optimising production (Huntingford & Adams 2005).

567 **Environmental Effects**

568

569 ***Environmental enrichment*** As benthic animals, shrimp are mainly constrained to two-
570 dimensional space rather than three-dimensional volume (Kumlu *et al.* 2001), although when
571 shrimp are stocked at high densities the water column becomes more important as they can
572 feed on suspended food particles (Tacon *et al.* 2013). Therefore, bottom substrate becomes an
573 important concern in an aquaculture setting. Several studies have demonstrated that artificial
574 substrates could increase the food supplement for shrimp (i.e. enhance the colonisation of
575 epiphytic biota) (Thompson *et al.* 2002; Burford *et al.* 2004; Moss & Moss 2004; Arnold *et*
576 *al.* 2005, 2006) as well as improve the water quality and control pathogenic bacteria
577 (Bratvold & Browdy 2001; Thompson *et al.* 2002; Moss & Moss 2004; Ballester *et al.* 2007;
578 Kumar *et al.* 2017). In *L. vannamei* the addition of artificial substrates has improved growth
579 and/or survival of postlarvae and juveniles, even at high stocking densities (Bratvold &
580 Browdy 2001; Moss & Moss 2004; Zarain-Herzberg *et al.* 2006). For example, Bratvold &
581 Browdy (2001) indicated that, in tanks containing AquaMats™, *L. vannamei* individuals
582 were up to 30% heavier than in tanks without mats.

583

584 Artificial substrates may confer several potential behavioural benefits for shrimp culture by
585 providing refuge for moulting shrimp, reducing aggressive interactions (e.g., cannibalism)
586 and increasing the surface area on which shrimp graze (Moss & Moss 2004; Arnold *et al.*

2006). It has also been suggested that for species that exhibit burrowing behaviour, sediment may be required for optimal production (Allan & Maguire 1995). Some studies have shown little burrowing activity exhibited by *L. vannamei* (Boddeke 1983; Robertson *et al.* 1993; Pontes & Arruda 2005a,b; Zhang *et al.* 2010; Da Costa *et al.* 2016;), thus the presence of sediment may have relatively little effect on its growth and behaviour (Bratvold & Browdy 2001). However, information is scarce and more research on this topic is needed, as according to Moctezuma and Blake (1981) *L. vannamei* shows a clear burrowing activity during diurnal hours and emerge at night, as do other *Penaeus* spp. (Hindley 1975; Boddeke 1983; Nunes *et al.* 1996). Moreover, Zhang *et al.* (2010) found behavioural changes in *L. vannamei* in relation to artificial substrates, demonstrating a clear change in shrimp distribution between tanks with and without artificial substrate. The proportion of shrimp on the bottom of tanks with artificial substrates was significantly lower than tanks without artificial substrates, with the differences becoming more distinct with a longer rearing time. This provides evidence of how artificial substrates can disperse shrimp from tank bottoms and thus alleviate the negative effect of high stocking density on shrimp production. The use of artificial substrate may also benefit feeding efficiency, stimulating more natural behavioural processes. Moreover, the role of artificial substrates in relation to the production of natural food supplement could encourage *L. vannamei* feeding activity.

605

Conspecific presence The presence of conspecifics and stocking density is an important consideration in aquaculture production. Normally, when held in higher densities, shrimp exhibit a reduction in growth and survival as a result of a combination of different factors including a decrease in favourable space and natural food sources and an increase in adverse behaviour such as cannibalism (Martin *et al.* 1998; Kautsky *et al.* 2000; Arnold *et al.* 2006; Zhang *et al.* 2010). It is also obvious that feeding success is density-dependent (Chavanich *et*

612 *al.* 2016) therefore stocking density is an important aspect when considering feeding
613 behaviour patterns (Da Costa *et al.* 2016). In general, *L. vannamei* is amenable to culture at
614 very high densities from 60 to 150 individuals/m² in pond culture, and even as high as 400
615 individuals/m² in controlled recirculated tank conditions (Briggs *et al.* 2004; Funge-Smith &
616 Briggs 2005). Such intensive culture systems, however, require a much higher degree of
617 control over environmental parameters (Briggs *et al.* 2004; Funge-Smith & Briggs 2005).
618 Moreover, high densities can compromise feeding management of shrimp production, as high
619 densities have more competition for access to feeding trays and that can restrict the number
620 of shrimp feeding (Sanchez *et al.* 2005). Therefore, feed management strategies need to be
621 implemented and automatic feed dispensers may present a promising solution to restricted
622 feeding at high densities by means of dispensing smaller amounts of feed many times a day
623 (Jory 2016). Stocking density affects growth of *L. vannamei* (Balakrishnan *et al.* 2011), and
624 stronger density-growth relationships have been observed at medium and higher densities (40
625 – 150 individuals/m²) than at moderately low densities (5 - 40 individuals/m²) (Sandifer *et al.*
626 1987, 1988; Wyban *et al.* 1987, 1988). According to Wyban *et al.* (1988), *L. vannamei*
627 shrimp stocked at a density of 45 individuals/m² presented an excellent growth and survival
628 rate, however at higher densities *L. vannamei* can often show an aggressive feeding
629 behaviour when food sources are limited (Zhang *et al.* 2010). Nevertheless, once again, this
630 may be mitigated through appropriate feed management.

631

632 There has been limited work conducted on the effects that different stocking densities have
633 on behaviours of *L. vannamei*. Da Costa *et al.* (2016) found that groups at low (50
634 individuals/m²), medium (75 individuals/m²) and high density (100 individuals/m²) displayed
635 differences in their individual behaviours. With regards to feeding, medium densities had the
636 lowest feeding frequency of all groups. Higher densities were observed to have a medium

637 feeding efficiency and lower densities performed best in terms of feeding (Da Costa *et al.*
638 2016). With regards to other behaviours, the study noted that there were, however, few
639 swimming behaviours at lower densities compared to higher stocking densities, most likely as
640 a trade-off due to increased feeding. As high frequency of swimming behaviour in *L.*
641 *vannamei* has been shown to be associated with stress (Taylor *et al.* 2004), it is also possible
642 that animals at a lower stocking density moved less frequently due to less chemical, tactile or
643 visual interference by other individuals.

644

645 **Photoperiod** In crustacean aquaculture, light conditions and photoperiod play important roles
646 in both behaviour (e.g. Pontes 2006; Santos *et al.* 2016) and physiology (e.g. Nakamura &
647 Echavarria 1989; Pontes & Arruda 2005b; Ravi & Manisseri 2013). Light conditions are
648 probably the most important single factor for determining the locomotor activity in penaeid
649 shrimp (Dall *et al.* 1990; Wassenberg & Hill 1994), therefore understanding the behaviours
650 which result in the activity pattern of species such as *L. vannamei* during the day/night is vital
651 for determining feeding schedules in aquaculture (Pontes 2006).

652

653 In general, penaeids exhibit circadian rhythms which are linked with diurnal burrowing habits
654 and emergence from the substratum in the dark (Hindley 1975; Boddeke 1983; Nunes *et al.*
655 1996). For example, *P. japonicus* is naturally active at night therefore captive animals are
656 usually fed after sunset (Nakamura & Echavarria 1989). However, *L. vannamei* belong to the
657 group of non-burrowing shrimp, and are likely to exhibit the same feeding and locomotor
658 activity during the day as well as at night, and do not usually burrow during the daytime
659 (Boddeke 1983; Robertson *et al.* 1993; Pontes & Arruda 2005a,b; Zhang *et al.* 2010; Da
660 Costa *et al.* 2016). However, some studies suggest feeding behaviour in *L. vannamei* is
661 heightened under light conditions. This may be a result of temperature fluctuations, with low

662 night-time water temperatures relative to the daytime temperatures reducing the activity of
663 the shrimp (Robertson *et al.* 1993). Additionally, *L. vannamei* seems to be less motile under
664 strong light than weak light conditions (Zhang *et al.* 2006). Sanudin *et al.* (2014) examined *L.*
665 *vannamei* larvae (0.5cm total length (TL)) and demonstrated a greater ingestion rate of feed
666 in light conditions compared to dark. As the shrimp grew (up to 1 – 1.5 cm TL), differences
667 between light and dark conditions did not influence the feeding activity to the same degree.
668 These findings may reflect the ontogenetic development of chemoreceptors to find food
669 sources compared to larvae stages that rely more on vision (Sanudin *et al.* 2014). However,
670 more research into ontological changes in chemoreception vs vision is necessary and presents
671 an exciting area for future studies.

672

673 The effect of light on feeding and growth efficiency may be dependent on whether trials are
674 located indoors or outdoors. For example, *L. vannamei* juveniles had better feeding and
675 growth efficiency during the light hours in trials performed inside (Robertson *et al.* 1993;
676 Tacon *et al.* 2002; Pontes & Arruda 2005b; Pontes *et al.* 2006). However outdoor trials found
677 no significant differences between diurnal and nocturnal food consumption patterns (Nunes *et*
678 *al.* 1996; Tacon *et al.* 2002). Regardless of these findings, *L. vannamei* shows the highest
679 ingestion of feed around the middle of the light phase (Lima *et al.* 2009) and it has been
680 determined that feed should be offered in the light phase and at a greater proportion 7 hours
681 after sunrise (Pontes *et al.* 2006). Substrate exploration behaviour in *L. vannamei*, indicative
682 of the search for feed, was more intense around this time (Pontes 2006). Moreover, some
683 studies have demonstrated the possibility of synchronising feeding activity with shrimp
684 circadian rhythms (Miguel & Aréchiga 1994; Santos *et al.* 2016); understanding such
685 behavioural rhythms is critical for feeding management for species such as *L. vannamei*. For
686 example, Santos *et al.* (2016) showed that the use of automatic self-feeding systems by *L.*

687 *vannamei* individuals was influenced by their circadian rhythms (i.e. shrimp activated the
688 feeders more at night). This information may have potential implications for commercial
689 systems that distribute feed automatically to shrimp (sometimes over 24 hours), a method
690 increasing in popularity in shrimp farms (Jory 2016; Molina & Espinoza 2018).

691

692

693 **Water Quality Effects**

694

695 **Salinity** Salinity is one of the most basic environmental parameters of marine shrimp culture
696 (Zhang *et al.* 2006). In general, euryhaline decapod crustaceans acclimated to sea water are
697 osmoconformers although some groups, such as the penaeid shrimp, have the capacity to
698 osmoregulate (Dall 1981; Dall *et al.* 1990). Osmotic regulation in penaeid shrimp is a
699 physiological property which can determine their distribution under different salinities
700 (Charmantier 1987; Vargas-Albores & Ochoa 1992). In *L. vannamei* the osmoregulatory
701 ability is better in juvenile phases, mostly at low salinities, declining naturally when they
702 reach subadult or adult stages (Gong *et al.* 2004; Bett & Vinatea 2009). This is reflected in
703 their natural distribution where postlarvae and juveniles inhabit brackish water but migrate as
704 adults to marine habitats (Vargas-Albores & Ochoa 1992; Ponce-Palafox *et al.* 1997). In
705 addition, size differences within a species may also affect the osmoregulatory capacity, with
706 smaller individuals being better regulators than those that are larger (Vargas-Albores &
707 Ochoa 1992).

708

709 The isosmotic point has been associated with optimum conditions for growth in penaeid
710 shrimp (between 20 – 25 ppt), however *L. vannamei* grows particularly well at low salinities
711 of around 10 – 20 ppt, below isosmotic conditions (Wyban & Sweeny 1991; Bray *et al.* 1994;

712 Rosas *et al.* 2001; Bett & Vinatea 2009; Ponce-Palafox *et al.* 2013). However, *Litopenaeus*
713 *vannamei* tolerates a wide range of salinities from 0.5 -50 ppt (Charmantier 1998; Briggs *et*
714 *al.* 2004; Funge-Smith & Briggs 2005; Bett & Vinatea 2009), levels that are currently found
715 within shrimp farming. Moreover, this ability makes the species a good candidate for inland
716 farms where they may be subject to high salinity fluctuations over a 24 hour period (Anger
717 1996; Díaz *et al.* 2001; Briggs *et al.* 2004; Funge-Smith & Briggs 2005).

718

719 In terms of how behaviour may be affected by salinity, there are a number of studies which
720 can provide better understanding. Gleeson *et al.* (1996) found that under low salinity
721 concentrations (< 12.5 ppt) or fresh water conditions (0 ppt) the functioning of the
722 aesthetasc/olfactory sensilla in crustaceans can be compromised, resulting in a decrease in
723 detection capacity. Rosas *et al.* (2001) found an increase in locomotor activity when *L.*
724 *vannamei* juveniles were maintained under 10 ppt. This may indicate an escape response to
725 those environmental conditions, as *L. vannamei* individuals reared under low salinity
726 conditions present low stress tolerance (Li *et al.* 2007). Davis *et al.* (2002) found similar
727 effects of low salinity where *L. vannamei* individuals were mainly found at the sides of the
728 tanks, displaying lethargic behaviours and a greater sensitivity to stress caused by handling,
729 temperature changes and low dissolved oxygen. There is little information on how feeding
730 behaviour is affected across different salinities, however, studies note a clear effect of salinity
731 on oxygen consumption for *L. vannamei*, which is related to feeding activity (i.e. higher
732 oxygen consumption/respiratory metabolism increases feeding activity). Nevertheless, this
733 relationship remains unclear as some studies observed a direct increase between oxygen
734 consumption and salinity concentration (e.g. Bett & Vinatea 2009; Zhang *et al.* 2009; Ponce-
735 Palafox *et al.* 2013) and others an inverse relationship between them (e.g. Rosas *et al.* 2001;

736 Li *et al.* 2007; Li *et al.* 2017). It is clear that further research is necessary to fully understand
737 the effect of salinity on shrimp behaviour.

738

739 **Temperature** Another important environmental factor influencing feeding and overall
740 metabolic rate in marine invertebrates is temperature (Kinne 1966; Armitage & Wall 1982;
741 Wyban *et al.* 1995; Magallón-Barajas *et al.* 2006), particularly in organisms with life cycles
742 involving estuarine areas that experience large temperature oscillations (Darsey 1990). In *L.*
743 *vannamei*, temperature and temperature-salinity interactions are the main factors modulating
744 growth, feeding rate and oxygen consumption (Wyban *et al.* 1988, 1995; Ponce-Palafox *et al.*
745 1997; Díaz *et al.* 2001; Ponce-Palafox *et al.* 2013). Several studies have concluded that
746 oxygen consumption between 24-30° C (at salinities of 15-25 ppt) results in greater stability
747 for *L. vannamei* cultures (Villarreal *et al.* 1994; Bett & Vinatea 2009; Valenzuela-Quiñonez
748 *et al.* 2011; Ponce-Palafox *et al.* 2013). Moreover, this temperature range overlays with the
749 optimal growth rate of the species (Wyban *et al.* 1995; Briggs *et al.* 2004). These
750 temperatures are normally associated with the native distribution of *L. vannamei* on the
751 Pacific coast. However, *L. vannamei* is able to tolerate a wide range of temperatures, from as
752 low as 15° C up to 33° C, but at reduced growth rates (Wyban & Sweeny 1991; Funge-Smith
753 & Briggs 2005). This adaptation is advantageous because juvenile *L. vannamei* often inhabit
754 semi-enclosed or closed estuaries and rock pools that are subject to high temperature
755 fluctuations over a 24 hour period (Anger, 1996). Tolerance of lower temperatures allows
756 culture during the cold season (October-February) in the Northern hemisphere (Briggs *et al.*
757 2004; Funge-Smith & Briggs 2005).

758

759 In the context of feeding behaviour, temperature has a considerable effect across all size
760 classes. Feeding rates and subsequent growth of *L. vannamei* are directly correlated with

761 temperature and fluctuate inversely with size (Wyban *et al.* 1995), with shrimp cultured at
762 higher temperatures growing faster as their metabolic rates are accelerated (Gong *et al.*
763 2004). According to Ponce-Palafox *et al.* (1997), juveniles of *L. vannamei* exhibited low food
764 consumption at 20° C compared with hyperactive animals at 35° C (salinities > 20 ppt).
765 Moreover, when they were offered unlimited food, shrimp maintained at 35° C had the
766 highest rate of food consumption. Wyban *et al.* (1995) found the same temperature effects in
767 relation to feeding activity and growth, particularly between 23 – 27° C, with shrimp being
768 extremely sensitive to small temperature changes. The study also concluded that *L. vannamei*
769 culture at low temperature conditions (23° C) resulted in problems in growth associated with
770 feeding activity which could be solved through use of more attractive diets. Hernández *et al.*
771 (2006) observed that the preferred feeding temperature of *L. vannamei* was within 26 – 31° C,
772 and according to Cuéllar-Anjel *et al.* (2010) feeding should be conducted at temperatures
773 above 26° C.

774

775 **pH** The pH of body fluids governs the ionization state of proteins and their physiological
776 function, and its control is vital for the proper functioning of animals (Reeves & Rahn 1979).
777 Studies have demonstrated the adverse effect of changes in pH on the chemo-responsiveness
778 of crustaceans (e.g. Allison *et al.* 1992; De la Haye *et al.* 2011, 2012), which have important
779 implications for feeding behaviour. Moreover, abnormal pH values have been reported to
780 cause a wide range of problems across a variety of crustacean species such as stunted growth
781 and reduced survival (Allan & Maguire 1992; Wang *et al.* 2002; Chen & Chen 2003; Wang
782 *et al.* 2009), disturbed ionoregulation (Morgan & McMahon 1982; Allan & Maguire 1992),
783 acid-base imbalance (Morgan & McMahon 1982), DNA damage (Wang *et al.* 2009) and
784 abnormal behaviours (Allison *et al.* 1992; Das & Sahu 2005). Therefore, controlling pH is
785 important due to the large potential fluctuations that may occur in aquaculture ponds.

786 The majority of studies on *L. vannamei* utilise pH parameters in the range of 7.4 – 8.2
787 (Samocha *et al.* 2004b; Zhang *et al.* 2006; Grey *et al.* 2009; Wang *et al.* 2009; Da Costa *et al.*
788 2016), and it has been found that changes in pH have important implications for survival. For
789 example, Wang *et al.* (2009), found 100% survival across a 24 hour period for *L. vannamei*
790 under conditions at pH 7.4. Survival was reduced however to 65% under more acidic
791 conditions (pH 5.6) and 35% under more alkaline conditions (pH 9.3). With regards to
792 feeding behaviours, a number of effects due to changes in pH have been identified in
793 crustaceans. Initially, low pH may reduce chemoreceptive perception by modifying the
794 charge distribution on the chemoreceptor cells of crustacean sensory organs (Tierney &
795 Atema, 1988). Moreover, changes in chemo-responsiveness might simply reflect reduced
796 activity levels, or reduced motivation to respond to chemical cues, occurring as a result of the
797 elevated metabolic load of maintaining acid-base balance under conditions of low pH (Spicer
798 *et al.* 2007). There may also be direct physical damage to the sensory organs as a result of
799 low pH conditions (De la Haye *et al.* 2012). Additionally, it has been shown that crustaceans
800 in low pH conditions significantly reduce antennular flicking which triggers less locomotor
801 activity, lack of stimulation and metabolic depression (Allison *et al.* 1992; De la Haye *et al.*
802 2011, 2012). For example, reduced pH compromises resource assessment and decision-
803 making behaviour in European hermit crabs (*P. bernhardus*) (De la Haye *et al.* 2011). Allison
804 *et al.* (1992) found similar results in freshwater crayfish (*C. bartoni*), which flicked its
805 antennules at a reduced rate and failed to locate a food source under low pH conditions (4.0)
806 instead of normal pH conditions (7.5). Therefore, the decreased flicking observed under low
807 pH conditions could imply a reduced ability to detect the chemical stimulus (De la Haye *et al.*
808 2011).

809

810 **Dissolved oxygen** Dissolved oxygen (DO) is a major limiting factor in crustacean aquaculture
811 (Martinez-Palacios *et al.* 1996; Cheng *et al.* 2003; Pérez-Rostro *et al.* 2004; Zhang *et al.*
812 2006). During the initial phases of shrimp development (larvae and juvenile forms), it is vital
813 to maintain adequate oxygen levels (Bett & Vinatea 2009). The bottom layer of pond waters,
814 where shrimp mostly remain, may become hypoxic or even anoxic due to decomposition of
815 accumulated organic matter and organism respiration (Cheng *et al.* 2003; Zhang *et al.* 2006).
816 Normally, DO values higher than 5 mg l⁻¹ have been suggested for intensive culture practices
817 (Cheng *et al.* 2003). Low DO concentrations negatively affect the behaviour and physiology
818 of crustaceans, such as growth, survival, respiration and circulation, metabolism, moulting
819 and feeding (Seidman & Lawrence 1985; Clark 1986; Aquacop *et al.* 1988; Allan & Maguire
820 1991).

821

822 The effect of hypoxia on growth, survival, feeding, moulting, behaviour, osmoregulatory
823 capacity and immune response of *L. vannamei* has been documented by some authors
824 (Aquacop *et al.* 1988; Hopkins *et al.* 1991; Charmantier *et al.* 1994; Pérez-Rostro *et al.* 2004;
825 Zhang *et al.* 2006). *Litopenaeus vannamei* tolerates low DO levels, although it does not grow
826 well below 2.8 - 3 mg l⁻¹, which is considered the limit of hypoxic conditions (Martinez-
827 Palacios *et al.* 1996; Vinatea *et al.* 2009). Lethal DO levels for *L. vannamei* are between 0.2 –
828 1 mg l⁻¹ (Hopkins *et al.* 1991; Pérez-Rostro *et al.* 2004) and it appears *L. vannamei* is an
829 oxygen conformer with modification to respiratory rate detected when DO levels decreased
830 below 5 mg l⁻¹ (Martinez-Palacios *et al.* 1996). It is possible to control some characteristics
831 of *L. vannamei* cultures to improve their resistance against hypoxic conditions. Zhang *et al.*
832 (2006) concluded that certain water parameters (22° C, 16.6 ppt salinity, pH 7.6) and strong
833 light conditions can help to counteract the effects of low DO levels. Nevertheless, exposure

834 time at low DO or high culture densities has a cumulative effect on the mortality rate or on
835 subsequent performance after recovery (Pérez-Rostro *et al.* 2004; Vinatea *et al.* 2009).

836

837 Under hypoxia, individuals respond by decreasing energy production as well as changes to
838 behaviour (Pérez-Rostro *et al.* 2004). Changes in general locomotor behaviour in *L.*
839 *vannamei* have been observed during hypoxic periods. When DO is at 50% saturation, *L.*
840 *vannamei* exhibits an increase in activity, making short but frequent random swimming
841 movements. Once the saturation value is below 50%, activity decreases, with slower
842 swimming speeds but with a clear and evident pattern of surface-seeking behaviour (Zhang *et*
843 *al.* 2006). Finally, activity ceases when DO levels fall below 1 mg l⁻¹ (lethal concentration)
844 (Allan & Maguire 1991; Pörtner *et al.* 2004; Zhang *et al.* 2006). This reduction in activity
845 may be a result of increased lactate concentrations in muscle and hepatopancreas after
846 exposure to hypoxic conditions (Mauro & Malecha 1984; Pérez-Rostro *et al.* 2004).
847 Additionally, shrimp exhibit increased ventilatory activity in low oxygen conditions
848 (Martinez-Palacios *et al.* 1996), which enhances CO₂ excretion from the haemolymph,
849 resulting in increased blood pH (Hagerman & Uglow 1984, 1985; Mauro & Malecha 1984).

850

851 There is little information on feeding behaviour and how it is affected by fluctuations of DO.
852 Cuéllar-Anjel *et al.* (2010) concluded that when DO concentrations fall below 4.5 mg/L
853 reduced feed consumption in *L. vannamei* occurs. In addition, if DO concentration is low for
854 a period of time (days or weeks), daily feed rations should be recalculated with the aim of
855 reducing or suspending feeding until normal water DO levels are attained.

856

857 **Nitrogenous compounds** Ammonia is the main nitrogenous end-product excreted by
858 crustaceans (Dall *et al.* 1990), and decomposing organic solids (e.g. uneaten food) can

859 produce an additional source of ammonia inside aquaculture systems (Chen & Lei 1990).
860 Therefore, nitrogenous compound concentration is directly correlated with stocking density
861 (Martin *et al.* 1998). Ammonia is utilised as a substrate by autotrophic nitrifying aerobic
862 bacteria, *Nitrosomas* and *Nitrobacter*, which oxidize it to nitrite and nitrate, respectively
863 (Chen & Kou 1992, Montoya *et al.* 2002). According to Martin *et al.* (1998), up to 38% of
864 the nitrogen entering ponds via the inflow and in feed pellets could accumulate in the
865 sediment, although it depends on the system used (e.g. biofloc systems can maintain
866 nitrogenous compounds at safe levels) (Vinatea *et al.* 2009; Maicá *et al.* 2014). *Litopenaeus*
867 *vannamei* presents a higher tolerance to total ammonia nitrogen (TAN) levels than other
868 penaeid shrimp species (Frías-Espericueta *et al.* 1999; Lin & Chen 2001; Schuler *et al.* 2010),
869 presenting a 96 hours LC50 with concentrations as high as 39.54-70.9 mg/l at high salinity
870 conditions (34-35 ppt) (Frías-Espericueta *et al.* 1999; Lin & Chen 2001). In addition, *L.*
871 *vannamei* also shows an increased tolerance to ammonia with age (Frías-Espericueta *et al.*
872 2000). However, this tolerance to ammonia can be compromised depending on culture
873 parameters. For example, it has been observed that TAN tolerance decreased to 24.39 mg/l
874 when salinity dropped to 15 ppt (Lin & Chen 2001) or to 9.33 mg/l at a salinity of 3 ppt (Li *et*
875 *al.* 2007). Low salinities may therefore increase ammonia excretion in penaeid shrimp (Rosas
876 *et al.* 1999), resulting in higher susceptibility of *L. vannamei* to ammonia toxicity with
877 salinity levels below 5 ppt (Li *et al.* 2007). Similarly, at high pH levels (pH 9) ammonia may
878 become very toxic for *L. vannamei* (Magallón-Barajas *et al.* 2006).

879

880 The accumulation of nitrogenous compounds inside culture systems can produce a series of
881 non-beneficial effects, such as immune suppression and disease incidence (Liu & Chen
882 2004), physical damage (e.g. hepatopancreas, gills) (Kuhn *et al.* 2010; Furtado *et al.* 2015),
883 cessation of feeding and growth (Frías-Espericueta *et al.* 2000), which result in higher

884 mortality rates in penaeid shrimp (Chen & Lei 1990). These setbacks mean the accumulation
885 of nitrogenous compounds is one of the most limiting parameters within shrimp aquaculture.
886 For that reason, safe levels of TAN, nitrites and nitrates for rearing *L. vannamei* were
887 estimated to be 2.44, 6.1 and 145 mg/l, respectively, at low salinities and 3.95, 25.7 and 177
888 mg/l at high salinities (Lin & Chen 2001, 2003; Kuhn *et al.* 2010; Furtado *et al.* 2015). When
889 ammonia levels are high, the first reaction of penaeids is the reduction or cessation of feeding
890 which in turn reduces the production of metabolic ammonia (Colt & Armstrong 1981).
891 Moreover, according to ammonia concentration increases in the water, the ammonia excreted
892 by shrimp is reduced. Consequently, the ammonia level in blood and tissues increases with
893 deleterious effects on the shrimp physiology and metabolism (Frías-Espéricueta *et al.* 2000;
894 Barbieri 2010). This is directly connected with how ammonia affects growth negatively in
895 penaeid shrimp. For example, Wickins (1976) reported ammonia concentrations as small as
896 0.22-0.69 mg/l can reduce growth in several penaeid shrimp species by 50%. Chen and Kou
897 (1992) found in *P. japonicus* juveniles that 40 days of 30 mg/l of ammonia produced a 86%
898 reduction in mass gain and a 17% reduction in length compared to controls. Furthermore,
899 there appears to be an important correlation between higher TAN levels and lower feeding
900 activity, although more research into effects of nitrogenous compounds on specific feeding
901 behaviours is required.

902

903

904 **HOW CAN WE USE BEHAVIOUR TO IMPROVE SHRIMP AQUACULTURE?**

905

906 Ethological studies in the context of shrimp farming are limited (e.g. Primavera & Leбата
907 1995; Soares *et al.* 2005; Silva *et al.* 2012; Da Costa *et al.* 2016), with those related to
908 feeding behaviour generally investigating responses to chemical stimulants (e.g. Costero &

909 Meyers 1993; Pittet *et al.* 1996; Nunes *et al.* 2006). The importance of chemical cues in
910 identifying and orientating towards potential food sources means that commercial diets must
911 be chemically attractive and include specific chemical cues that can be rapidly recognized as
912 a food source and initiate shrimp feeding behaviour (Zimmer-Faust 1989; Sanchez *et al.*
913 2005; Grey *et al.* 2009). Feeding effectors (attractants and palatability factors) are compounds
914 that attract animals via chemical stimuli and enhance palatability of food (Lee & Meyers
915 1997; Smith *et al.* 2005; Suresh *et al.* 2011). As such, compounds that can enhance feeding
916 behaviour have received a lot of attention as a potential refinement in crustacean aquaculture
917 (e.g. Hartati & Briggs 1993; Smith *et al.* 2005; Nunes *et al.* 2006; Grey *et al.* 2009),
918 especially for species such as *L. vannamei* (Heinen 1980; Costero & Meyers 1993; Gadiant
919 & Schai, 1994; Peñafiorida & Virtanen 1996; Nunes *et al.* 2006). Table 2 summarises feeding
920 studies in shrimp which have examined the use of potential feed effectors. Feeding attractants
921 can elicit behavioural and physiological responses from the shrimp's olfactory and gustatory
922 systems, stimulating the different feeding behaviour phases discussed above (Lee & Meyers
923 1996) (see behavioural responses section). For example, a range of food effectors tested by
924 Nunes *et al.* (2006) in *L. vannamei* resulted in improved feeding efficiency correlated with
925 reduced time spent on perception, orientation and locomotion, and more time spent feeding.

926

927 In addition to the chemical utilised, it is important to consider the amount which is used in
928 feed (i.e. the inclusion level) when understanding effects on behaviour. Obviously, the level
929 of feed enhancer needs to be high enough to stimulate shrimp behaviour (Nunes *et al.* 2006).
930 Derby *et al.* (2016) found a positive relationship between the inclusion level of krill meal
931 within feed and the increase in attractability and feeding consumption of *L. vannamei* to that
932 feed in the first 60 minutes after the food was presented. However, detection of the chemical
933 does not necessarily imply that the diet will be acceptable or consumed and assimilated

934 efficiently; indeed the attractiveness of the diet may become attenuated with time (Lee &
935 Meyers 1997). It is also important that the level of attractant is not too high as desensitization
936 can occur (Pittet *et al.* 1996). Moreover, water quality can affect the crustacean's ability to
937 detect and respond to chemicals (Lee & Meyers 1996; Koehl 2006, 2011; De la Haye *et al.*
938 2012).

939

940 Even if an individual shrimp is attracted to a food source, there are numerous external and
941 internal influences which may alter its behaviour (see Fig. 1). Another area of behavioural
942 research that has been used to the benefit of aquaculture practice has focused on optimal
943 feeding strategies and how feed is presented. Although recently some countries have adopted
944 more complex feeding protocols (e.g. automatic feeders, acoustic feeding-control) (Bador *et*
945 *al.* 2013; Jory 2016), most feed management strategies have remained largely unaltered since
946 the mid-1980s (e.g. manual broadcasting) (Carvalho & Nunes 2006; Jory 2016). Feed is
947 commonly offered between two to four times per day with the ration based on shrimp
948 biomass and the eaten-uneaten feed ratio (Jory 1995; Pontes *et al.* 2008; Lima *et al.* 2009).
949 Feed is generally supplied to the shrimp in feeding trays which are a good tool for the
950 estimation of how much shrimp are eating daily and allow spatial feed distribution
951 throughout ponds (Bador 2000; Cuéllar-Anjel *et al.* 2010). Frequency of feeding may change
952 seasonally (Seiffert & Andreatta 2004) and distribution of the daily ration in different
953 percentages is common practice (Jory 1995). However, understanding how individuals
954 respond to different feed presentations will be important in maximising efficiencies.

955

956 Feed has to be consumed by shrimp in the first two hours after feed has been introduced into
957 the water, before changes in the physical and chemical characteristics of the pellets occur,
958 leading to a reduction in feed attractant properties and nutrients which leach into the water

959 (Sick *et al.* 1973, Smith *et al.* 2002; Carvalho & Nunes 2006; Cuéllar-Anjel *et al.* 2010).
960 Water absorption also makes the pellets more difficult for shrimp to manipulate into their
961 buccal cavity (Sick *et al.* 1973). While *P. monodon* has been observed to consume pellets
962 from feeding trays which had been in the water for up to 4 hours (Smith *et al.* 2002), uneaten
963 food crumbles, and is consequently not consumed (Martinez-Cordova *et al.* 1998), leading to
964 deterioration in water quality (Nunes *et al.* 1996; Sanchez *et al.* 2005). Ensuring that food is
965 provided at the most appropriate times and rations is, therefore, important. In some studies,
966 an increased feeding frequency has had a positive effect on shrimp growth (Robertson *et al.*
967 1993; Wyban *et al.* 1995; Tacon *et al.* 2002) whereas other studies have found no direct
968 relationship between these two factors (Velasco *et al.* 1999; Smith *et al.* 2002; Carvalho &
969 Nunes 2006). Greater behavioural exploration levels, feed conversion ratios and growth rates
970 in *L. vannamei* resulted from three to four feedings per day (Pontes *et al.* 2008; Lima *et al.*
971 2009); however, feeding more than twice per day may not be desirable for aquaculture as it is
972 labour-intensive (Carvalho & Nunes 2006). More than four feedings per day could also elicit
973 a loss of stimulation to the shrimp and reduce overall feeding. As previously mentioned,
974 automatic feeders are being implemented more frequently in *L. vannamei* aquaculture (e.g.
975 AQ1 systems). Jescovitch *et al.* (2018), testing different feeding protocols, determined that
976 automatic feeders produced the best results in terms of pond biomass of shrimp. However,
977 they also resulted in the highest levels of nitrogenous compounds. This technology does
978 however present a large potential application for shrimp farming, but its use must be properly
979 adjusted to shrimp feeding activity, an area where behavioural studies could provide useful
980 information.

981

982 A full analysis of daily behavioural feeding patterns can help refine feeding practices (Lima
983 *et al.* 2009). Verifying the time of the day when shrimp are most likely to perform searching

984 behaviours allows feed provision at a time when shrimp are most likely to eat it (Da Costa *et*
985 *al.* 2016). For example, introduction of feed elicited feeding behaviour arousal and
986 exploration in *L. vannamei* within 30 minutes, with the most efficient feeding rates occurring
987 in the morning and afternoon compared to the night (Pontes & Arruda 2005a). Although *L.*
988 *vannamei* individuals were more active during the night, this was not influenced by the
989 presence of food. Pontes *et al.* (2006) showed feeding *L. vannamei* once a day at random
990 times resulted in substrate exploration in both light and dark phases, suggesting that the
991 search for food occurs both during the day and night. Studies with other shrimp species, such
992 as the Southern brown shrimp (*Penaeus subtilis* Pérez Farfante), have found the greatest
993 feeding activity 10 minutes after onset (Nunes *et al.* 1997) and that peaks in ingestion of
994 natural food occurred around 30 minutes after the addition of artificial feed (Nunes *et al.*
995 1996). Taken together, these behavioural results suggest that feeding times spread out over
996 the course of the day may stimulate search behaviours and lead to greater ingestion of feed
997 and is why continuous feeding is now often used in shrimp farming. There are many other
998 factors which may determine the success of food distribution, however, few have been
999 studied in detail in shrimp aquaculture. In finfish aquaculture, distribution of food and
1000 feeding times is an important consideration, for example, in the formation of social
1001 hierarchies (Chandroo *et al.* 2004; Bégout *et al.* 2012; Martins *et al.* 2012), and similar
1002 behaviours may affect shrimp feeding.

1003 **Table 2.** Studies on the behaviour of shrimp in response to feeding effectors. Water quality parameters that shrimp were held under prior to and/or during behavioural
 1004 observations are given, whether behaviours were observed in shrimp held individually or in groups and whether anything is known about the moult stage or sex. The
 1005 presence of any substrate or environmental enrichment is noted. Feeding effectors listed are the most effective for that particular study.

Feeding Effectors (selection)	Inclusion level (%)	Base Feed	Species	Behaviours Measured	Water Quality Parameters	Juveniles/Adults	Moult Stage	Single (S) or Group (G)	Substrate or Environmental Enrichment	Reference
Taurine Yeast extract	1.5	Control diet (using 1.5% α -cellulose in place of an attractant)	<i>Penaeus monodon</i>	Diets in paired choice	29 \pm 1°C, 25 ppt, 12:12 h	Juveniles	-	G, 8 shrimp	30% water exchange/day	Hartati & Briggs, 1993
Heat-dried squid Krill hydrolysate Fish hydrolysate	3, 9, 15	Commercial feed (35% CP)	<i>Litopenaeus vannamei</i>	None	28 \pm 0.5°C, 32 ppt	Juveniles	-	G, 8 shrimp	80% water exchange/day	Córdova-Murueta & García- Carreño, 2002
Krill meal	4	Wheat-flour-casein based feed (16% and 45% CP)	<i>Litopenaeus vannamei</i>	Feeding activity within feeding trays (predetermined time intervals)	28-29°C, 27.3-28.2 ppt, DO 5.8-6 mg/l, 12:12h	Adults	Same moult status	G, 50 shrimp	50% water exchange/day	Sanchez <i>et al.</i> , 2005
Squid meal Crustacean meal Krill meal	1, 2.5, 5	Base feed (terrestrial source)	<i>Penaeus monodon</i>	Diets in paired choice using feeding trays	27 \pm 2°C, 12:12h	Juveniles	Inter-moult	G, 12 shrimp	-	Smith <i>et al.</i> , 2005
Amino acid complex Whole squid protein hydrolysate Condensed fish soluble protein	0.5, 1, 3	Neutral gelatine pellets	<i>Litopenaeus vannamei</i>	Detection, orientation+ locomotion and feeding activity with paired choice using Y-maze	33 ppt	Juveniles	-	S	-	Nunes <i>et al.</i> , 2006

Dimethyl sulphide Trimethylamine oxide Trimethylamine	0.5, 0.01	Commercial feed	<i>Penaeus monodon</i>	Time spent to approach and pick up the feed	27±0.2°C, 30±2ppt, pH 8.0±1	Juveniles	-	G, 5 shrimp	-	Ahamad-Ali <i>et al.</i> , 2007
Squid liver meal Krill meal	3	Poultry by- product meal at 20% + base feed (46% CP)	<i>Penaeus stylirostris</i>	Diets in paired comparisons (dietary choice)	28-31°C, 27-30 ppt, pH 7-8.2, DO > 5mg/l	Juveniles	-	G, 10 shrimp	-	Suresh <i>et al.</i> , 2011
Krill meal	0, 1, 3, 6	Commercial feed	<i>Litopenaeus vannamei</i>	Consumption rate within 60 and 180 min	25-27°C, 32- 35 ppt, 12:12 h	Juveniles	-	S G, 36 shrimp	-	Derby <i>et al.</i> , 2016
Krill meal solution	0.00133, 0.0133, 0.133, 1.33, 13.3 mg/ml	Commercial feed	<i>Litopenaeus vannamei</i>	Specific behaviours measured such as 'Probe', 'Grab' in response to food.	25-27°C, 32- 35 ppt, 12:12 h	Juveniles	-	S G, 36 shrimp	-	Derby <i>et al.</i> , 2016

1006

1007

1008 **CONCLUSIONS**

1009

1010 Solutions to the problems associated with feed management and efficiency in crustaceans
1011 have been addressed by several nutritional and behavioural studies in crustacean aquaculture.
1012 However, what is clear from this work is it is vital that nutritional and behavioural studies are
1013 developed together to understand, for example, how the study of chemical components is
1014 important in enhancing feeding behaviours, such as arousal, search initiation and food
1015 location (Derby *et al.* 2016). Considering these types of studies in combination may also
1016 useful for improving palatability of commercial feeds and enhancing ingestion (Holland &
1017 Borski 1993; Lee & Meyers 1996; Samocha *et al.* 2004a; Sanchez *et al.* 2005; Smith *et al.*
1018 2005; Suresh *et al.* 2011).

1019

1020 Very little work has thoroughly examined the sequence of feeding behaviour patterns, which
1021 can only be achieved through continuous behavioural monitoring linked to different feeding
1022 phases. In *L. vannamei*, nutritional studies have shown how different strategies, such as
1023 changing culture parameters (e.g. Rosas *et al.* 2001; Zhang *et al.* 2006; Bett & Vinatea 2009;
1024 Ponce-Palafox *et al.* 2013) or testing novel additives (e.g. Córdova-Murueta & García-
1025 Carreño 2002; Sanchez *et al.* 2005; Nunes *et al.* 2006; Derby *et al.* 2016), can help solve
1026 problems associated with feeding efficiency in the species. However, these studies are mainly
1027 focused on outputs such as survival or growth rate, with very few studies obtaining data on
1028 feeding behaviour patterns (e.g. Pontes & Arruda 2005a,b; Nunes *et al.* 2006). In addition,
1029 work is often carried out under laboratory conditions which then require scaling up to
1030 realistic farming conditions (Tacon 1996). There is also a need to focus research at the
1031 species level as many of the influences on shrimp behaviour will be species-specific. The
1032 majority of studies to date have not controlled for, or at least not reported, individual

1033 variation in behavioural response, thus these influences are largely unaccounted for in our
1034 understanding.

1035

1036 Behavioural research in finfish aquaculture is extensive (e.g. Chandroo *et al.* 2004;
1037 Dingemans *et al.* 2009; Bégout *et al.* 2012; Martins *et al.* 2012; Ibarra-Zatarain *et al.* 2016),
1038 and while driven primarily by welfare considerations, has led to significant refinements in
1039 feeding practice and efficiency. Recent reviews highlight the importance of applied behaviour
1040 in finfish aquaculture. For example, Zhou *et al.* (2017) defended the use of technological
1041 feeding control devices using behavioural research. Castanheira *et al.* (2017) highlighted the
1042 importance of coping styles in finfish aquaculture, and the idea that proactive (i.e. bold) and
1043 reactive (i.e. shy) fish have different fundamental requirements. Selection or accommodation
1044 of desirable traits can, therefore, increase growth and reduce aggression within an aquaculture
1045 environment. Our understanding of personality and associated behavioural traits in
1046 crustaceans is extensive, yet the literature is lacking a focus towards species in aquaculture.
1047 Thus, a large potential exists to improve feeding efficiencies in shrimp aquaculture with
1048 focused research on shrimp behaviour.

1049

1050

1051 **REFERENCES**

1052

1053 Accioly IV, Lima-Filho PA, Santos TL, Barbosa ACA, Campos LBS, Souza JV, Araújo WC,
1054 Wagner, FM (2013) Sexual dimorphism in *Litopenaeus vannamei* (Decapoda)
1055 identified by geometric morphometrics. *Pan-American Journal of Aquatic Sciences* **8**:
1056 276-281.

1057 Ache BW (1982) Chemoreception and Thermoreception. In: Atwood HL, Sandeman DC
1058 (eds.) *The Biology of Crustacea, vol 3. Neurobiology: Structure and Function*, pp. 369-
1059 398. Academic Press, NY, USA.

1060 Aggio JF, Tieu R, Wei A, Derby CD (2012) Oesophageal chemoreceptors of blue crabs,
1061 *Callinectes sapidus*, sense chemical deterrents and can block ingestion of food. *The*
1062 *Journal of Experimental Biology* **215**:1700-1710.

1063 Ahamad-Ali S, Gopal C, Ramana JV (2007) Attractant and growth promoting properties of
1064 some feed materials and chemicals incorporated in the diets for *Penaeus monodon*
1065 (Fabricius). *Indian Journal of Fisheries* **54**: 67-73.

1066 Allan GL, Maguire GB (1991) Lethal levels of low dissolved oxygen and effects of short-
1067 term oxygen stress on subsequent growth of juvenile *Penaeus*
1068 *monodon*. *Aquaculture* **94**: 27-37.

1069 Allan GL, Maguire GB (1992) Effects of pH and salinity on survival, growth and
1070 osmoregulation in *Penaeus monodon* Fabricius. *Aquaculture* **107**: 33-47.

1071 Allan GL, Maguire GB (1995) Effect of sediment on growth and acute ammonia toxicity for
1072 the school prawn, *Metapenaeus macleayi* (Haswell). *Aquaculture* **131**: 59-71.

1073 Allen WV (1971) Amino acid and fatty acid composition of tissues of the Dungeness crab
1074 (*Cancer magister*). *Journal of the Fisheries Board of Canada* **28**: 1191-1195.

1075 Allison V, Dunham DW, Harvey HH (1992) Low pH alters response to food in the crayfish
1076 *Cambarus bartoni*. *Canadian Journal of Zoology* **70**: 2416-2420.

1077 Amaya EA, Davis DA, Rouse DB (2007) Replacement of fish meal in practical diets for the
1078 Pacific white shrimp (*Litopenaeus vannamei*) reared under pond conditions.
1079 *Aquaculture* **262**: 393-401.

1080 Ameyaw-Akumfi C (1977) Feeding chemoreceptor sites in the crayfish *Procambarus clarkii*
1081 (Girard). *Crustaceana* **33**: 259-264.

- 1082 Anger K (1996) Salinity tolerance of the larvae and first juveniles of a semiterrestrial grapsid
1083 crab, *Armases miersii* (Rathbun). *The Journal of Experimental Marine Biology and*
1084 *Ecology* **202**: 205-223.
- 1085 Aquacop, Bedier E, Soyez C (1988) Effects of dissolved oxygen concentration on survival
1086 and growth of *Penaeus vannamei* and *Penaeus stylirostris*. *Journal of the World*
1087 *Aquaculture Society* **19**: 13A.
- 1088 Armitage KB, Wall TJ (1982) The effects of body size, starvation and temperature
1089 acclimation on oxygen consumption of the crayfish *Orconectes nais*. *Comparative*
1090 *Biochemistry and Physiology Part A: Physiology* **73**: 63-68.
- 1091 Arnold SJ, Sellars MJ, Crocos PJ, Coman GJ (2005) Response of juvenile brown tiger shrimp
1092 (*Penaeus esculentus*) to intensive culture conditions in a flow through tank system with
1093 three-dimensional artificial substrate. *Aquaculture* **246**: 231-238.
- 1094 Arnold SJ, Sellars MJ, Crocos PJ, Coman GJ (2006) Intensive production of juvenile tiger
1095 shrimp *Penaeus monodon*: an evaluation of stocking density and artificial substrates.
1096 *Aquaculture* **261**: 890-896.
- 1097 Atema J (1985) Chemoreception in the sea: adaptations of chemoreceptors and behaviour to
1098 aquatic stimulus conditions. *Symposia of the Society for Experimental Biology* **39**: 387-
1099 423.
- 1100 Atema J (1995) Chemical signals in the marine environment: dispersal, detection and
1101 temporal signal analysis. *Proceeding of the National Academy of Sciences of the United*
1102 *States of America* **92**: 62-66.
- 1103 Bador RF (2000) Uso de charolas de alimentación para el cultivo de Camarón en Sudamérica.
1104 In: Civera-Cerecedo R, Pérez-Estrada CJ, Ricque-Marie D, Cruz-Suárez LE (eds.)
1105 *Avances en Nutrición Acuícola IV*, pp. 540-549. Memorias del IV Simposium

1106 Internacional de Nutrición Acuícola, Universidad Autónoma de Nuevo León, La Paz,
1107 México.

1108 Bador R, Blyth P, Dodd R (2013) Acoustic control improves feeding productivity at shrimp
1109 farms. *Global Aquaculture Advocate* [Cited 30 May 2018]. Available from URL:
1110 [https://www.aquaculturealliance.org/advocate/acoustic-control-improves-feeding-](https://www.aquaculturealliance.org/advocate/acoustic-control-improves-feeding-productivity-at-shrimp-farms/)
1111 [productivity-at-shrimp-farms/](https://www.aquaculturealliance.org/advocate/acoustic-control-improves-feeding-productivity-at-shrimp-farms/).

1112 Balakrishnan G, Peyail S, Ramachandran K, Theivasigamani A, Savji KA, Chokkaiah M,
1113 Nataraj P (2011) Growth of cultured white leg shrimp *Litopenaeus vannamei* (Boone
1114 1931) in different stocking density. *Advances in Applied Science Research* **2**:107-113.

1115 Ballester ELC, Wasielesky W, Cavalli RO, Abreu PC (2007) Nursery of the pink shrimp
1116 *Farfantepenaeus paulensis* in cages with artificial substrates: biofilm composition and
1117 shrimp performance. *Aquaculture* **269**: 355-362.

1118 Barbato JC, Daniel PC (1997) Chemosensory activation of an antennular grooming behavior
1119 in the spiny lobster, *Panulirus argus*, is tuned narrowly to L-glutamate. *The Biological*
1120 *Bulletin* **193**:107-115.

1121 Barbieri E (2010) Acute toxicity of ammonia in white shrimp (*Litopenaeus schmitti*)
1122 (Burkenroad, 1936, Crustacea) at different salinity levels. *Aquaculture* **306**: 329-333.

1123 Bégout ML, Kadri S, Huntingford F, Damsgård B (2012) Tools for studying the behaviour of
1124 farmed fish. In: Huntingford F, Jobling M, Kadri S (eds.) *Aquaculture and Behavior*,
1125 pp. 65-86. Wiley-Blackwell Publishing, Oxford.

1126 Bett C, Vinatea L (2009) Combined effect of body weight, temperature and salinity on
1127 shrimp *Litopenaeus vannamei* oxygen consumption rate. *Brazilian Journal of*
1128 *Oceanography* **57**: 305-314.

- 1129 Blewett TA, Goss GG (2017) A novel pathway of nutrient absorption in crustaceans:
1130 branchial amino acid uptake in the green shore crab (*Carcinus maenas*). *Proceedings of*
1131 *the Royal Society of London B: Biological Sciences* **284**: 20171298.
- 1132 Boddeke R (1983) Survival strategies of penaeid shrimps and their significance for shrimp
1133 culture. In: Roger GL, Day R, Lim A (eds.), *Proceeding of the First International*
1134 *Conference of Warm water Aquaculture*, pp. 514-523. Louisiana, USA.
- 1135 Bondad-Reantaso MG, Subasinghe RP, Josupeit H, Cai J, Zhou X (2012) The role of
1136 crustacean fisheries and aquaculture in global food security: past, present and
1137 future. *Journal of Invertebrate Pathology* **110**:158-165.
- 1138 Boyd CE, Tucker C, McNevin A, Bostick K, Clay J (2007) Indicators of resource use
1139 efficiency and environmental performance in fish and crustacean aquaculture. *Reviews*
1140 *in Fisheries Science* **15**: 327-360.
- 1141 Bratland S, Stien LH, Braithwaite VA, Juell JE, Folkedal O, Nilsson J, Oppedal F,
1142 Fosseidengen JE, Kristiansen TS (2010) From fright to anticipation: using aversive
1143 light stimuli to investigate reward conditioning in large groups of Atlantic salmon
1144 (*Salmo salar*). *Aquaculture International* **18**: 991 – 1001.
- 1145 Bratvold D, Browdy CL (2001) Effects of sand sediment and vertical surfaces (AquaMats™)
1146 on production, water quality, and microbial ecology in an intensive *Litopenaeus*
1147 *vannamei* culture system. *Aquaculture* **195**: 81-94.
- 1148 Bray WA, Lawrence AL, Leung-Trujillo JR (1994) The effect of salinity on growth and
1149 survival of *Penaeus vannamei*, with observations on the interaction of IHHN virus and
1150 salinity. *Aquaculture* **122**: 133-146.
- 1151 Briffa M, Rundle SD, Fryer A (2008) Comparing the strength of behavioural plasticity and
1152 consistency across situations: animal personalities in the hermit crab *Pagurus*

1153 *bernhardus*. *Proceedings of the Royal Society of London B: Biological Sciences* **275**:
1154 1305-1311.

1155 Briffa M, Weiss A (2010) Animal personality. *Current Biology* **20**: 912-914.

1156 Briffa M, Twyman C (2011) Do I stand out or blend in? Conspicuousness awareness and
1157 consistent behavioural differences in hermit crabs. *Biology Letters* **7**: 330-332.

1158 Briffa M, Sneddon LU, Wilson AJ (2015) Animal personality as a cause and consequence of
1159 contest behaviour. *Biology Letters* **11**: 20141007.

1160 Briffa M, Sneddon LU (2016) Proximate mechanisms of animal personality among-
1161 individual behavioural variation in animals. *Behaviour* **153**: 1509-1515.

1162 Briggs M, Funge-Smith S (1994) A nutrient budget of some intensive marine shrimp ponds in
1163 Thailand. *Aquaculture and Fisheries Management* **25**: 789-811.

1164 Briggs M, Funge-Smith S, Subasinghe R, Phillips M (2004) *Introduction and Movement of*
1165 *Penaeus vannamei and Penaeus stylirostris in Asia and the Pacific*. FAO RAP
1166 Publication 2004/ 10. FAO RAP, Bangkok, Thailand.

1167 Burford MA, Sellars MJ, Arnold SJ, Keys SJ, Crocos PJ, Preston NP (2004) Contribution of
1168 the natural biota associated with substrates to the nutritional requirements of the post-
1169 larval shrimp, *Penaeus esculentus* (Haswell), in high-density rearing systems.
1170 *Aquaculture Research* **35**: 508-515.

1171 Campos-Ramos R, Garza-Torres R, Guerrero-Tortolero DA, Maeda-Martínez AM, Obregón-
1172 Barboza H (2006) Environmental sex determination, external sex differentiation and
1173 structure of the androgenic gland in the Pacific white shrimp *Litopenaeus vannamei*
1174 (Boone). *Aquaculture Research* **37**: 1583-1593.

1175 Carr WES (1978) Chemoreception in the shrimp, *Palaemonetes pugio*: the role of amino
1176 acids and betaine in elicitation of a response by extracts. *Comparative Biochemistry*
1177 *and Physiology Part A: Physiology* **61**: 127-131.

1178 Carr WES, Netherton III JC, Milstead ML (1984) Chemoattractants of the shrimp,
1179 *Palaemonetes pugio*: Variability in responsiveness and the stimulatory capacity of
1180 mixtures containing amino acids, quaternary ammonium compounds, purines and other
1181 substances. *Comparative Biochemistry and Physiology Part A: Physiology* **77**: 469-474.

1182 Carr WES, Derby CD (1986) Chemically stimulated feeding behaviour in marine animals:
1183 importance of chemical mixture and involvement of mixture interactions. *Journal of*
1184 *Chemical Ecology* **12**: 989-1011.

1185 Carvalho EA, Nunes AJP (2006) Effects of feeding frequency on feed leaching loss and
1186 grow-out patterns of the white shrimp *Litopenaeus vannamei* fed under a diurnal
1187 feeding regime in pond enclosures. *Aquaculture* **252**: 494-502.

1188 Castanheira MF, Herrera M, Costas B, Conceição LEC, Martins CIM (2013) Linking cortisol
1189 responsiveness and aggressive behaviour in gilthead seabream *Sparus aurata*:
1190 indication of divergent coping styles. *Applied Animal Behaviour Science* **143**: 75 – 81.

1191 Castanheira MF, Conceição LEC, Millot S, Rey S, Bégout ML, Damsgård B, Kristiansen T,
1192 Höglund E, Øverli Ø, Martins CIM (2017). Coping styles in farmed fish: consequences
1193 for aquaculture. *Reviews in Aquaculture* **9**: 23-41.

1194 Cate HS, Derby CD (2001) Morphology and distribution of setae on the antennules of the
1195 Caribbean spiny lobster *Panulirus argus* reveal new types of bimodal chemo-
1196 mechanosensilla. *Cell and Tissue Research* **304**: 439-454.

1197 Chamberlain GW, Lawrence AL (1981) Maturation, reproduction, and growth of *Penaeus*
1198 *vannamei* and *P. stylirostris* fed natural diets. *Journal of the World Aquaculture*
1199 *Society* **12**: 209-224.

1200 Chamberlain GW (2010) History of shrimp farming. In: Alday-Sanz V (ed.) *The Shrimp*
1201 *Book*, pp. 1-34. Nottingham University Press, United Kingdom.

- 1202 Chan S, Rankin SM, Keeley LL (1988) Characterization of the moult stages in *Penaeus*
1203 *vannamei*: Setogenesis and hemolymph levels of total protein, ecdysteroids, and
1204 glucose. *The Biological Bulletin* **175**:185-192.
- 1205 Chandroo KP, Duncan IJH, Moccia RD (2004) Can fish suffer?: perspectives on sentience,
1206 pain, fear and stress. *Applied Animal Behaviour Science* **86**: 225-250.
- 1207 Chanratchakool P, Turnbull JF, Funge-Smith SJ, Limsuwan C (1998) *Health Management in*
1208 *Shrimp Ponds*, 3rd edn. Aquatic Animal Health Research Institute, Bangkok, Thailand.
- 1209 Charmantier G (1987) L'osmoregulation chez les crevettes Penaeidae (Crustacea, Decapoda).
1210 *Oceanis* **13**: 179-196.
- 1211 Charmantier G, Soyez C, Aquacop (1994) Effect of molt stage and hypoxia on
1212 osmoregulatory capacity in the penaeid shrimp *Penaeus vannamei*. *The Journal of*
1213 *Experimental Marine Biology and Ecology* **178**: 233-246.
- 1214 Charmantier G (1998) Ontogeny of osmoregulation in crustaceans: a review. *Invertebrate*
1215 *Reproduction and Development* **33**: 177-190.
- 1216 Chavanich S, Viyakarn V, Senanan W, Panutrakul S (2016) Laboratory assessment of
1217 feeding-behavior interactions between the introduced Pacific white shrimp *Litopenaeus*
1218 *vannamei* (Boone, 1931) (Penaeidae) and five native shrimps plus a crab species in
1219 Thailand. *Aquatic Invasions* **11**: 67-74.
- 1220 Chen J, Lei S (1990) Toxicity of ammonia and nitrite to *Penaeus mondon* juveniles. *Journal*
1221 *of the World Aquaculture Society* **21**: 300-306.
- 1222 Chen J, Kou Y (1992) Effects of ammonia on growth and molting of *Penaeus japonicus*
1223 juveniles. *Aquaculture* **104**: 249-260.
- 1224 Chen S, Chen J (2003) Effects of pH on survival, growth, molting and feeding of giant
1225 freshwater prawn *Macrobrachium rosenbergii*. *Aquaculture* **218**: 613-623.

- 1226 Cheng W, Liu C, Yan D, Chen J (2002) Hemolymph, oxyhemocyanin, protein, osmolality
1227 and electrolyte levels of whiteleg shrimp *Litopenaeus vannamei* in relation to size and
1228 molt stage. *Aquaculture* **211**: 325-339.
- 1229 Cheng W, Liu C, Kuo C (2003) Effects of dissolved oxygen on hemolymph parameters of
1230 freshwater giant prawn, *Macrobrachium rosenbergii* (de Man). *Aquaculture* **220**: 843-
1231 856.
- 1232 Cheroske AG, Cronin TW, Durham MF, Caldwell RL (2009) Adaptive signaling behavior in
1233 stomatopods under varying light conditions. *Marine and Freshwater Behaviour and*
1234 *Physiology* **42**: 219-232.
- 1235 Chow S, Sandifer PA (1991) Differences in growth, morphometric traits, and male sexual
1236 maturity among Pacific white shrimp, *Penaeus vannamei*, from different commercial
1237 hatcheries. *Aquaculture* **92**: 165-178.
- 1238 Clark JV (1986) Inhibition of moulting in *Penaeus semisulcatus* (De Haan) by long-term
1239 hypoxia. *Aquaculture* **52**: 253-254.
- 1240 Colt JE, Armstrong DA (1981) Nitrogen toxicity to crustaceans, fish and molluscs. In: Allen
1241 LJ, Kinney EC (eds.) *Proceedings of the Bio-Engineering Symposium for Fish Culture.*
1242 *Fish Culture Section, American Fisheries Society*, pp. 34-47. Northeast Society of
1243 Conservation Engineers, Bethesda, Maryland, USA.
- 1244 Comoglio LI, Gaxiola G, Roque A, Cuzon G, Amin O (2004) The effect of starvation on
1245 refeeding, digestive enzyme activity, oxygen consumption, and ammonia excretion in
1246 juvenile white shrimp *Litopenaeus vannamei*. *Journal of Shellfish Research* **23**: 243-
1247 249.
- 1248 Comoglio L, Smolko L, Amin O (2005) Effects of starvation on oxygen consumption,
1249 ammonia excretion and biochemical composition of the hepatopancreas on adult males
1250 of the false southern king crab *Paralomis granulosa* (Crustacea, Decapoda).

- 1251 *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular*
1252 *Biology* **140**: 411-416.
- 1253 Córdova-Murueta JH, García-Carreño FL (2002) Nutritive value of squid and hydrolyzed
1254 protein supplement in shrimp feed. *Aquaculture* **210**: 371-384.
- 1255 Costero M, Meyers SP (1993) Evaluation of chemoreception by *Penaeus vannamei* under
1256 experimental conditions. *The Progressive Fish-Culturist* **55**:157-162.
- 1257 Cruz-Suárez LE, Nieto-López M, Guajardo-Barbosa C, Tapia-Salazar M, Scholz U, Ricque-
1258 Marie D (2007) Replacement of fish meal with poultry by-product meal in practical
1259 diets for *Litopenaeus vannamei*, and digestibility of the tested ingredients and diets.
1260 *Aquaculture* **272**: 466-476.
- 1261 Cuéllar-Anjel J, Lara C, Morales V, De Gracia A, García-Suárez O (2010) *Manual of best*
1262 *management practices for white shrimp Penaeus vannamei farming*, 1st edn. New
1263 Concept Publications, Panama.
- 1264 Cutts CJ, Metcalfe NB, Taylor AC (1998) Aggression and growth depression in juvenile
1265 Atlantic salmon: the consequences of individual variation in standard metabolic
1266 rate. *Journal of Fish Biology* **52**: 1026-1037.
- 1267 Cuzon G, Cahu C, Aldrin JF, Messenger JL, Stéphan G, Mével M (1980) Starvation effect on
1268 metabolism of *Penaeus japonicus*. *Proceedings of the World Mariculture Society* **11**:
1269 410-423.
- 1270 Cuzon G, Lawrence A, Gaxiola G, Rosas C, Guillaume J (2004) Nutrition of *Litopenaeus*
1271 *vannamei* reared in tanks or in ponds. *Aquaculture* **235**: 513-551.
- 1272 Da Costa FP, Gomes BSF, Pereira SDNA, Arruda, FM (2016) Influence of stocking density
1273 on the behaviour of juvenile *Litopenaeus vannamei* (Boone, 1931). *Aquaculture*
1274 *Research* **47**: 912-924.

- 1275 Dall W (1981) Osmoregulatory ability and juvenile habitat preference in some penaeid
1276 prawns. *The Journal of Experimental Marine Biology and Ecology* **54**: 55-64.
- 1277 Dall W (1986) Estimation of routine metabolic rate in a penaeid prawn, *Penaeus esculentus*
1278 Haswell. *The Journal of Experimental Marine Biology & Ecology* **96**: 57-74.
- 1279 Dall W, Hill BJ, Rothlisberg PC, Staples DJ (1990) *The Biology of the Penaeidae. Advances*
1280 *in Marine Biology*. Academic Press, London, UK.
- 1281 Daniel PC, Derby CD (1988) Behavioral olfactory discrimination of mixtures in the spiny
1282 lobster (*Panulirus argus*) based on a habituation paradigm. *Chemical Senses* **13**: 385-
1283 395.
- 1284 Daniel PC, Derby CD (1991) Mixture suppression in behaviour: The antennular flick
1285 response in the spiny lobster towards binary odorant mixtures. *Physiology & Behavior*
1286 **49**: 591-601.
- 1287 Darsey KT (1990) *The influence of temperature, salinity on the growth and survival of*
1288 *California brown shrimp Penaeus californiensis* (Holmes). Doctoral dissertation, M.
1289 Sc. Thesis, University of Arizona, USA.
- 1290 Das S, Sahu BK (2005) Interaction of pH with mercuric chloride toxicity to penaeid prawns
1291 from a tropical estuary, East Coast of India: enhanced toxicity at low pH. *Chemosphere*
1292 **58**: 1241-1248.
- 1293 Davis DA, Arnold CR (2000) Replacement of fish meal in practical diets for the Pacific white
1294 shrimp, *Litopenaeus vannamei*. *Aquaculture* **185**: 291-298.
- 1295 Davis DA, Saoud IP, McGraw WJ, Rouse DB (2002) Considerations for *Litopenaeus*
1296 *vannamei* reared in inland low salinity waters. In: Cruz-Suárez LE, Ricque-Marie D,
1297 Tapia-Salazar M, Gaxiola-Cortes MG, Simoes N (eds.) *Avances en Nutrición Acuícola*
1298 *VI*, pp. 73-90. Memorias del VI Simposium Internacional de Nutrición Acuícola,
1299 Cancún, Quintana Roo, México.

- 1300 Davis DA, Amaya E, Venero J, Zelaya O, Rouse DB (2006) A case study on feed
1301 management to improving production and economic returns for the semi-intensive
1302 pond production of *Litopenaeus vannamei*. In: Cruz-Suárez LE, Ricque-Marie D,
1303 Tapia-Salazar M, Nieto-López MG, Villarreal-Cavazos DA, Puello-Cruz AN, García-
1304 Ortega A (eds.) *Avances en Nutrición Acuícola VIII*, pp. 282-303. Memorias del VIII
1305 Simposium Internacional de Nutrición Acuícola, Universidad Autónoma de Nuevo
1306 León, Monterrey, Nuevo León, México.
- 1307 De la Haye KL, Spicer JI, Widdicombe S, Briffa M (2011) Reduced sea water pH disrupts
1308 resource assessment and decision making in the hermit crab *Pagurus bernhardus*.
1309 *Animal Behaviour* **82**: 495-501.
- 1310 De la Haye KL, Spicer JI, Widdicombe S, Briffa M (2012) Reduced pH sea water disrupts
1311 chemo-responsive behaviour in an intertidal crustacean. *The Journal of Experimental*
1312 *Marine Biology & Ecology* **412**:134-140.
- 1313 De Oliveira-Cesar JR, Zhao B, Malecha S, Ako H, Yang J (2006) Morphological and
1314 biochemical changes in the muscle of the marine shrimp *Litopenaeus vannamei* during
1315 the molt cycle. *Aquaculture* **261**: 688-694.
- 1316 Dennenmoser S, Thiel M (2007) Competition for food and mates by dominant and
1317 subordinate male rock shrimp, *Rhynchocinetes typus*. *Behaviour* **144**: 33-59.
- 1318 Derby CD, Atema J (1981) Selective improvement in responses to prey odors by the lobster,
1319 *Homarus americanus*, following feeding experience. *Journal of chemical ecology* **7**:
1320 1073-1080.
- 1321 Derby CD, Atema J (1982) The function of chemo- and mechanoreceptors in lobster
1322 (*Homarus americanus*) feeding behaviour. *The Journal of Experimental Biology* **98**:
1323 317-327.

- 1324 Derby CD (2000) Learning from spiny lobsters about chemosensory coding of mixtures.
1325 *Physiology & Behaviour* **69**: 203-209.
- 1326 Derby CD, Steullet P (2001) Why do animals have so many receptors? the role of multiple
1327 chemosensors in animal perception. *The Biological Bulletin* **200**: 211-215.
- 1328 Derby CD, Steullet P, Horner AJ, Cate HS (2001) The sensory basis of feeding behaviour in
1329 the Caribbean spiny lobster, *Panulirus argus*. *Marine & Freshwater Research* **52**:
1330 1339-1350.
- 1331 Derby CD, Sorensen PW (2008) Neural processing, perception, and behavioural responses to
1332 natural chemical stimuli by fish and crustaceans. *Journal of Chemical Ecology* **34**: 898-
1333 914.
- 1334 Derby CD, Elsayed FH, Williams SA, González C, Choe M, Bharadwaj AS, Chamberlain
1335 GW (2016) Krill meal enhances performance of feed pellets through concentration-
1336 dependent prolongation of consumption by Pacific white shrimp, *Litopenaeus*
1337 *vannamei*. *Aquaculture* **458**: 13-20.
- 1338 Dersjant-Li Y (2002) The use of soy protein in aquafeeds. In: Cruz-Suárez LE, Ricque-Marie
1339 D, Tapia-Salazar M, Gaxiola-Cortés MG, Simoes N (eds.) *Avances en Nutrición*
1340 *Acuícola VI*, pp. 541-558. Memorias del VI Simposium Internacional de Nutrición
1341 Acuícola, Cancún, Quintana Roo, México.
- 1342 Devine DV, Atema J (1982) Function of chemoreceptor organs in spatial orientation of the
1343 lobster, *Homarus americanus*: differences and overlap. *The Biological Bulletin* **163**:
1344 144-153.
- 1345 DeWitt TJ, Sih A, Wilson DS (1998) Costs and limits of phenotypic plasticity. *Trends in*
1346 *Ecology & Evolution* **13**: 77-81.

- 1347 Díaz GA, Smith SG, Serafy JE, Ault JS (2001) Allometry of the growth of pink shrimp
1348 *Farfantepenaeus duorarum* in a subtropical bay. *Transactions of the American*
1349 *Fisheries Society* **130**: 328-335.
- 1350 Dingemanse NJ, Van der Plas F, Wright J, Réale D, Schrama M, Roff DA, Van der Zee E,
1351 Barber I (2009) Individual experience and evolutionary history of predation affect
1352 expression of heritable variation in fish personality and morphology. *Proceedings of the*
1353 *Royal Society of London B: Biological Sciences* **276**: 1285-1293.
- 1354 Dingemanse NJ, Kazem AJ, Réale D, Wright J (2010) Behavioural reaction norms: animal
1355 personality meets individual plasticity. *Trends in Ecology & Evolution* **25**: 81-89.
- 1356 Dunham DW, Ciruna KA, Harvey HH (1997) Chemosensory role of antennules in the
1357 behavioural integration of feeding by the crayfish *Cambarus bartonii*. *Journal of*
1358 *Crustacean Biology* **17**: 27-32.
- 1359 Engdahl S (1997) Effects of manganese and hypoxia on the chemoreception behaviour of the
1360 Norway lobster, *Nephrops norvegicus*. Master Thesis. Department of Marine Ecology,
1361 Göteborg University, Sweden.
- 1362 Engle CR, McNevin A, Racine P, Boyd CE, Paungkaew D, Viriyatum R, Tinh HQ, Minh HN
1363 (2017) Economics of sustainable intensification of aquaculture: evidence from shrimp
1364 farms in Vietnam and Thailand. *Journal of the World Aquaculture Society* **48**: 227-239.
- 1365 FAO (2016) *The State of World Fisheries and Aquaculture 2016*. FAO, Rome, Italy.
- 1366 Felix S (2013) *Advances in shrimp aquaculture management*. Daya Publishing House, New
1367 Delhi, India.
- 1368 Fernández I, Oliva M, Carrillo O, Van Wormhoudt A (1997) Digestive enzyme activities of
1369 *Penaeus notialis* during reproduction and moulting cycle. *Comparative Biochemistry &*
1370 *Physiology Part A: Physiology* **118**: 1267-1271.

1371 FIGIS (2015) FAO Statistics. Global Aquaculture Production 1950-2015 [Cited 7 November
1372 2017]. Available from URL: <http://www.fao.org/figis/>.

1373 Flegel TW (2009) Current status of viral diseases in Asian shrimp aquaculture. *The Israeli*
1374 *Journal of Aquaculture - Bamigdeh* **61**: 229-239.

1375 Folkedal O, Fernö A, Nederlof MAJ, Fosseidengen JE, Cerqueira M, Olsen RE, Nilsson J
1376 (2018) Habituation and conditioning in gilthead sea bream (*Sparus aurata*): effects of
1377 aversive stimuli, reward and social hierarchies. *Aquaculture Research* **49**: 335 – 340.

1378 Fox JM, Lawrence AL, Smith F (2004) Development of a low-fish meal feed formulation for
1379 commercial production of *Litopenaeus vannamei*. In: Cruz-Suárez LE, Ricque-Marie
1380 D, Nieto-López MG, Villarreal D, Scholz U, González M (eds.) *Avances en Nutrición*
1381 *Acuícola VII*, pp. 238-258. Memorias del VII Simposium Internacional de Nutrición
1382 Acuícola, Hermosillo, Sonora, México.

1383 Frías-Espéricueta MG, Harfush-Melendez M, Osuna-López JI, Páez-Osuna F (1999) Acute
1384 toxicity of ammonia to juvenile shrimp *Penaeus vannamei* Boone. *Bulletin of*
1385 *Environmental Contamination & Toxicology* **62**: 646-652.

1386 Frías-Espéricueta MG, Harfush-Melendez M, Páez-Osuna F (2000) Effects of ammonia on
1387 mortality and feeding of postlarvae shrimp *Litopenaeus vannamei*. *Bulletin of*
1388 *Environmental Contamination & Toxicology* **65**: 98-103.

1389 Funge-Smith S, Briggs M (2005) The introduction of *Penaeus vannamei* and *P. stylirostris*
1390 into the Asia-Pacific region. In: Bartley DM, Bhujel RC, Funge-Smith S, Olin PG,
1391 Phillips MJ (eds.) *International Mechanisms for the Control and Responsible use of*
1392 *Alien Species in Aquatic Ecosystems, Report of an Ad Hoc Expert Consultation*. Food
1393 and Agriculture Organization of the United Nations (FAO), Rome, Italy.

1394 Furtado PS, Campos BR, Serra FP, Klosterhoff M, Romano LA, Wasielesky W (2015)
1395 Effects of nitrate toxicity in the Pacific white shrimp, *Litopenaeus vannamei*, reared
1396 with biofloc technology (BFT). *Aquaculture International* **23**: 315-327.

1397 Fuss CM, Ogren LH (1966) Factors affecting activity and burrowing habits of the pink
1398 shrimp, *Penaeus duorarum* Burkenroad. *The Biological Bulletin* **130**: 170-191.

1399 Fuzessery ZM, Childress JJ (1975) Comparative chemosensitivity to amino acids and their
1400 role in the feeding activity of bathypelagic and littoral crustaceans. *The Biological*
1401 *Bulletin* **149**: 522-538.

1402 Gadiant M, Schai E (1994) Leaching of various vitamins for shrimp feed. *Aquaculture* **124**:
1403 201-205.

1404 Garcia DK, Faggart MA, Rhoades L, Alcivar-Warren AA, Wyban JA, Carr WH, Sweeney
1405 JN, Ebert KM (1994) Genetic diversity of cultured *Penaeus vannamei* shrimp using
1406 three molecular genetic techniques. *Molecular Marine Biology & Biotechnology* **3**:
1407 270-280.

1408 Gatlin DM, Barrows FT, Brown P, Dabrowski K, Gaylord TG, Hardy RW, Herman E, Hu G,
1409 Krogdahl Å, Nelson R, Overturf K, Rust M, Sealey W, Skonberg D, Souza EJ, Stone D,
1410 Wilson R, Wurtele E (2007) Expanding the utilization of sustainable plant products in
1411 aquafeeds: a review. *Aquaculture Research* **38**: 551-579.

1412 Gherardi F, Aquiloni L, Tricarico E (2012) Behavioral plasticity, behavioral syndromes and
1413 animal personality in crustacean decapods: an imperfect map is better than no
1414 map. *Current Zoology* **58**: 567-579.

1415 Gitterle T, Rye M, Salte R, Cock J, Johansen H, Lozano C, Suárez JA, Gjerde B (2005)
1416 Genetic (co)variation in harvest body weight and survival in *Penaeus (Litopenaeus)*
1417 *vannamei* under standard commercial conditions. *Aquaculture* **243**: 83-92.

1418 Gopal C, Gopikrishna G, Krishna G, Jahageerdar SS, Rye M, Hayes BJ, Paulpandi S, Kiran
1419 RP, Pillai SM, Ravichandran P, Ponniah AG, Kumar D (2010) Weight and time of
1420 onset of female-superior sexual dimorphism in pond reared *Penaeus*
1421 *monodon*. *Aquaculture* **300**: 237-239.

1422 Goldman JA, Koehl MAR (2001) Fluid dynamic design of lobster olfactory organs: high
1423 speed kinematic analysis of antennule flicking by *Panulirus argus*. *Chemical*
1424 *Senses* **26**: 385-398.

1425 Gleeson RA, Carr WES, Trapido-Rosenthal HG (1993) Morphological characteristics
1426 facilitating stimulus access and removal in the olfactory organ of the spiny lobster,
1427 *Panulirus argus*: insight from the design. *Chemical Senses* **18**: 67-75.

1428 Gleeson RA, McDowell LM, Aldrich HC (1996) Structure of the aesthetasc (olfactory)
1429 sensilla of the blue crab, *Callinectes sapidus*: transformations as a function of salinity.
1430 *Cell & Tissue Research* **284**: 279-288.

1431 Gong H, Jiang DH, Lightner DV, Collins C, Brock D (2004) A dietary modification approach
1432 to improve the osmoregulatory capacity of *Litopenaeus vannamei* cultured in the
1433 Arizona desert. *Aquaculture Nutrition* **10**: 227-236.

1434 Grasso FW, Basil JA (2002) How lobsters, crayfishes, and crabs locate sources of odor:
1435 current perspectives and future directions. *Current Opinion in Neurobiology* **12**: 721-
1436 727.

1437 Grey M, Forster I, Dominy W, Ako H, Giesen AF (2009) Validation of a feeding stimulant
1438 bioassay using fish hydrolysates for the Pacific white shrimp, *Litopenaeus vannamei*.
1439 *Journal of the World Aquaculture Society* **40**: 547-555.

1440 Griffith DRW, Wigglesworth JM (1993) Growth rhythms in the shrimp *Penaeus vannamei*
1441 and *P. schmitti*. *Marine Biology* **115**: 295-299.

- 1442 Grünert U, Ache BW (1988) Ultrastructure of the aesthetasc (olfactory) sensilla of the spiny
1443 lobster, *Panulirus argus*. *Cell & Tissue Research* **251**: 95-103.
- 1444 Hagerman L, Uglow RF (1984) The influence of hypoxia on the blood regulation of the
1445 brackish water shrimp *Palaemonetes varians* Leach. *Journal of Experimental Marine*
1446 *Biology & Ecology* **76**: 157-165.
- 1447 Hagerman L, Uglow RF (1985) Effects of hypoxia on the respiratory and circulatory
1448 regulation of *Nephrops norvegicus*. *Marine Biology* **87**: 273-278.
- 1449 Hall D (2004) Explaining the diversity of Southeast Asian shrimp aquaculture. *Journal of*
1450 *Agrarian Change* **4**: 315-335.
- 1451 Hallberg E, Johansson KUI, Elofsson R (1992) The aesthetasc concept: structural variations
1452 of putative olfactory receptor cell complexes in crustacea. *Microscopy Research &*
1453 *Technique* **22**: 325-335.
- 1454 Hansford SW, Hewitt DR (1994) Growth and nutrient digestibility by male and female
1455 *Penaeus monodon*: evidence of sexual dimorphism. *Aquaculture* **125**: 147-154.
- 1456 Hasan MR, New MB (2013) *On-farm feeding and feed management in aquaculture*. FAO
1457 Fisheries & Aquaculture Technical Paper No. 583, Rome, Italy.
- 1458 Hardy RW, Tacon AGJ (2002) Fish meal: historical uses, production trends and future
1459 outlook for sustainable supplies. In: Stickney RR, McVey JP (eds.) *Responsible Marine*
1460 *Aquaculture*, pp. 311-325. CABI Publishing, New York, USA.
- 1461 Harpaz S, Kahan D, Galun R, Moore I (1987) Responses of freshwater prawn,
1462 *Macrobrachium rosenbergii*, to chemical attractants. *Journal of Chemical Ecology*
1463 **13**: 1957-1965.
- 1464 Harpaz S, Steiner JE (1987) Quantitative analysis of feeding behaviour stereotypes and of the
1465 rejection of aversive-tasting food in the freshwater prawn *Macrobrachium*
1466 *rosenbergii*. *Annals of the New York Academy of Sciences* **510**: 347-348.

- 1467 Harpaz S, Steiner JE (1990) Analysis of betaine-induced feeding behaviour in the prawn
1468 *Macrobrachium rosenbergii* (De Man, 1879) (Decapoda, Caridea). *Crustaceana* **58**:
1469 175-185.
- 1470 Hartati R, Briggs MRP (1993) Effect of feeding attractants on the behaviour and performance
1471 of juvenile *Penaeus monodon* Fabricius. *Aquaculture & Fisheries Management* **24**:
1472 613-624.
- 1473 Hay ME (2011) Crustaceans as powerful models in aquatic chemical ecology. In: Breithaupt
1474 T, Thiel M (eds.) *Chemical Communication in Crustaceans*, pp. 41-62. Springer
1475 Science & Business Media, New York, USA.
- 1476 Hazlett BA (1971) Chemical and chemotactic stimulation of feeding behavior in the hermit
1477 crab *Petrochirus diogenes*. *Comparative Biochemistry & Physiology Part A*:
1478 *Physiology* **39**: 665-670.
- 1479 Hazlett BA (1995) Behavioral plasticity in crustacea: why not more?. *The Journal of*
1480 *Experimental Marine Biology & Ecology* **193**: 57-66.
- 1481 Heinen JM (1980) Chemoreception in decapod crustacea and chemical feeding stimulants as
1482 potential feed additives. *Proceedings of the World Mariculture Society* **11**: 319-334.
- 1483 Hernández C, Sarmiento-Pardo J, González-Rodríguez B, de la Parra IA (2004) Replacement
1484 of fish meal with co-extruded wet tuna viscera and corn meal in diets for white shrimp
1485 (*Litopenaeus vannamei* Boone). *Aquaculture Research* **35**: 1153-1157.
- 1486 Hernández M, Bückle LF, Palacios E, Barón B (2006) Preferential behavior of white shrimp
1487 *Litopenaeus vannamei* (Boone 1931) by progressive temperature-salinity simultaneous
1488 interaction. *Journal of Thermal Biology* **31**: 565-572.
- 1489 Hertrampf JW, Piedad-Pascual F (2000) *Handbook on Ingredients for Aquaculture Feeds*, 1st
1490 edn. Kluwer Academic Publishers, Dordrecht, Netherlands.

- 1491 Hervant F, Mathieu J, Barré H, Simon K, Pinon C (1997) Comparative study on the
1492 behavioral, ventilatory, and respiratory responses of hypogean and epigean crustaceans
1493 to long-term starvation and subsequent feeding. *Comparative Biochemistry &*
1494 *Physiology Part A: Physiology* **118**: 1277-1283.
- 1495 Hindley JPR (1975) Effects of endogenous and some exogenous factors on the activity of the
1496 juvenile banana prawn *Penaeus merguensis*. *Marine Biology* **29**: 1-8.
- 1497 Hochachka PW, Lutz PL (2001) Mechanism, origin, and evolution of anoxia tolerance in
1498 animals. *Comparative Biochemistry & Physiology Part B: Biochemistry & Molecular*
1499 *Biology* **130**: 435-459.
- 1500 Holland KN, Borski RJ (1993) A palatability bioassay for determining ingestive stimuli in the
1501 marine shrimp *Penaeus vannamei*. *Aquaculture* **109**: 153-164.
- 1502 Hopkins JS, Stokes AD, Browdy CL, Sandifer PA (1991) The relationship between feeding
1503 rate, paddlewheel aeration rate and expected dawn dissolved oxygen in intensive
1504 shrimp ponds. *Aquacultural Engineering* **10**: 281-290.
- 1505 Horner AJ, Ngo V, Steullet P, Keller T, Weissburg MJ, Derby CD (2000) The role of
1506 different types of antennular sensilla in orientation by Caribbean spiny lobsters to a
1507 natural odor stimulus under controlled flow conditions. *Chemical Senses* **25**: 670
1508 (abstract).
- 1509 Horner AJ, Weissburg MJ, Derby CD (2004) Dual antennular chemosensory pathways can
1510 mediate orientation by Caribbean spiny lobster in naturalistic flow conditions. *The*
1511 *Journal of Experimental Biology* **207**: 3785-3796.
- 1512 Huntingford F, Adams C (2005) Behavioural syndromes in farmed fish: implications for
1513 production and welfare. *Behaviour* **142**: 1207-1221.

1514 Ibarra-Zatarain Z, Fatsini E, Rey S, Chereguini O, Martin I, Rasines I, Alcaraz C, Duncan N
1515 (2016) Characterization of stress coping style in Senegalese sole (*Solea senegalensis*)
1516 juveniles and breeders for aquaculture. *The Royal Society Open Science* **3**: 160495.

1517 Jeffs A (2010) Status and challenges for advancing lobster aquaculture. *Journal of the Marine*
1518 *Biological Association of India* **52**: 320-326.

1519 Jescovitch LN, Ullman C, Rhodes M, Davis DA (2018) Effects of different feed
1520 management treatments on water quality for Pacific white shrimp *Litopenaeus*
1521 *vannamei*. *Aquaculture Research* **49**: 526-531.

1522 Jory DE (1995) Feed management practices for a healthy pond environment. In: Browdy CL,
1523 Hopkins JS (eds.) *Swimming Through Troubled Water, Proceedings of the Special*
1524 *Session on Shrimp Farming*, pp. 118-143. World Aquaculture Society, Louisiana, USA.

1525 Jory D, Cabrera T (2012) Marine shrimp. In: Lucas JS, Southgate PC (eds.) *Aquaculture:*
1526 *Farming Aquatic Animals and Plants*, pp. 476-513. Blackwell Publishing Ltd, Oxford,
1527 United Kingdom.

1528 Jory (2016) The proper feed management of commercial shrimp feeds; Part 2. Global
1529 Aquaculture Advocate [Cited 30 May 2018]. Available from URL:
1530 [https://www.aquaculturealliance.org/advocate/the-proper-management-of-commercial-](https://www.aquaculturealliance.org/advocate/the-proper-management-of-commercial-shrimp-feeds-part-2/)
1531 [shrimp-feeds-part-2/](https://www.aquaculturealliance.org/advocate/the-proper-management-of-commercial-shrimp-feeds-part-2/)

1532 Juinio-Menez MA, Gotanco RR (2004) Status of piny lobster resources of the Philippines. In:
1533 Williams KC (ed.) *Spiny Lobster Ecology and Exploitation in the South China Sea*
1534 *Region*, pp. 3-6. Australian Centre for International Agricultural Research, Canberra,
1535 Australia.

1536 Karavanich C, Atema J (1998) Olfactory recognition of urine signals in dominance fights
1537 between male lobster, *Homarus americanus*. *Behaviour* **135**: 719-730.

- 1538 Kautsky N, Rönnbäck P, Tedengren M, Troell M (2000) Ecosystem perspectives on
1539 management of disease in shrimp pond farming. *Aquaculture* **191**: 145-161.
- 1540 Kinne O (1966) Physiological aspects of animal life in estuaries with special reference to
1541 salinity. *Netherlands Journal of Sea Research* **3**: 222-244.
- 1542 Klages M, Muyakshin S, Soltwedel T, Arntz WE (2002) Mechanoreception, a possible
1543 mechanism for food fall detection in deep-sea scavengers. *Deep Sea Research Part I:
1544 Oceanographic Research Papers* **49**: 143-155.
- 1545 Koehl MAR (2006) The fluid mechanics of arthropod sniffing in turbulent odor
1546 plumes. *Chemical Senses* **31**: 93-105.
- 1547 Koehl MAR (2011) Hydrodynamics of sniffing by crustaceans. In: Breithaupt T, Thiel M
1548 (eds.) *Chemical Communication in Crustaceans*, pp. 85-102. Springer Science &
1549 Business Media, New York, USA.
- 1550 Krång AS, Rosenqvist G (2006) Effects of manganese on chemically induced food search
1551 behaviour of the Norway lobster, *Nephrops norvegicus* (L.) *Aquatic Toxicology* **78**:
1552 284-291.
- 1553 Kuhn DD, Smith SA, Boardman GD, Angier MW, Marsh L, Flick GJ (2010) Chronic toxicity
1554 of nitrate to Pacific white shrimp, *Litopenaeus vannamei*: impacts on survival, growth,
1555 antennae length, and pathology. *Aquaculture* **309**: 109-114.
- 1556 Kumar VS, Pandey PK, Anand T, Bhuvaneswari R, Kumar S (2017) Effect of periphyton
1557 (aquamat) on water quality, nitrogen budget, microbial ecology, and growth parameters
1558 of *Litopenaeus vannamei* in a semi-intensive culture system. *Aquaculture* **479**: 240-
1559 249.
- 1560 Kumlu M, Eroldogan OT, Saglamtimur B (2001) The effects of salinity and added substrates
1561 on growth and survival of *Metapenaeus monoceros* (Decapoda: Penaeidae) post-larvae.
1562 *Aquaculture* **196**: 177-188.

- 1563 Kurmaly K, Jones DA, Yule AB (1990) Acceptability and digestion of diets fed to larval
1564 stages of *Homarus gammarus* and the role of dietary conditioning behaviour. *Marine*
1565 *Biology* **106**: 181-190.
- 1566 Lawrence, AL, Lee PG (1997) Research in the Americas. In: D'Abramo LR, Conklin DE,
1567 Akiyama DM (eds.) *Crustacean Nutrition*, pp. 566-587. World Aquaculture Society,
1568 Los Angeles, USA.
- 1569 Lee PG, Meyers SP (1996) Chemoattraction and feeding stimulation in crustaceans.
1570 *Aquaculture Nutrition* **2**: 157-164.
- 1571 Lee PG, Meyers SP (1997) Chemoattraction and feeding stimulation. In: D'Abramo LR,
1572 Conklin DE, Akiyama DM (eds.) *Crustacean Nutrition*, pp. 292-352. World
1573 Aquaculture Society, Los Angeles, USA.
- 1574 Legrand JJ, Legrand-Hamelin E, Juchault P (1987) Sex determination in
1575 Crustacea. *Biological Reviews* **62**: 439-470.
- 1576 Li E, Chen L, Zeng C, Chen X, Yu N, Lai Q, Qin JG (2007) Growth, body composition,
1577 respiration and ambient ammonia nitrogen tolerance of the juvenile white shrimp,
1578 *Litopenaeus vannamei*, at different salinities. *Aquaculture* **265**: 385-390.
- 1579 Li E, Wang X, Chen K, Xu C, Qin JG, Chen L (2017) Physiological change and nutritional
1580 requirement of Pacific white shrimp *Litopenaeus vannamei* at low salinity. *Reviews in*
1581 *Aquaculture* **9**: 57-75.
- 1582 Liao IC, Chien Y (2011) The Pacific white shrimp, *Litopenaeus vannamei*, in Asia: the
1583 world's most widely cultured alien crustacean. In: Galil BS, Clark, PF, Carlton JT
1584 (eds.) *In the Wrong Place-Alien Marine Crustaceans: Distribution, Biology and*
1585 *Impacts*, pp. 489-519. Springer & Business media, Berlin, Germany.
- 1586 Lim C, Dominy W (1990) Evaluation of soybean meal as a replacement for marine animal
1587 protein in diets for shrimp (*Penaeus vannamei*). *Aquaculture* **87**: 53-63.

- 1588 Lin Y, Chen J (2001) Acute toxicity of ammonia on *Litopenaeus vannamei* Boone juveniles
1589 at different salinity levels. *Journal of Experimental Marine Biology & Ecology* **259**:
1590 109-119.
- 1591 Lin Y, Chen J (2003) Acute toxicity of nitrite on *Litopenaeus vannamei* (Boone) juveniles at
1592 different salinity levels. *Aquaculture* **224**: 193-201.
- 1593 Lin X, Pan J, Xu Z, Li Z, Li H (2008) Effect of periodic starvation on feeding, growth and
1594 defecation of *Litopenaeus vannamei*. *Acta Hydrobiologica Sinica* **32**: 403-407.
- 1595 Lima PP, Pontes CS, Arruda MF (2009) Activity pattern of the marine shrimp *Litopenaeus*
1596 *vannamei* (Boone 1931) in laboratory as a function of different feeding frequencies.
1597 *Aquaculture Research* **41**: 53-60.
- 1598 Liu C, Chen J (2004) Effect of ammonia on the immune response of white shrimp
1599 *Litopenaeus vannamei* and its susceptibility to *Vibrio alginolyticus*. *Fish & Shellfish*
1600 *Immunology* **16**: 321-334.
- 1601 Maciel JC, Francisco CJ, Miranda-Filho KC (2018) Compensatory growth and feed
1602 restriction in marine shrimp production, with emphasis on biofloc technology.
1603 *Aquaculture International* **26**: 203-212.
- 1604 Magallón-Barajas FJ, Servín-Villegas R, Portillo-Clark G, López-Moreno B (2006)
1605 *Litopenaeus vannamei* (Boone) post-larval survival related to age, temperature, pH and
1606 ammonium concentration. *Aquaculture Research* **37**: 492-499.
- 1607 Maicá PF, Borba MR, Martins TG, Wasielesky W (2014) Effect of salinity on performance
1608 and body composition of Pacific white shrimp juveniles reared in a super-intensive
1609 system. *Revista Brasileira de Zootecnia* **47**: 343-350.
- 1610 Martin M, Veran Y, Guelorget O, Pham D (1998) Shrimp rearing: stocking density, growth,
1611 impact on sediment, waste output and their relationships studied through the nitrogen
1612 budget in rearing ponds. *Aquaculture* **164**: 135-149.

1613 Martinez-Cordova LR, Porchas-Cornejo MA, Villarreal-Colemnares H, Calderon-Perez JA,
1614 Naranjo-Paramo J (1998) Evaluation of three feeding strategies on the culture of white
1615 shrimp *Penaeus vannamei* Boone 1931 in low water exchange ponds. *Aquacultural*
1616 *Engineering* **17**: 21-28.

1617 Martinez-Palacios CA, Ross LG, Jimenez-Valenzuela L (1996) The effects of temperature
1618 and body weight on the oxygen consumption of *Penaeus vannamei*, Boone,
1619 1931. *Journal of Aquaculture in the Tropics* **11**: 59-65.

1620 Martins CIM, Galhardo L, Noble C, Damsgård B, Spedicato MT, Zupa W, Beauchaud M,
1621 Kulczykowska E, Massabuau JC, Carter T, Planellas SR, Kristiansen T (2012)
1622 Behavioural indicators of welfare in farmed fish. *Fish Physiology & Biochemistry* **38**:
1623 17-41.

1624 Mauro NA, Malecha SR (1984) The effects of hypoxia on blood pH and lactate levels in
1625 *Macrobrachium rosenbergii* (De Man). *Comparative Biochemistry & Physiology Part*
1626 *A: Physiology* **77**: 627-630.

1627 Mead KS, Koehl MAR, O'Donnell MJ (1999) Stomatopod sniffing: the scaling of
1628 chemosensory sensillae and flicking behavior with body size. *Journal of Experimental*
1629 *Marine Biology & Ecology* **241**: 235-261.

1630 Metcalfe NB (1986) Intraspecific variation in competitive ability and food intake in
1631 salmonids: consequences for energy budgets and growth rates. *Journal of Fish*
1632 *Biology* **28**: 525-531.

1633 Miguel FF, Aréchiga H (1994) Circadian locomotor activity and its entrainment by food in
1634 the crayfish *Procambarus clarki*. *The Journal of Experimental Biology* **190**: 9-21.

1635 Mjos K, Grasso F, Atema J (1999) Antennule use by the American lobster, *Homarus*
1636 *americanus*, during chemo-orientation in three turbulent odor plumes. *The Biological*
1637 *Bulletin* **197**: 249-250.

1638 Moctezuma MA, Blake BF (1981) Burrowing activity in *Penaeus vannamei* Boone from the
1639 Caimanero-Huizache lagoon system on the Pacific coast of Mexico. *Bulletin of Marine*
1640 *Science* **31**: 312-317.

1641 Molina C, Cadena E, Orellana F (2000) Alimentación de camarones en relación a la actividad
1642 enzimática como una respuesta natural al ritmo circadiano y ciclo de muda. In: Cruz-
1643 Suárez LE, Ricque-Marie, D, Tapia-Salazar M, Olvera-Novoa MA, Civera-Cerecedo R
1644 (eds.) *Avances en Nutrición Acuícola V*, pp. 358-380. Memorias del V Simposium
1645 Internacional de Nutrición Acuícola, Universidad Autónoma de Nuevo León, Mérida,
1646 Yucatán, México.

1647 Molina-Poveda C, Escobar V, Gamboa-Delgado J, Cadena E, Orellana F, Piña R (2002)
1648 Estrategia de alimentación de acuerdo a la demanda fisiológica del juvenil *Litopenaeus*
1649 *vannamei* (Boone). In: Cruz-Suárez LE, Ricque-Marie D, Tapia-Salazar M, Gaxiola-
1650 Cortés MG, Simoes N (eds.). *Avances en Nutrición Acuícola VI*, pp. 98-113. Memorias
1651 del VI Simposium Internacional de Nutrición Acuícola, Cancún, Quintana Roo,
1652 México.

1653 Molina C, Espinoza M (2018) Demand feeding through automatic dispensing systems.
1654 Global Aquaculture Advocate [Cited 4 June 2018]. Available from URL:
1655 [https://www.aquaculturealliance.org/advocate/demand-feeding-through-automatic-](https://www.aquaculturealliance.org/advocate/demand-feeding-through-automatic-dispensing-systems/)
1656 [dispensing-systems/](https://www.aquaculturealliance.org/advocate/demand-feeding-through-automatic-dispensing-systems/).

1657 Montoya RA, Lawrence AL, Grant WE, Velasco M (2002) Simulation of inorganic nitrogen
1658 dynamics and shrimp survival in an intensive shrimp culture system. *Aquaculture*
1659 *Research* **33**: 81-94.

1660 Moore PA, Atema J, Gerhardt GA (1991) Fluid dynamics and microscale chemical
1661 movement in the chemosensory appendages of the lobster, *Homarus*
1662 *americanus*. *Chemical Senses* **16**: 663-674.

- 1663 Moore PA, Grills JL (1999) Chemical orientation to food by the crayfish *Orconectes rusticus*:
1664 influence of hydrodynamics. *Animal Behaviour* **58**: 953-963.
- 1665 Morgan DO, McMahon BR (1982) Acid tolerance and effects of sublethal acid exposure on
1666 iono-regulation and acid-base status in two crayfish *Procambarus clarki* and
1667 *Orconectes rusticus*. *The Journal of Experimental Biology* **97**: 241-252.
- 1668 Moss DR, Henning OL, Moss SM (2002) Sexual growth dimorphism in penaeid shrimp.
1669 Potential for all female culture?. *Global Aquaculture Advocate* **5**: 60-61.
- 1670 Moss KRK, Moss SM (2004) Effects of artificial substrate and stocking density on the
1671 nursery production of Pacific white shrimp *Litopenaeus vannamei*. *Journal of the*
1672 *World Aquaculture Society* **35**: 536-542.
- 1673 Moss DR, Moss SM (2006) Effects of gender and size on feed acquisition in the Pacific white
1674 shrimp *Litopenaeus vannamei*. *Journal of the World Aquaculture Society* **37**: 161-167.
- 1675 Nakamura K, Echavarria I (1989) Artificial controls of feeding rhythm of the prawn *Penaeus*
1676 *japonicus*. *Nippon Suisan Gakkaishi* **55**: 1325-1329.
- 1677 Nash, C. 2011. *The history of aquaculture*, 1st edn. Blackwell Publishing Ltd, Ames, Iowa,
1678 USA.
- 1679 Naylor RL, Hardy RW, Bureau DP, Chiu A, Elliott M, Farrell AP, Forster I, Gatlin DM,
1680 Goldberg RJ, Hua K, Nichols PD (2009) Feeding aquaculture in an era of finite
1681 resources. *Proceeding of the National Academy of Sciences of the United States of*
1682 *America* **106**: 15103-15110.
- 1683 Neiland AE, Soley N, Varley JB, Whitmarsh DJ (2001) Shrimp aquaculture: economic
1684 perspectives for policy development. *Marine Policy* **25**: 265-279.
- 1685 Nunes AJP, Goddard S, Gesteira TCV (1996) Feeding activity patterns of the Southern
1686 brown shrimp *Penaeus subtilis* under semi-intensive culture in NE Brazil. *Aquaculture*
1687 **144**: 371-386.

1688 Nunes AJP, Gesteira TCV, Goddard S (1997) Food ingestion and assimilation by the
1689 Southern brown shrimp *Penaeus subtilis* under semi-intensive culture in NE Brazil.
1690 *Aquaculture* **149**: 121-136.

1691 Nunes AJP, Parsons GJ (1999) Feeding levels of the Southern brown shrimp *Penaeus subtilis*
1692 in response to food dispersal. *Journal of the World Aquaculture Society* **30**: 331- 348.

1693 Nunes AJP, Sá MVC, Andriola-Neto FF, Lemos D (2006) Behavioural response to selected
1694 feed attractants and stimulants in Pacific white shrimp, *Litopenaeus vannamei*.
1695 *Aquaculture* **260**: 244-254.

1696 Palacios E, Bonilla A, Luna D, Racotta IS (2004) Survival, Na⁺/K⁺ -ATPase and lipid
1697 responses to salinity challenge in fed and starved white Pacific shrimp (*Litopenaeus*
1698 *vannamei*) postlarvae. *Aquaculture* **234**: 497-511.

1699 Pearson WH, Sugarman PC, Woodruff DL (1979) Thresholds for detection and feeding
1700 behavior in the Dungeness crab, *Cancer magister* (Dana). *Journal of Experimental*
1701 *Marine Biology & Ecology* **39**: 65-78.

1702 Peñaflorida VD, Virtanen E (1996) Growth, survival and feed conversion of juvenile shrimp
1703 (*Penaeus monodon*) fed a betaine/amino acid additive. *The Israeli Journal of*
1704 *Aquaculture - Bamidgeh* **48**: 3-9.

1705 Pérez-Rostro CI, Ramirez JL, Ibarra AM (1999) Maternal and cage effects on genetic
1706 parameter estimation for Pacific white shrimp *Penaeus vannamei* Boone. *Aquaculture*
1707 *Research* **30**: 681-693.

1708 Pérez-Rostro CI, Ibarra AM (2003) Heritabilities and genetic correlation of size traits at
1709 harvest size in sexually dimorphic Pacific white shrimp (*Litopenaeus vannamei*) grown
1710 in two environments. *Aquaculture Research* **34**: 1079-1085.

1711 Pérez-Rostro CI, Racotta IS, Ibarra AM (2004) Decreased genetic variation in metabolic
1712 variables of *Litopenaeus vannamei* shrimp after exposure to acute hypoxia. *Journal of*
1713 *Experimental Marine Biology & Ecology* **302**: 189-200.

1714 Pittet AO, Ellis JC, Lee PG (1996) Methodology for the identification and quantitative
1715 measurement of chemical stimulants for penaeid shrimp. *Aquaculture Nutrition* **2**: 175-
1716 182.

1717 Ponce-Palafox JT, Martínez-Palacios CA, Ross LG (1997) The effects of salinity and
1718 temperature on the growth and survival rates of juvenile white shrimp, *Penaeus*
1719 *vannamei*, Boone, 1931. *Aquaculture* **157**: 107-115.

1720 Ponce-Palafox JT, Ruíz-Luna A, García-Ulloa M, Esparza-Leal HM, Arredondo-Figueroa JL,
1721 Martínez-Palacios CA, Ross LG (2013) A response-surface analysis of the relative
1722 importance of the temperature, salinity and body weight on the respiratory metabolism
1723 of the white shrimp *Litopenaeus vannamei* (Boone, 1931). *Marine & Freshwater*
1724 *Behaviour & Physiology* **46**: 399-417.

1725 Pontes CS, Arruda, MF (2005a) Comportamento de *Litopenaeus vannamei* (Boone)
1726 (Crustacea, Decapoda, Penaeidae) em função da oferta do alimento artificial nas fases
1727 clara e escura do período de 24 horas. *Revista Brasileira de Zoologia* **22**: 648-652.

1728 Pontes CS, Arruda MF (2005b) Acesso ao alimento artificial e enchimento do trato digestivo
1729 de juvenis do camarão marinho *Litopenaeus vannamei* (Boone) (Crustacea, Decapoda,
1730 Penaeidae) durante as fases clara e escuro do período de 24 horas. *Revista Brasileira de*
1731 *Zoologia* **22**: 1039-1043.

1732 Pontes CS (2006) Padrão de deslocamento do camarão marinho *Litopenaeus vannamei*
1733 (Boone) (Crustacea, Decapoda, Penaeidae) nas fases clara e escura ao longo de 24
1734 horas. *Revista Brasileira de Zoologia* **23**: 223-227.

- 1735 Pontes CS, Arruda MF, Menezes AA, Lima PP (2006) Daily activity pattern of the marine
1736 shrimp *Litopenaeus vannamei* (Boone 1931) juveniles under laboratory conditions.
1737 *Aquaculture Research* **37**: 1001-1006.
- 1738 Pontes CS, Lima PP, Arruda MF (2008) Feeding responses of juvenile shrimp *Litopenaeus*
1739 *vannamei* (Boone) fed at different frequencies under laboratory conditions. *Aquaculture*
1740 *Research* **39**: 1416-1422.
- 1741 Pörtner HO, Langenbuch M, Reipschläger A (2004) Biological impact of elevated ocean CO₂
1742 concentrations: lessons from animal physiology and earth history. *Journal of*
1743 *Oceanography* **60**: 705-718.
- 1744 Price RB, Ache BW (1977) Peripheral modification of chemosensory information in the
1745 spiny lobster. *Comparative Biochemistry & Physiology Part A: Physiology* **57**: 249-
1746 253.
- 1747 Primavera JH, Leбата J (1995) Diel activity patterns in *Metapenaeus* and *Penaeus* juveniles.
1748 *Hydrobiologia* **295**: 295-302.
- 1749 Ravi R, Manisseri MK (2013) The effect of different pH and photoperiod regimens on the
1750 survival rate and developmental period of the larvae of *Portunus pelagicus* (Decapoda,
1751 Brachyura, Portunidae). *Iranian Journal of Fisheries Sciences* **12**: 490-499.
- 1752 Reeder PB, Ache BW (1980) Chemotaxis in the Florida spiny lobster, *Panulirus*
1753 *argus*. *Animal Behaviour* **28**: 831-839.
- 1754 Reeves RB, Rahn H (1979) Patterns in vertebrate acid-base regulation. In: Wood SC, Lenfant
1755 C (eds.) *Evolution of Respiratory Processes, a Comparative Approach*, pp. 225-252.
1756 Marcel Dekker, New York, USA.
- 1757 Rimmer MA, Sugama K, Rakhmawati D, Rofiq R, Habgood RH (2013) A review and SWOT
1758 analysis of aquaculture development in Indonesia. *Reviews in Aquaculture* **5**: 255-279.

- 1759 Rittschof D (1992) Chemosensation in the daily life of crabs. *American Zoologist* **32**: 363-
1760 369.
- 1761 Robertson JR, Fudge JA, Vermeer GK (1981) Chemical and live feeding stimulants of the
1762 sand fiddler crab, *Uca pugilator* (Bosc). *Journal of Experimental Marine Biology &*
1763 *Ecology* **53**: 47-64.
- 1764 Robertson L, Bray W, Leung-Trujillo J, Lawrence A (1987) Practical molt staging of
1765 *Penaeus setiferus* and *Penaeus stylirostris*. *Journal of the World Aquaculture Society*
1766 **18**: 180-185.
- 1767 Robertson L, Lawrence AL, Castille FL (1993) Effect of feeding frequency and feeding time
1768 on growth of *Penaeus vannamei* (Boone). *Aquaculture & Fisheries Management* **24**: 1-
1769 6.
- 1770 Rodríguez J, Bayot B, Amano Y, Panchana F, De Blas I, Alday V, Calderón J (2003) White
1771 spot syndrome virus infection in cultured *Penaeus vannamei* (Boone) in Ecuador with
1772 emphasis on histopathology and ultrastructure. *Journal of Fish Diseases* **26**: 439-450.
- 1773 Rola WR, Hasan MR (2007) *Economics of Aquaculture Feeding Practices: A Synthesis of*
1774 *Case Studies Undertaken in Six Asian Countries*. FAO, Rome, Italy; Report No.: 505.
- 1775 Rosas C, Martinez E, Gaxiola G, Brito R, Sánchez A, Soto LA (1999) The effect of dissolved
1776 oxygen and salinity on oxygen consumption, ammonia excretion and osmotic pressure
1777 of *Penaeus setiferus* (Linnaeus) juveniles. *Journal of Experimental Marine Biology &*
1778 *Ecology* **234**: 41-57.
- 1779 Rosas C, López N, Mercado P, Martínez E (2001) Effect of salinity acclimation on oxygen
1780 consumption of juveniles of the white shrimp *Litopenaeus vannamei*. *Journal of*
1781 *Crustacean Biology* **21**: 912-922.
- 1782 Rosenberry B (2001) *World shrimp farming 2001*. Shrimp News International, San Diego,
1783 California, USA.

1784 Rusaini R, Owens L (2011) A simple technique to stage the moult of *Penaeus*
1785 *monodon*. *Asian Fisheries Science* **24**: 1-11.

1786 Samocha TM, Davis DA, Saoud IP, DeBault K (2004a) Substitution of fish meal by co-
1787 extruded soybean poultry by-product meal in practical diets for the Pacific white
1788 shrimp, *Litopenaeus vannamei*. *Aquaculture* **231**: 197-203.

1789 Samocha TM, Lawrence AL, Collins CA, Castille FL, Bray WA, Davies CJ, Lee PG, Wood
1790 GF (2004b) Production of the Pacific white shrimp, *Litopenaeus vannamei*, in high-
1791 density greenhouse-enclosed raceways using low salinity groundwater. *Journal of*
1792 *Applied Aquaculture* **15**: 1-19.

1793 Sanchez DR, Fox JM, Lawrence AL, Castille FL, Dunsford B (2005) A methodology for
1794 evaluation of dietary feeding stimulants for the Pacific white shrimp, *Litopenaeus*
1795 *vannamei*. *Journal of the World Aquaculture Society* **36**:14-23.

1796 Sandifer PA, Hopkins JS, Stokes AD (1987) Intensive culture potential of *Penaeus vannamei*.
1797 *Journal of the World Aquaculture Society* **18**: 94-100.

1798 Sandifer PA, Hopkins JS, Stokes AD (1988) Intensification of shrimp culture in earthen
1799 ponds in South Carolina: progress and prospects. *Journal of the World Aquaculture*
1800 *Society* **19**: 218-226.

1801 Santos ADA, López-Olmeda JF, Sánchez-Vázquez FJ, Fortes-Silva R (2016)
1802 Synchronization to light and mealtime of the circadian rhythms of self-feeding behavior
1803 and locomotor activity of white shrimps (*Litopenaeus vannamei*). *Comparative*
1804 *Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **199**: 54-61.

1805 Sanudin N, Tuzan AD, Yong ASK (2014) Feeding activity and growth performance of
1806 shrimp post larvae *Litopenaeus vannamei* under light and dark conditions. *Journal of*
1807 *Agricultural Science* **6**: 103-109.

- 1808 Schmidt M, Van Ekeris L, Ache BW (1992) Antennular projections to the midbrain of the
1809 spiny lobster. I. Sensory innervation of the lateral and medial antennular neuropils. *The*
1810 *Journal of Comparative Neurology* **318**: 277-290.
- 1811 Schmidt M, Derby CD (2005) Non-olfactory chemoreceptors in asymmetric setae activate
1812 antennular grooming behavior in the Caribbean spiny lobster *Panulirus argus*. *The*
1813 *Journal of Experimental Biology* **208**: 233-248.
- 1814 Schmitt BC, Ache BW (1979) Olfaction: responses of a decapod crustacean are enhanced by
1815 flicking. *Science* **205**: 204-206.
- 1816 Schuler DJ, Boardman GD, Kuhn DD, Flick GJ (2010) Acute toxicity of ammonia and nitrite
1817 to Pacific white shrimp, *Litopenaeus vannamei*, at low salinities. *Journal of the World*
1818 *Aquaculture Society* **41**: 438-446.
- 1819 Seidman ER, Lawrence AL (1985) Growth, feed digestibility, and proximate body
1820 composition of juvenile *Penaeus vannamei* and *Penaeus monodon* grown at different
1821 dissolved oxygen levels. *Journal of the World Mariculture Society* **16**: 333-346.
- 1822 Seiffert WQ, Andreatta ER (2004) El manejo de la alimentación y la sostenibilidad en el
1823 cultivo de camarones en el brasil. In: Cruz Suárez LE, Ricque-Marie D, Nieto-López
1824 MG, Villarreal D, Scholz U, González M (eds.) *Avances en Nutrición Acuícola VII*, pp.
1825 707-720. Memorias del VII Simposium Internacional de Nutrición Acuícola,
1826 Hermosillo, Sonora, México.
- 1827 Shuranova Z, Burmistrov Y, Abramson CI (2005) Habituation to a novel environment in the
1828 crayfish *Procambarus cubensis*. *Journal of Crustacean Biology* **25**: 488-494.
- 1829 Sick LV, White D, Baptist G (1973) The effect of duration of feeding, amount of food, light
1830 intensity, and animal size on rate of ingestion of pelleted food by juvenile penaeid
1831 shrimp. *The Progressive Fish-Culturist* **35**: 22-26.

- 1832 Sih A, Bell A, Johnson JC (2004) Behavioral syndromes: an ecological and evolutionary
1833 overview. *Trends in Ecology & Evolution* **19**: 372-378.
- 1834 Silva PF, Medeiros MS, Silva HPA, Arruda MF (2012) A study of feeding in the shrimp
1835 *Farfantepenaeus subtilis* indicates the value of species level behavioral data for
1836 optimizing culture management. *Marine & Freshwater Behaviour & Physiology* **45**:
1837 121-134.
- 1838 Smith DM, Burford MA, Tabrett SJ, Irvin SJ, Ward L (2002) The effect of feeding frequency
1839 on water quality and growth of the black tiger shrimp (*Penaeus monodon*). *Aquaculture*
1840 **207**: 125-136.
- 1841 Smith DM, Tabrett SJ, Barclay MC, Irvin SJ (2005) The efficacy of ingredients included in
1842 shrimp feeds to stimulate intake. *Aquaculture Nutrition* **11**: 263-272.
- 1843 Smith DV, Tabrett SJ (2013) The use of passive acoustics to measure feed consumption by
1844 *Penaeus monodon* (giant tiger prawn) in cultured systems. *Aquacultural Engineering*
1845 **57**: 38-47.
- 1846 Snow PJ (1973) The antennular activities of the hermit crab, *Pagurus alaskensis*
1847 (Benedict). *The Journal of Experimental Biology* **58**: 745-765.
- 1848 Snow PJ (1975) Central patterning and reflex control of antennular flicking in the hermit crab
1849 *Pagurus alaskensis* (Benedict). *The Journal of Experimental Biology* **63**: 17-32.
- 1850 Soares R, Peixoto S, Wasielesky W, D'Incao F (2005) Feeding rhythms and diet of
1851 *Farfantepenaeus paulensis* under pen culture in Patos Lagoon estuary, Brazil. *Journal*
1852 *of Experimental Marine Biology & Ecology* **322**: 167-176.
- 1853 Sookying D, Davis DA, Da Silva FSD (2013) A review of the development and application
1854 of soybean-based diets for Pacific white shrimp *Litopenaeus vannamei*. *Aquaculture*
1855 *Nutrition* **19**: 441-448.

1856 Spicer JI, Raffo A, Widdicombe S (2007) Influence of CO₂-related seawater acidification on
1857 extracellular acid-base balance in the velvet swimming crab *Necora puber*. *Marine*
1858 *Biology* **151**: 1117-1125.

1859 Staples DJ, Heales DS (1991) Temperature and salinity optima for growth and survival of
1860 juvenile banana prawns *Penaeus merguensis*. *Journal of Experimental Marine Biology*
1861 *& Ecology* **154**: 251-274.

1862 Steiner JE, Harpaz S (1987) Behavior stereotypes of food acceptance and of the rejection of
1863 'bitter' food in the freshwater prawn *Macrobrachium rosenbergii*. *Chemical Senses* **12**:
1864 89-97.

1865 Steullet P, Flavus T, Radman D, Hamidani G, Zhou M, Dudar O, Hill R, Derby CD (1999)
1866 The aesthetasc-olfactory lobe pathway of spiny lobsters is not necessary for odor-
1867 activated searching behavior, odor-associative learning, and discrimination of complex
1868 odors. *Chemical Senses* **24**: 613 (abstract).

1869 Steullet P, Dudar O, Flavus T, Zhou M, Derby CD (2001) Selective ablation of antennular
1870 sensilla on the Caribbean spiny lobster *Panulirus argus* suggest that dual antennular
1871 chemosensory pathways mediate odorant activation of searching and localization of
1872 food. *The Journal of Experimental Biology* **204**: 4259-4269.

1873 Steullet P, Krutzfeldt DR, Hamidani G, Flavus T, Ngo V, Derby CD (2002) Dual antennular
1874 chemosensory pathways mediate odor-associative learning and odor discrimination in
1875 the Caribbean spiny lobster *Panulirus argus*. *The Journal of Experimental Biology* **205**:
1876 851-867.

1877 Stuck KC, Watts SA, Wang SY (1996) Biochemical responses during starvation and
1878 subsequent recovery in postlarval Pacific white shrimp, *Penaeus vannamei*. *Marine*
1879 *Biology* **125**: 33-45.

- 1880 Sun M, Hassan SG, Li D (2016) Models for estimating feed intake in aquaculture: a review.
1881 *Computers & Electronics in Aquaculture* **127**: 425-438.
- 1882 Suresh AV, Kumaraguru-Vasagam KP, Nates S (2011) Attractability and palatability of
1883 protein ingredients of aquatic and terrestrial animal origin, and their practical value for
1884 blue shrimp, *Litopenaeus stylirostris* fed diets formulated with high levels of poultry
1885 byproduct meal. *Aquaculture* **319**: 132-140.
- 1886 Tacon AGJ (1996) Nutritional studies in crustaceans and the problems of applying research
1887 findings to practical farming systems. *Aquaculture Nutrition* **1**: 165-174.
- 1888 Tacon AGJ, Barg UC (1998) Major challenges to feed development for marine and
1889 diadromous finfish and crustacean species. In: De Silva SS (ed.) *Tropical Mariculture*,
1890 pp. 171-208. Academic Press, San Diego, California, USA.
- 1891 Tacon AGJ, Cody JJ, Conquest LD, Divakaran S, Forster IP, Decamp OE (2002) Effect of
1892 culture system on the nutrition and growth performance of Pacific white shrimp
1893 *Litopenaeus vannamei* (Boone) fed different diets. *Aquaculture Nutrition* **8**: 121-137.
- 1894 Tacon AGJ, Jory D, Nunes A (2013) Shrimp feed management: issues and perspectives. In:
1895 Hasan MR and New MB (eds.) *On-Farm Feeding and Feed Management in*
1896 *Aquaculture*, pp. 481-488. FAO Fisheries and Aquaculture Technical Paper No. 583,
1897 Rome, FAO.
- 1898 Taylor J, Vinatea L, Ozorio R, Schuweitzer R, Andreatta ER (2004) Minimizing the effects
1899 of stress during eyestalk ablation of *Litopenaeus vannamei* females with topical
1900 anaesthetic and a coagulating agent. *Aquaculture* **233**: 173-179.
- 1901 Thiel M, Breithaupt T (2011) Chemical communication in crustaceans: research challenges
1902 for the twenty-first century. In: Breithaupt T, Thiel M (eds.) *Chemical Communication*
1903 *in Crustaceans*, pp. 3-22. Springer Science & Business Media, New York, USA.

- 1904 Thompson FL, Abreu PC, Wasielesky W (2002) Importance of biofilm for water quality and
1905 nourishment in intensive shrimp culture. *Aquaculture* **203**: 263-278.
- 1906 Thuy NTB, Ngoc NB (2004) Current status and exploitation of wild spiny lobsters in
1907 Vietnamese waters. In: Williams KC (ed.) *Spiny lobster ecology and exploitation in the*
1908 *south China sea region*, pp. 13-16. Australian Centre for International Agricultural
1909 Research, Canberra, Australia.
- 1910 Tierney AJ, Atema J (1988) Amino acid chemoreception: effects of pH on receptors and
1911 stimuli. *Journal of Chemical Ecology* **14**: 135-141.
- 1912 Vainikka A, Rantala MJ, Niemelä P, Hirvonen H, Kortet R (2011) Boldness as a consistent
1913 personality trait in the noble crayfish, *Astacus astacus*. *Acta Ethologica* **14**: 17-25.
- 1914 Valenzuela-Quiñónez W, Rodríguez-Quiroz G, Ponce-Palafox JT, Esparza-Leal HM (2011)
1915 Efecto de diferentes combinaciones de temperatura y salinidad sobre el consumo
1916 específico de oxígeno en el camarón blanco *Litopenaeus vannamei*. *Revista de Biología*
1917 *Marina y Oceanografía* **46**: 303-311.
- 1918 Vargas-Albores F, Ochoa JL (1992) Variation of pH, osmolality, sodium and potassium
1919 concentrations in the haemolymph of sub-adult blue shrimp (*Penaeus stylirostris*)
1920 according to size. *Comparative Biochemistry and Physiology Part A: Physiology* **102**:
1921 1-5.
- 1922 Vega-Villasante F, Nolasco-Soria H, Civera-Cerecedo R, González-Valdés R, Oliva-Suárez
1923 M (2000) Alternativa para la alimentación del camarón en cultivo: el manejo de la
1924 muda. In: Cruz-Suárez L, Ricque-Marie D, Tapia-Salazar M, Olvera-Novoa M, Civera-
1925 Cercedo R (eds.) *Avances en Nutrición Acuícola V*, pp. 313-320. Memorias del V
1926 Simposium Internacional en Nutrición Acuícola, Mérida, Yucatán, México.

- 1927 Velasco M, Lawrence AL, Castille FL (1999) Effect of variations in daily feeding frequency
1928 and ration size on growth of shrimp, *Litopenaeus vannamei* (Boone), in zero-water
1929 exchange culture tanks. *Aquaculture* **179**: 141-148.
- 1930 Vijayakumaran M, Venkatesan R, Murugan TS, Kumar TS, Jha DK, Remany MC, Thilakam
1931 JML, Jahan SS, Dharani G, Kathirolu S, Selvan K (2009) Farming of spiny lobsters in
1932 sea cages in India. *New Zealand Journal of Marine & Freshwater Research* **43**: 623-
1933 634.
- 1934 Villarreal H, Hinojosa P, Naranjo J (1994) Effect of temperature and salinity on the oxygen
1935 consumption of laboratory produced *Penaeus vannamei* postlarvae. *Comparative*
1936 *Biochemistry & Physiology Part A: Physiology* **108**: 331-336.
- 1937 Vinatea L, Olivera-Gálvez A, Venero J, Leffler J, Browdy C (2009) Oxygen consumption of
1938 *Litopenaeus vannamei* juveniles in heterotrophic medium with zero water exchange.
1939 *Pesquisa Agropecuária Brasileira* **44**: 534-538.
- 1940 Wajsbrodt N, Gasith A, Krom MD, Samocha TM (1990) Effect of dissolved oxygen and the
1941 molt stage on the acute toxicity of ammonia to juvenile green tiger prawn *Penaeus*
1942 *semisulcatus*. *Environmental Toxicology & Chemistry* **9**: 497-504.
- 1943 Wang W, Wang A, Chen L, Liu Y, Sun R (2002) Effects of pH on survival, phosphorus
1944 concentration, adenylate energy charge and Na⁺ - K⁺ ATPase activities of *Penaeus*
1945 *chinensis* Osbeck juveniles. *Aquatic Toxicology* **60**: 75-83.
- 1946 Wang W, Zhou J, Wang P, Tian T, Zheng Y, Liu Y, Mai W, Wang A (2009) Oxidative stress,
1947 DNA damage and antioxidant enzyme gene expression in the Pacific white shrimp,
1948 *Litopenaeus vannamei* when exposed to acute pH stress. *Comparative Biochemistry &*
1949 *Physiology Part C: Toxicology & Pharmacology* **150**: 428-435.

- 1950 Wassenberg TJ, Hill BJ (1994) Laboratory study of the effect of light on the emergence
1951 behaviour of eight species of commercially important adult penaeid prawns. *Australian*
1952 *Journal of Marine & Freshwater Research* **45**: 43-50.
- 1953 Watanabe T (2002) Strategies for further development of aquatic feeds. *Fisheries Science* **68**:
1954 242-252.
- 1955 Wickins JF (1976) The tolerance of warm-water prawns to recirculated water. *Aquaculture* **9**:
1956 19-37.
- 1957 Wilcox JR, Jeffries HP (1976) Hydration in the sand shrimp *Crangon septemspinosa*: relation
1958 to diet. *The Biological Bulletin* **150**: 522-530.
- 1959 Wight K, Francis L, Eldridge D (1990) Food aversion learning by the hermit crab *Pagurus*
1960 *granosimanus*. *The Biological Bulletin* **178**: 205-209.
- 1961 Wilkens LA, Schmitz B, Herrnkind WF (1996) Antennal responses to hydrodynamic and
1962 tactile stimuli in the spiny lobster *Panulirus argus*. *The Biological Bulletin* **191**: 187-
1963 198.
- 1964 Wilson DS, Coleman K, Clark AB, Biederman L (1993) Shy-bold continuum in pumpkinseed
1965 sunfish (*Lepomis gibbosus*): an ecological study of a psychological trait. *Journal of*
1966 *Comparative Psychology* **107**: 250-260.
- 1967 Wilson DS (1998) Adaptive individual differences within single populations. *Philosophical*
1968 *Transactions of the Royal Society of London B: Biological Sciences* **353**: 199-205.
- 1969 Wroblewska J, Whalley S, Fischetti M, Daniel PC (2002) Identification of chemosensory
1970 sensilla activating antennular grooming behaviour in the Caribbean spiny lobster,
1971 *Panulirus argus*. *Chemical Senses* **27**: 769-778.
- 1972 Wu L, Dong S (2002) The effects of repetitive 'starvation-and-refeeding' cycles on the
1973 compensatory growth response in Chinese shrimp, *Fenneropenaeus chinensis* (Osbeck,
1974 1765) (Decapoda, Penaeidae). *Crustaceana* **74**: 1225-1239.

- 1975 Wyban JA, Lee CS, Sato VT, Sweeney JN, Richards WK (1987) Effect of stocking density
1976 on shrimp growth rates in manure-fertilized ponds. *Aquaculture* **61**: 23-32.
- 1977 Wyban JA, Sweeney JN, Kanna RA (1988) Shrimp yields and economic potential of
1978 intensive round pond systems. *Journal of the World Aquaculture Society* **19**: 210-217.
- 1979 Wyban JA, Sweeney JN (1991) *The Oceanic Institute Shrimp Manual: Intensive Shrimp*
1980 *Production Technology*. The Oceanic Institute, Makapuu Point, Honolulu, Hawaii,
1981 USA.
- 1982 Wyban J, Walsh WA, Godin DM (1995) Temperature effects on growth, feeding rate and
1983 feed conversion of the Pacific white shrimp (*Penaeus vannamei*). *Aquaculture* **138**:
1984 267-279.
- 1985 Yi D, Reardon T, Stringer R (2016) Shrimp aquaculture technology change in Indonesia: are
1986 small farmers included?. *Aquaculture* **493**: 436-445.
- 1987 You K, Yang H, Liu Y, Liu S, Zhou Y, Zhang T (2006) Effects of different light sources and
1988 illumination methods on growth and body color of shrimp *Litopenaeus vannamei*.
1989 *Aquaculture* **252**: 557-565.
- 1990 Zarain-Herzberg M, Campa-Córdova AI, Cavalli RO (2006) Biological viability of producing
1991 white shrimp *Litopenaeus vannamei* in seawater floating cages. *Aquaculture* **259**: 283-
1992 289.
- 1993 Zhang P, Zhang X, Li J, Huang G (2006) The effects of body weight, temperature, salinity,
1994 pH, light intensity and feeding condition on lethal DO levels of whiteleg shrimp,
1995 *Litopenaeus vannamei* (Boone, 1931). *Aquaculture* **256**: 579-587.
- 1996 Zhang P, Zhang X, Li J, Gao T (2009) Effect of salinity on survival, growth, oxygen
1997 consumption and ammonia-N excretion of juvenile whiteleg shrimp, *Litopenaeus*
1998 *vannamei*. *Aquaculture Research* **40**: 1419-1427.

- 1999 Zhang B, Lin W, Huang J, Wang Y, Xu R (2010) Effects of artificial substrates on the
2000 growth, survival and spatial distribution of *Litopenaeus vannamei* in the intensive
2001 culture condition. *Iranian Journal of Fisheries Sciences* **9**: 293-304.
- 2002 Zhou C, Xu D, Lin K, Sun C, Yang X (2017) Intelligent feeding control methods in
2003 aquaculture with an emphasis on fish: a review. *Reviews in Aquaculture* **0**: 1-19
- 2004 Zimmer-Faust RK, Case JF (1983) A proposed dual role of odor in foraging by the California
2005 spiny lobster, *Panulirus interruptus* (Randall). *The Biological Bulletin* **164**: 341-353.
- 2006 Zimmer-Faust RK, Michel WC, Tyre JE, Case JF (1984) Chemical induction of feeding in
2007 California spiny lobster, *Panulirus interruptus* (Randall): responses to molecular
2008 weight fractions of abalone. *Journal of Chemical Ecology* **10**: 957-971.
- 2009 Zimmer-Faust RK (1987) Crustacean chemical perception: towards a theory on optimal
2010 chemoreception. *The Biological Bulletin* **172**:10-29.
- 2011 Zimmer-Faust RK (1989) The relationship between chemoreception and foraging behavior in
2012 crustaceans. *Limnology & Oceanography* **34**: 1367-1374.
- 2013 Zimmer-Faust RK (1991) Chemical signal-to-noise detection by spiny lobsters. *The*
2014 *Biological Bulletin* **181**: 419-426.
- 2015 Zion B, Karplus I, Barki A (2010) Generalization and discrimination of positive and negative
2016 acoustic stimuli in the common carp (*Cyprinus carpio*). *Behavioural Processes* **83**:
2017 306–310.
- 2018 Zion B, Barki A, Grinshpon J, Rosenfeld L, Karplus I (2011a) Retention of acoustic
2019 conditioning in St. Peter's fish *Sarotherodon galilaeus*. *Journal of Fish Biology* **78**:
2020 838–847.
- 2021 Zion B, Karplus I, Grinshpon J, Rosenfeld L, Barki A (2011b) Periodic reinforcement of
2022 acoustically conditioned behaviour in St. Peter's fish, *Sarotherodon galilaeus*, for
2023 ranching purposes. *Aquaculture* **315**: 394–399.