



## UWS Academic Portal

### 99mTc nanocolloid indocyanine green

Gambini, Juan; Silvera, Enzo; Musetti, Matias; Quinn, Thomas; Yang, Guang Zhong; Matalonga, Santiago; Hermida, Juan; Alonso, Omar; Cabral, Pablo

*Published in:*  
The Journal of Nuclear Medicine

Published: 01/05/2019

*Document Version*  
Peer reviewed version

[Link to publication on the UWS Academic Portal](#)

*Citation for published version (APA):*

Gambini, J., Silvera, E., Musetti, M., Quinn, T., Yang, G. Z., Matalonga, S., Hermida, J., Alonso, O., & Cabral, P. (2019). 99mTc nanocolloid indocyanine green: an hybrid tracer for breast sentinel node procedures. *The Journal of Nuclear Medicine*, 60(Supplement 1), [1231].

[http://jnm.snmjournals.org/content/60/supplement\\_1/1231.abstract?sid=da0a99ee-dbc8-4c6b-8751-14bca2bbdd6a](http://jnm.snmjournals.org/content/60/supplement_1/1231.abstract?sid=da0a99ee-dbc8-4c6b-8751-14bca2bbdd6a)

#### General rights

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

#### Take down policy

If you believe that this document breaches copyright please contact [pure@uws.ac.uk](mailto:pure@uws.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

**Introduction:** Breast sentinel lymph node biopsy (SLNB) evaluates if cancer has spread outside the breast, by allowing the identification of the node that has the highest probability of harboring metastasis. Once it is identified and removed during surgery, histologic examination looks for the presence of metastasis. If metastasis are identified, the patient will have an axillary dissection performed, if no metastasis are identified, the patient can spare the axillary dissection with its associated morbidity. SLNB procedures are performed with radiotracers or with dyes, or a combination of both. Recently, the hybrid tracer (fluorescent-radioactive)  $^{99m}\text{Tc}$  nanocolloid indocyanine green ( $^{99m}\text{Tc}$  NC ICG) has been described. The aim of our work is to describe our experience with the hybrid tracer in breast SLN biopsy. Materials Patients were injected periareolarly subcutaneous in the same quadrant of the lesions. Planar Scintigraphic and SPECT/CT (Mediso Anyscan) images were later acquired. During the SLNB a gamma probe (Europrobe) and our own developed near infrared (NIR) visualization system was used to localize the SLN. This combination allowed surgeons to guide them to the SLN using the acoustic cue provided from the gamma probe and the images from the NIR visualization system. Once the nodes were found they were removed and examined histopathologically. Patients who had positive SLNB underwent axillary dissection.

**Results:** SLNB with  $^{99m}\text{Tc}$  NC ICG allowed us to use their complementary fluorescent and radioactive properties in order to find the SLN. In this way, the gamma probe guided us to the region where the node was, and fluorescence made it easy to remove it and spare the rest of surrounding tissue. In this way we could remove 38 nodes, being 4 positive. All radioactive nodes were fluorescent.

**Conclusions:** SLNB with  $^{99m}\text{Tc}$  NC ICG could be performed without complications, having no adverse effects on patients. The hybrid tracer adds a visual cue with to the procedure aiding surgeons on SLN localization and removal.