



UWS Academic Portal

Gene expression during THP-1 differentiation is influenced by vitamin D3 and not vibrational mechanostimulation

Simakou, Theodoros; Freeburn, Robin; Henriquez, Fiona L.

Published in:
PeerJ

DOI:
[10.7717/peerj.11773](https://doi.org/10.7717/peerj.11773)

Published: 14/07/2021

Document Version
Publisher's PDF, also known as Version of record

[Link to publication on the UWS Academic Portal](#)

Citation for published version (APA):

Simakou, T., Freeburn, R., & Henriquez, F. L. (2021). Gene expression during THP-1 differentiation is influenced by vitamin D3 and not vibrational mechanostimulation. *PeerJ*, 9, [e11773]. <https://doi.org/10.7717/peerj.11773>

General rights

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact pure@uws.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Gene expression during THP-1 differentiation is influenced by vitamin D3 and not vibrational mechanostimulation

Theodoros Simakou^{1,2}, Robin Freeburn¹ and Fiona L. Henriquez¹

¹School of Health and Life Sciences, University of West of Scotland, Paisley, United Kingdom

²Institute of Infection, Immunity and Inflammation, University of Glasgow, Glasgow, United Kingdom

ABSTRACT

Background. In injury or infection, monocytes migrate into the affected tissues from circulation and differentiate into macrophages which are subsequently involved in the inflammatory responses. Macrophage differentiation and activation have been studied in response to multiple chemokines and cytokines. However, mechanical, and physical stimuli can also influence macrophage differentiation, activation, cytokine production, and phagocytic activity.

Methods. In this study the macrophage differentiation from THP-1 monocytes was assessed upon the stimulation with 1,25-dihydroxyvitamin D3 and 1,000 Hz vibrations, using qPCR for quantification of transcript expression. Vitamin D binds the vitamin D receptor (VDR) and subsequently modulates the expression of a variety of genes in monocytes. The effects of the 1,000 Hz vibrational stimulation, and the combined treatment of vitamin D3 and 1000 Hz vibrations were unknown. The differentiation of macrophages was assessed by looking at transcription of macrophage markers (e.g., *CD14*, *CD36*), antigen presenting molecules (e.g., *HLA-DRA*), transcription factors (e.g., *LEF-1*, *TCF7L2*), and mechanosensors (e.g., *PIEZO1* and *PKD2*).

Results. The results showed that vitamin D3 induced THP-1 macrophage differentiation, which was characterized by upregulation of *CD14* and *CD36*, downregulation of *HLA-DRA*, upregulation of the *PKD2* (*TRPP2*), and an inverse relationship between *TCF7L2* and *LEF-1*, which were upregulated and downregulated respectively. The 1,000 Hz vibrations were sensed from the cells which upregulated *PIEZO1* and *TCF3*, but they did not induce expression of genes that would indicate macrophage differentiation. The mRNA transcription profile in the cells stimulated with the combined treatment was comparable to that of the cells stimulated by the vitamin only. The 1,000 Hz vibrations slightly weakened the effect of the vitamin for the regulation of *CD36* and *HLA-DMB* in the suspension cells, but without causing changes in the regulation patterns. The only exception was the upregulation of *TCF3* in the suspension cells, which was influenced by the vibrations. In the adherent cells, the vitamin D3 cancelled the upregulating effect of the 1,000 Hz vibrations and downregulated *TCF3*. The vitamin also cancelled the upregulation of *PIEZO1* gene by the 1,000 Hz vibrations in the combined treatment.

Conclusion. The mechanical stimulation with 1,000 Hz vibrations resulted in upregulation of *PIEZO1* in THP-1 cells, but it did not affect the differentiation process which was investigated in this study. Vitamin D3 induced THP-1 macrophage differentiation and could potentially influence M2 polarization as observed by upregulation of *CD36* and

Submitted 17 February 2021

Accepted 23 June 2021

Published 14 July 2021

Corresponding author

Fiona L. Henriquez,
fiona.henriquez@uws.ac.uk

Academic editor

Liudmila Leppik

Additional Information and
Declarations can be found on
page 21

DOI 10.7717/peerj.11773

© Copyright
2021 Simakou et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

downregulation of *HLA-DRA*. In addition, in THP-1 cells undergoing the combined stimulation, the gene expression patterns were influenced by vitamin D3, which also ablated the effect of the mechanical stimulus on *PIEZO1* upregulation.

Subjects Biochemistry, Bioengineering, Cell Biology, Molecular Biology

Keywords THP-1, Vitamin D3, Macrophage differentiation, *PIEZO1*, *PKD2*, Mechanosensation, TCF/LEF, Vibrational Stimulation

INTRODUCTION

Macrophages play important roles in health and disease through phagocytosis of pathogenic microorganisms, by releasing inflammatory mediators, by inducing and maintaining inflammation, and by removing apoptotic cells and repairing tissues (*Gordon, 2007; Mosser & Edwards, 2008*). Tissue-resident macrophages, derived from the yolk sac at the embryonic stage, are replicated in tissues to maintain cell number, and have different morphology and function depending on the tissue where they reside (*Lavin et al., 2015*). However, in the case of tissue injury or infection, monocytes derived from bone marrow circulating in peripheral blood, migrate to the affected tissues where they differentiate into macrophages, and are subsequently involved in the inflammatory response (*Shi & Pamer, 2011*).

THP-1 cells are human immortalized monocytes derived from acute monocytic leukemia and have been extensively used to study macrophage differentiation, functions, signaling pathways, and nutrient and drug transport (*Chanput, Mes & Wichers, 2014; Bosshart & Heinzelmann, 2016; Nurminen, Seuter & Carlberg, 2019*). In this study we investigated the THP-1 responses towards stimulation with vitamin D3 (1,25-dihydroxyvitamin D3), 1,000 Hz nanovibrations or both, in order to study the expression of genes that could indicate differentiation or mechanosensitivity changes in these cells. In the following text, the combined treatment refers to the combination of 50 nM vitamin D3 and 1,000 Hz vibrations. The cell responses were investigated separately in adherent and suspension THP-1 monocytes upon each stimulation, to avoid averaging results for both the cell types within the same population and consider their differences (*Fig. S2*).

Vitamin D has been shown to promote monocyte differentiation into macrophages and targets multiple genes (*Nurminen, Seuter & Carlberg, 2019*). The active form of vitamin D, 1,25-dihydroxyvitamin D3, is a lipophilic molecule which easily passes through biological membranes and binds with high-affinity to the receptor and transcription factor vitamin D receptor (VDR), which is primarily located in the nucleus (*Hausler et al., 2013*). The activation of vitamin D target genes is explained by the chromatin model (*Nurminen, Seuter & Carlberg, 2019*). The ligand-activated VDR molecules bind to a wide variety of enhancer regions that carry suitable binding motifs and are located within accessible chromatin. With the help of pioneer factors, such as PU.1, CEBPA, and GABPA, VDR increases the accessibility of chromatin at and around these enhancer regions (*Seuter, Neme & Carlberg, 2017; Seuter, Neme & Carlberg, 2018; Nurminen et al., 2019*). In THP-1 cells, 1,25-dihydroxyvitamin D3 stimulation significantly affects the binding strength of transcription factor CTCF to topologically associating domain (TAD) anchors, which

results in about 600 TADs becoming sensitive to vitamin D (Neme, Seuter & Carlberg, 2016). Looping of activated DNA-bound VDR to a transcription start site (TSS) at these promoter regions results in increased chromatin accessibility as well as of H3K27ac and H3K4me3 marks (Seuter, Neme & Carlberg, 2016; Nurminen et al., 2019). All these vitamin D-triggered changes in the local chromatin structure at enhancer and promoter regions finally lead to the activation of RNA polymerase II assembled on the respective TSSs and the start of mRNA synthesis. The vitamin may also affect gene expression by increasing the expression and the activity of transcription factors other than VDR, such as BCL6, NFE2, POU4F2, and ELF4 (Nurminen et al., 2015).

The effects of vitamin D have been studied in the context of macrophage differentiation from monocytes, but they are also extended into the effector macrophage responses (Hewison, 2010). In fact, normal human macrophages are able to synthesize 1,25-dihydroxyvitamin D₃ when stimulated with interferon gamma (IFN γ) (Phillip Koeffler et al., 1985). The vitamin D is involved in the regulation of T cell and macrophage effector functions, primarily via localized autocrine or paracrine synthesis of 1,25-dihydroxyvitamin D₃ from its precursor 25-hydroxyvitamin D₃ (Hewison, 2010). In addition, vitamin D deficiency is prevalent in multiple autoimmune diseases, such as multiple sclerosis, type 1 diabetes, systemic lupus erythematosus, and alopecia areata, and it is highly associated with the risk of autoimmunity (Yang et al., 2013; Lin, Meng & Song, 2019). Vitamin D has been implicated in prevention and protection from autoimmune diseases by immunomodulation of macrophage, dendritic cell, and T cell responses (Hewison, 2010; Yang et al., 2013).

In the recent years, interest has been given to the mechanobiology of macrophages, which like other immune cells have evolved mechanisms to perceive and respond to the mechanical forces around them (Kim et al., 2019). The cellular functions of tissue-resident macrophages and monocyte-derived macrophages are affected by the tissue-specific microenvironment, which can create many types of mechanical stress on cells (McWhorter, Davis & Liu, 2015; Mennens, Van den Dries & Cambi, 2017). Stiffness and topography, which are mechanical properties of the extracellular matrix, regulate the differentiation, proliferation, and function of macrophages such as phagocytosis (Patel et al., 2012). In monocytes, the PIEZO1 mechanotransduction in response to cyclical hydrostatic pressure, results in HIF1 α stabilization and secretion of molecules, such as endothelin-1 (EDN1), and neutrophil chemoattractant CXCL2 (Solis et al., 2019). In addition, macrophages in tissues are exposed to alterations of pressure which affect the secretion of cytokines such as IL-6, TNF- α and IL-1 β , (Ferrier, McEvoy & Andrew, 2000; Mevov et al., 2002). Other mechanical forces that these cells experience originate from dynamic mechanical loading, such as continuous and cyclic stretch and compression (McWhorter, Davis & Liu, 2015; Mennens, Van den Dries & Cambi, 2017).

Just like normal monocytes, THP-1 cells have shown to respond to mechanical stressors. For example, in models of atherosclerosis, biomechanical strain on THP-1 cells can induce expression of the class A scavenger receptor, an important lipoprotein receptor in atherogenesis (Yamamoto, Ikeda & Shimada, 2003). In addition, DNA microarray analysis has shown that cyclic mechanical strain in THP-1 cells induces expression of genes, some encoding for inflammatory markers such as IL-8 and IEX-1 (Yamamoto, Ikeda & Shimada,

2003). In these cells, biomechanical deformation influences the degradation of extracellular matrix, monocyte differentiation, and promotion of atherosclerosis (Yamamoto, Ikeda & Shimada, 2003). In addition, as THP-1 cells differentiate they become adherent, a process which may result in altered mechanosensitivity (Tsuchiya et al., 1982; Schwende et al., 1996).

In this study, the cells were mechanically stimulated using 1,000 Hz vibrations. The vibrational stimulation of 1,000 Hz frequency and nano-scale amplitude has been used to study *in vitro* osteogenic differentiation with successful results (Nikukar et al., 2013; Pemberton et al., 2015; Tsimbouri, 2015; Robertson et al., 2018), and in this study it was used to investigate any effect it may have on the differentiation of macrophages from THP-1 monocytes. Assessment of macrophage differentiation in response to externally applied vibrational stimuli can provide insights into monocyte mechanosensitivity and enquire the therapeutic effects of vibrational treatments in inflammatory diseases.

From a technical point of view, the experiments of this study were designed to give an insight into the differentiation process of THP-1 monocytes into macrophages under different stimulation parameters, compare between treatments, and look into mechanosensor mRNA expression.

MATERIALS & METHODS

THP-1 monocyte growth

THP-1 cells (ATCC[®] TIB-202[™]) were reconstituted from -80°C storage and allowed to recover for 2 weeks in cultures, splitting when confluency reached around 8×10^5 cells/mL. The culture medium needed for cell growth was composed of RPMI-1640 with L-glutamine (Capricorn Scientific; RPMI-HA), 10% Foetal Bovine Serum (FBS) (Gibco; A3160802) and 1% Antibiotic-Antimycotic 100X mix (Gibco; 15240062). The cells were cultured at 37°C , 5% CO_2 until ready for the experiments.

Experimental set up

The THP-1 cells were collected from T75 flasks (25 mL suspension) and pelleted by centrifugation at 1,500 rpm for 10 min. The experiment involved 4 replicates of untreated cells, 4 replicates of cells treated with 50 nM 1,25-dihydroxyvitamin D3 (Sigma-Aldrich, D1530), 4 replicates of cells treated with 1,000 Hz vibrations (amplitude range of 30 - 60 nm), and 4 replicates of cells treated simultaneously with 50 nM 1,25-dihydroxyvitamin D3 and 1,000 Hz stimulation. The cells underwent stimulation for 3 days (72 h). No medium or vitamin D3 replacement occurred for this duration of time. The cell density per each replicate at the start of the experiment was 1.5×10^5 cells/mL, in one mL suspension plated on 24-well plates (Thermo Fisher Scientific; 142475). The experiments took place at 37°C , 5% CO_2 , and 95% air incubator (LEEC 190D CO_2).

Preparation of the vibrational device

Plates (24-well plates) which would be clamped on the bioreactor had magnet sheets (First4Magnets; D-F4MA43MHP) attached 48 h before the start of the experiment, for better adhesion and removal of air pockets with time. In addition, the vibrational device

(nicknamed Nanokicking bioreactor) was incubated at 37 °C for 2 days prior to the start of experiments, which was the temperature at which the bioreactor was calibrated. Incubation prior to the experiment was also useful for avoiding condensation upon immediate translocation of the bioreactor from room temperature to incubator environment. The experiments took place in fanless incubator LEEC 190D to avoid additional external vibrations. The bioreactor's stability and generated vibrations were assessed using laser interferometry every 3 months (diagram in [Fig. S1](#)). The platform of the bioreactor was generating vibrations of 1, 000 Hz frequency and amplitude range 30–60 nm at the time of the experiments.

RNA extraction

The RNA was extracted separately for the suspension and adherent cells. Cell suspension was slowly removed and added to sterile RNase-free 1.5 mL tubes. The cells in suspension were pelleted by centrifugation at 3,000 rpm for 5 min. The supernatant was discarded and 1 mL Trizol reagent (Invitrogen; AM9738) was added to homogenize the pellet. For the adherent cells, 1 mL Trizol reagent was added directly in the wells. The lysed cells were homogenized using a 25 g syringe. The RNA extraction from the lysed cells in Trizol solution was done by separating the aqueous phase after addition of 0.2 mL chloroform and centrifugation at 13,000 rpm for 15 min at 4 °C. The RNA was washed with isopropanol and 75% ethanol and stored in 30 µL of nuclease-free water (Gibco; 10977035). Quantification of the RNA in ng/µL was done on Nanodrop 1000, using the RNA nucleic acid program.

DNase treatment

The DNase treatment was performed following the protocol of DNA-free Kit (Thermo Fisher Scientific, AM1906), in order to degrade any genomic DNA that contaminated the RNA solutions during extraction. The maximum RNA concentration for each sample was 5 µg per 50 µL DNase reaction. Removal of genomic DNA contamination allowed efficient detection of amplification during the real-time PCR.

Complementary DNA synthesis

The synthesis of cDNA was done as instructed on the protocol of High-Capacity cDNA Reverse transcription Kit (Applied Biosystems; 4368814). The reaction was comprised of 10 µL of 2X RT Mastermix and 10 µL of purified RNA solution from the previous step. Reaction was started by warming at 25 °C for 10 min, followed by incubation at 37 °C for 2 h for the synthesis of the cDNA, and termination of reaction at 85 °C for 5 min. The newly synthesized cDNA was stored at –20 °C until used for PCR reactions.

Real-time PCR

Real-time PCR was used to quantify gene expression in adherent and suspension THP-1 cells. The PCR amplifications were performed in 25 µL reactions containing 12.5 µL PowerUP SYBR Green Mastermix (Applied Biosystems; A25742); 0.5 µL Forward Primer and 0.5 µL Reverse Primer for the respective genes, 1µL of cDNA and topped up to 25µL with nuclease free water (Gibco; 10977035).

The primer pair used for amplification of the housekeeper *RPL37A* were *RPL37A* forward 5'-ATTGAAATCAGCCAGCACGC-3' and *RPL37A* reverse 5'-AGGAACCACAGTGCCAGATCC-3'. The primer pair used for amplification of the housekeeper *ACTB* were *ACTB* forward 5'-ATTGCCGACAGGATGCAGAA-3' and *ACTB* reverse 5'-GCTGATCCACATCTGCTGGAA-3'. The primer pair used for amplification of *CD36* were *CD36* forward 5'-TCACTGCGACATGATTAATGGTACA-3' and *CD36* reverse 5'-ACGTCGGATTCAAATACAGCATAGAT-3'. The primer pair used for amplification of *CD14* were *CD14* forward 5'-ACGCCAGAACCTTGTGAGC-3' and *CD14* reverse 5'-GCATGGATCTCCACCTCTACTG-3'. The primer pair for amplification of *HLA-DRA* were *HLA-DRA* forward 5'-TAAGGCACATGGAGGTGATG-3' and *HLA-DRA* reverse 5'-GTACGGAGCAATCGAAGAGG-3'. The primer pair used for amplification of *HLA-DMB* were *HLA-DMB* forward 5'-CTCTCACAGCACCTCAACCA-3' and *HLA-DMB* reverse 5'-TAGAAGCCCCACACATAGCA-3'. The primer pair used for amplification of *PIEZO1* were *PIEZO1* forward 5'-CATCTTGGTGGTCTCCTCTGTCT-3' and *PIEZO1* reverse 5'-CTGGCATCCACATCCCTCTCATC-3'. The primer pair used for detection of *PKD1* were *PKD1* forward 5'-CGCCGCTTCACTAGCTTCGAC-3' and *PKD1* reverse 5'-ACGCTCCAGAGGGAGTCCAC-3'. The primer pair used for amplification of *PKD2* were *PKD2* forward 5'-GCGAGGTCTCTGGGGAAC-3' and *PKD2* reverse 5'-TACACATGGAGCTCATCATGC-3'. The primer pair used for amplification of *NFAT2* were *NFAT2* forward 5'-CACTCCTGCTGCCTTACACA-3' and *NFAT2* reverse 5'-AAGATGCGAGCATGCGACTA-3'. The primer pair used for amplification of *TCF3* were *TCF3* forward 5'-TGACCTCCTGGACTTCAGC-3' and *TCF3* reverse 5'-ACCTGAACCTCCGAAGTGC-3'. The primer pair used for amplification of *TCF4* were *TCF4* forward 5'-AGTGCGATGTTTTACCTCC-3' and *TCF4* reverse 5'-CCTGAGCTACTTCTGTCTTC-3'. The primer pair used for the amplification of *TCF7L2* were *TCF7L2* forward 5'-CCGGGAAAGTTTGGGAAGAAG-3' and *TCF7L2* reverse 5'-ACTGAAAATGGAGGGTTCGG-3'. The primer pair used for amplification of *LEF-1* were *LEF-1* forward 5'-GACAGTGACCTAATGCACGT-3' and *LEF-1* reverse 5'-CCACCTTCTGCCAAGAATCT-3'.

The primers for TCFs and LEF-1 transcription factors were designed and tested by Dr. Robin Freeburn. Primers amplifying *PIEZO1* were designed using the NCBI primer design tool for the mRNA sequence [NM_001142864.4](#), and primers amplifying *CD14* were designed similarly for the mRNA sequences [NM_001174105.2](#) (*CD14* mRNA transcript variant 4), [NM_001040021.3](#) (*CD14* mRNA transcript variant 2), [NM_000591.4](#) (*CD14* mRNA transcript variant 1) and [NM_001174104.1](#) (*CD14* mRNA transcript variant 3). The primers for *NFAT2* were obtained from [Dagna, Pritchett & Lusso \(2013\)](#), primers for *HLA-DRA* and *HLA-DMB* were obtained from [Ulbricht et al. \(2012\)](#), primers for *PKD1* and *PKD2* were obtained from [Dalagiorgou et al. \(2013\)](#), and primers from *CD36*, *ACTB* and *RPL37A* were obtained from [Maefß, Sendelbach & Lorkowski \(2010\)](#).

The efficiency of primers taken from existing literature has been assessed in published papers ([Fukuda, Mitsuoka & Schmid-Schönbein, 2004](#); [Maefß, Sendelbach & Lorkowski, 2010](#); [Ulbricht et al., 2012](#); [Dagna, Pritchett & Lusso, 2013](#); [Dalagiorgou et al., 2013](#)). The primer efficiency was assessed prior to the experiments and was around 97% for all the investigated

genes. Similar PCR efficiency for each primer is necessary for relative quantification using the $\Delta\Delta C_T$ method (Livak & Schmittgen, 2001). The PCR efficiency was also assessed by melt curve analysis. The collected C_T values were used for the $\Delta\Delta C_T$ relative quantification of expression, comparing the stimulated cells to the untreated controls. The ΔC_T was obtained by comparison of C_T s of genes of interest to the mean C_T of two housekeeping genes *RPL37A* and *ACTB*. These housekeeping genes are considered to be the best for the analysis of RNA expression in THP-1 cells (Maefß, Sendelbach & Lorkowski, 2010).

Statistical analysis

The gene expression data are presented as mean of four replicates \pm SEM, with little exception where some particular genes were not detected in all replicates. The analysis of statistical significance between the stimulated cells versus controls, and between each type of stimulation was done using unpaired *T* test with Welch's correction. Statistical analysis was carried out using GraphPad Prism[®] version 6. *P* values < 0.05 were accepted as significant.

RESULTS

Regulation of genes encoding macrophage markers and antigen presenting molecules

Stimulation with vitamin D3, which also served as a positive control for the induction of differentiation, resulted in upregulation of the *CD14* and *CD36* mRNA in both adherent and suspension cells (Figs. 1A and 2A). Vitamin D3 also downregulated the *HLA-DRA* expression in adherent and suspension cells (Figs. 1A and 2A). The mRNA of *HLA-DMB* was upregulated for vitamin D3 stimulation in suspension cells (Fig. 1A). The *HLA-DMB* was not regulated in response to the vitamin D3 in the adherent cells (Fig. 2A).

The 1,000 Hz stimulation caused upregulation of *CD36* and downregulation of *HLA-DMB* in suspension cells (Fig. 1B), whereas in adherent cells it only downregulated the *HLA-DRA* (Fig. 2B).

The combined stimulation induced upregulation of *CD14* and *CD36* in both adherent and suspension cells (Figs. 1C and 2C). The *HLA-DRA* was downregulated in both cell types compared to the respective unstimulated control (Figs. 1C and 2C), whereas *HLA-DMB* was upregulated in suspension cells (Fig. 1C).

The expression values of *CD14*, *CD36*, *HLA-DRA* and *HLA-DMB* in stimulated cells versus controls are shown in Tables 1 and 2, for suspension and adherent cells respectively.

A comparison between the treatments was performed for the above genes in suspension (Fig. 3) and adherent cells (Fig. 4). The comparison between treatments is shown in detail in Table 3.

CD14 was upregulated only in response to the vitamin D3, as the mRNA levels were comparable to the cells stimulated by the vitamin only (Figs. 3A and 4A). Similarly, the upregulation of *CD36* in the adherent cells was only in response to the vitamin D3 in the combined treatment (Fig. 3A). In suspension cells undergoing the combined treatment, the 1,000 Hz stimulation weakened the upregulation of *CD36* by the vitamin D3, which was still higher than the upregulation caused by the 1,000 Hz vibrational stimulation




Figure 1 Gene expression in response to different stimulations in suspension THP-1 cells, compared to the unstimulated suspension THP-1 cells. (A) mRNA regulation in response to stimulation with 50nM of 1,25-dihydroxyvitamin D3. (B) mRNA regulation in response to 1,000 Hz vibrations (amplitude 30–60 nm). (C) mRNA regulation in response to the combined vitamin D (50 nM) and 1,000 Hz (30–60 nm amplitude) vibrations. Data presented as mean of four replicates \pm SEM. Statistical analysis between stimulated and control values was assessed by unpaired *T* test with Welch's correction. *P* values lower than 0.05 were considered statistically significant.

Full-size DOI: 10.7717/peerj.11773/fig-1

alone (Table 3). In the combined treatment, the 1,000 Hz vibrations also weakened the upregulation of *HLA-DMB* by the vitamin in the suspension cells (Table 3). Interestingly, the *HLA-DRA* was downregulated from all treatments at the same level in the adherent




Figure 2 Gene expression in response to different stimulations in adherent THP-1 cells, compared to the unstimulated adherent THP-1 cells. (A) mRNA regulation in response to stimulation with 50nM of 1,25-dihydroxyvitamin D3. (B) mRNA regulation in response to 1,000 Hz vibrations (amplitude 30–60 nm). (C) mRNA regulation in response to the combined vitamin D (50 nM) and 1,000 Hz (30–60 nm amplitude) vibrations. Data presented as mean of four replicates \pm SEM. Statistical analysis between stimulated and control values was assessed by unpaired *T* test with Welch's correction. *P* values lower than 0.05 were considered statistically significant.

Full-size DOI: 10.7717/peerj.11773/fig-2

Table 1 Expression of genes in stimulated suspension THP-1 cells compared to the unstimulated suspension cells at 72 h. Statistical analysis was performed using unpaired *T* test with Welch's correction. Fold change ($2^{-\Delta\Delta Ct}$) values higher than 1 indicate upregulation, whereas values between 0 and 1 indicate downregulation of mRNA transcripts in stimulated cells.

THP-1 cells in suspension				
Stimulation	mRNA	Roles	Fold change ($2^{-\Delta\Delta Ct}$) Stimulated cells vs Control	<i>P</i> value
50 nM 1,25(OH) ₂ D3 (72 h)	<i>CD14</i>	Macrophage marker	573.92	0.0161
	<i>CD36</i>	Macrophage marker	3.66	0.0004
	<i>HLA-DRA</i>	Antigen presentation	0.16	0.0001
	<i>HLA-DMB</i>	Antigen presentation	2.17	0.0016
	<i>PKD2</i>	Mechanosensory non-selective cation channel	2.32	0.0004
	<i>TCF4</i>	Transcription factor (unknown roles in macrophages)	0.5	0.001
	<i>TCF7L2</i>	Transcription factor (proliferation and differentiation)	1.85	0.0052
Vibrations 1,000 Hz (30–60 nm) (72 h)	<i>LEF-1</i>	Transcription factor (proliferation and differentiation)	0.37	0.0005
	<i>CD36</i>	Macrophage marker	1.43	0.013
	<i>HLA-DMB</i>	Antigen presentation	0.86	0.0404
	<i>PIEZO1</i>	Mechanosensory channel	1.39	0.0441
50 nM 1,25(OH) ₂ D3 + Vibrations 1,000 Hz (30–60 nm) (72 h)	<i>TCF3</i>	Transcription factor (unknown roles in macrophages)	1.69	0.0182
	<i>CD14</i>	Macrophage marker	428.9	0.0359
	<i>CD36</i>	Macrophage marker	2.66	0.0009
	<i>HLA-DRA</i>	Antigen presentation	0.19	<0.0001
	<i>HLA-DMB</i>	Antigen presentation	1.51	0.0071
	<i>PKD2</i>	Mechanosensory non-selective cation channel	2.38	0.0021
	<i>TCF3</i>	Transcription factor (unknown roles in macrophages)	1.27	0.0142
	<i>TCF4</i>	Transcription factor (unknown roles in macrophages)	0.55	0.0026
	<i>NFAT2</i>	Transcription factor (undefined roles in macrophages)	0.69	0.0434
	<i>TCF7L2</i>	Transcription factor (proliferation and differentiation)	2.1	0.0063
	<i>LEF-1</i>	Transcription factor (proliferation and differentiation)	0.66	0.0011

cells (Fig. 4A), but only the vitamin downregulated this gene in suspension cells (Fig. 3A; Table 3).

Regulation of genes encoding transcription factor

The stimulation with vitamin D3 downregulated *NFAT2* and *TCF3* in adherent cells (Fig. 2A). The *TCF4* and *LEF-1* were downregulated in both adherent and suspension cells stimulated with the vitamin (Figs. 1A and 2A). The *TCF7L2* mRNA was upregulated in

Table 2 Expression of genes in stimulated adherent THP-1 cells compared to the unstimulated adherent cells at 72 h. Statistical analysis was performed using unpaired T test with Welch's correction. Fold change ($2^{-\Delta\Delta Ct}$) values higher than 1 indicate upregulation, whereas values between 0 and 1 indicate downregulation of mRNA transcripts in stimulated cells.

Stimulation	mRNA	THP-1 cells adhered		P value
		Roles	Fold change ($2^{-\Delta\Delta Ct}$) Stimulated cells vs Control	
50 nM 1,25(OH) ₂ D3 (72 h)	<i>CD14</i>	Macrophage marker	650.9	0.0026
	<i>CD36</i>	Macrophage marker	3.13	0.0073
	<i>HLA-DRA</i>	Antigen presentation	0.09	0.0011
	<i>PKD2</i>	Mechanosensory non-selective cation channel	2.47	0.0096
	<i>TCF3</i>	Transcription factor (unknown roles in macrophages)	0.63	0.0134
	<i>TCF4</i>	Transcription factor (unknown roles in macrophages)	0.48	0.0032
	<i>NFAT2</i>	Transcription factor (undefined roles in macrophages)	0.52	0.0458
	<i>TCF7L2</i>	Transcription factor (proliferation and differentiation)	2.63	0.0313
Vibrations 1,000 Hz (30–60 nm) (72 h)	<i>LEF-1</i>	Transcription factor (proliferation and differentiation)	0.3	0.0491
	<i>HLA-DRA</i>	Antigen presentation	0.07	0.0022
	<i>PIEZO1</i>	Mechanosensory channel	11.44	0.0247
	<i>PKD2</i>	Mechanosensory non-selective cation channel	0.6	0.0236
	<i>NFAT2</i>	Transcription factor (undefined roles in macrophages)	0.12	0.0004
50 nM 1,25(OH) ₂ D3 + Vibrations 1,000 Hz (30–60 nm) (72 h)	<i>TCF3</i>	Transcription factor (unknown roles in macrophages)	4.73	0.04
	<i>CD14</i>	Macrophage marker	542.09	0.0011
	<i>CD36</i>	Macrophage marker	2.27	0.0227
	<i>HLA-DRA</i>	Antigen presentation	0.09	0.002
	<i>PKD2</i>	Mechanosensory non-selective cation channel	1.95	0.0013
	<i>TCF3</i>	Transcription factor (unknown roles in macrophages)	0.69	0.0232
	<i>TCF4</i>	Transcription factor (unknown roles in macrophages)	0.37	0.0016
	<i>NFAT2</i>	Transcription factor (undefined roles in macrophages)	0.31	0.0006
<i>TCF7L2</i>	Transcription factor (proliferation and differentiation)	2.36	0.0116	

response to the stimulation with vitamin D3 in suspension (Fig. 1A), and adherent cells (Fig. 2A).

The 1,000 Hz vibrational stimulation upregulated *TCF3* in both adherent and suspension cells compared to the respective controls (Figs. 1B and 2B). This type of stimulation also




Figure 3 Comparison of fold change values for different genes between the treatments in stimulated THP-1 suspension cells. Each treatment values were compared to the others using unpaired *T* test with Welch's correction. *P* values lower than 0.05 were considered statistically significant. Genes investigated encode for markers of macrophage differentiation (A), transcription factors (B), and mechanosensors (C).
[Full-size !\[\]\(b345a1c4255362eec3746050dd71ccac_img.jpg\) DOI: 10.7717/peerj.11773/fig-3](https://doi.org/10.7717/peerj.11773/fig-3)

downregulated *NFAT2* in the adherent cells (Fig. 2B). The mRNA expression of *TCF4*, *TCF7L2* and *LEF-1* were not affected by the vibrational stimulation (Figs. 1B and 2B).

The combined stimulation downregulated *NFAT2* in both adherent and suspension cells (Figs. 1C and 2C). The *TCF3* mRNA was downregulated in the adherent cells (Fig. 2C), but upregulated in the suspension cells (Fig. 1C). The *TCF4* was downregulated in both cell types, and *TCF7L2* was upregulated in both cell types (Figs. 1C and 2C). The *LEF-1* was downregulated in the suspension cells (Fig. 1C).




Figure 4 Comparison of fold change values for different genes between the treatments in stimulated THP-1 adherent cells. Each treatment values were compared to the others using unpaired *T* test with Welch's correction. *P* values lower than 0.05 were considered statistically significant. Genes investigated encode for markers of macrophage differentiation (A), transcription factors (B), and mechanosensors (C). Full-size [DOI: 10.7717/peerj.11773/fig-4](https://doi.org/10.7717/peerj.11773/fig-4)

The expression values of *TCF3*, *TCF4*, *TCF7L2* and *LEF-1* in stimulated cells versus controls are also shown in [Tables 1](#) and [2](#), for suspension and adherent cells respectively.

A comparison between the treatments was performed for these genes encoding transcription factors in suspension ([Fig. 3B](#)) and adherent cells ([Fig. 4B](#)), and shown in [Table 3](#).

Table 3 Comparison of THP-1 gene expression between different treatments. The arrows indicate upregulation or downregulation of the genes when comparing the different stimuli. Statistical analysis was performed using unpaired *T* test with Welch's correction.

mRNA	50 nM Vitamin D3 vs 1,000 Hz		50 nM Vitamin D3 vs 50bnM Vitamin D3+1,000 Hz		50 nM Vitamin D3+1,000 Hz vs 1,000 Hz	
	Adherent	Suspension	Adherent	Suspension	Adherent	Suspension
<i>CD14</i>	↑ 559.5 Fold (<i>p</i> = 0.003)	↑ 564.9 Fold (<i>p</i> = 0.016)	No difference (<i>p</i> = 0.24)	No difference (<i>p</i> = 0.41)	↑ 466.0 Fold (<i>p</i> = 0.001)	↑ 422.2 Fold (<i>p</i> = 0.036)
<i>CD36</i>	↑ 9.28 Fold (<i>p</i> = 0.004)	↑ 2.55 Fold (<i>p</i> = 0.0003)	No difference (<i>p</i> = 0.14)	↑ 1.38 Fold (<i>p</i> = 0.008)	↑ 6.73 Fold (<i>p</i> = 0.006)	↑ 1.85 Fold (<i>p</i> = 0.002)
<i>HLA-DRA</i>	No difference (<i>p</i> = 0.57)	↓ 0.15 Fold (<i>p</i> = 0.0001)	No difference (<i>p</i> = 0.87)	No difference (<i>p</i> = 0.14)	No difference (<i>p</i> = 0.39)	↓ 0.18 Fold (<i>p</i> < 0.0001)
<i>HLA-DMB</i>	No difference (<i>p</i> = 0.58)	↑ 2.51 Fold (<i>p</i> = 0.0008)	No difference (<i>p</i> = 0.09)	↑ 1.43 Fold (<i>p</i> = 0.006)	No difference (<i>p</i> = 0.61)	↑ 1.75 Fold (<i>p</i> = 0.002)
<i>NFAT2</i>	No difference (<i>p</i> = 0.075)	No difference (<i>p</i> = 0.14)	No difference (<i>p</i> = 0.28)	No difference (<i>p</i> = 0.15)	No difference (<i>p</i> = 0.052)	↓ 0.84 Fold (<i>p</i> = 0.02)
<i>TCF3</i>	↓ 0.13 Fold (<i>p</i> = 0.031)	↓ 0.58 Fold (<i>p</i> = 0.013)	No difference (<i>p</i> = 0.44)	↓ 0.77 Fold (<i>p</i> = 0.045)	↓ 0.15 Fold (<i>p</i> = 0.009)	No difference (<i>p</i> = 0.08)
<i>TCF4</i>	↓ 0.18 Fold (<i>p</i> = 0.032)	↓ 0.54 Fold (<i>p</i> < 0.0001)	No difference (<i>p</i> = 0.28)	No difference (<i>p</i> = 0.13)	↓ 0.14 Fold (<i>p</i> = 0.029)	↓ 0.59 Fold (<i>p</i> < 0.0001)
<i>TCF7L2</i>	No difference (<i>p</i> = 0.32)	↑ 1.84 Fold (<i>p</i> = 0.0034)	No difference (<i>p</i> = 0.63)	No difference (<i>p</i> = 0.29)	No difference (<i>p</i> = 0.46)	↑ 2.09 Fold (<i>p</i> = 0.0047)
<i>LEF-1</i>	n/a	↓ 0.36 Fold (<i>p</i> = 0.0014)	No difference (<i>p</i> = 0.3)	↓ 0.56 Fold (<i>p</i> = 0.011)	n/a	↓ 0.65 Fold (<i>p</i> = 0.0027)
<i>PKD1</i>	No difference (<i>p</i> = 0.2)	No difference (<i>p</i> = 0.06)	No difference (<i>p</i> = 0.15)	No difference (<i>p</i> = 0.91)	No difference (<i>p</i> = 0.54)	No difference (<i>p</i> = 0.055)
<i>PKD2</i>	↑ 4.09 Fold (<i>p</i> = 0.0036)	↑ 2.2 Fold (<i>p</i> < 0.0001)	No difference (<i>p</i> = 0.16)	No difference (<i>p</i> = 0.74)	↑ 3.24 Fold (<i>p</i> = 0.0001)	↑ 2.3 Fold (<i>p</i> = 0.0039)
<i>PIEZO1</i>	↓ 0.11 Fold (<i>p</i> = 0.026)	↓ 0.59 Fold (<i>p</i> = 0.015)	No difference (<i>p</i> = 0.42)	No difference (<i>p</i> = 0.478)	↓ 0.12 Fold (<i>p</i> = 0.027)	No difference (<i>p</i> = 0.052)

The *NFAT2* mRNA was downregulated in adherent cells for all the treatments, without difference between each other (Fig. 4B). In the suspension cells, the *NFAT2* was downregulated only for the combined stimulation (Fig. 3B).

In adherent cells, the *TCF3* mRNA was downregulated in response to vitamin D3 but upregulated for the 1,000 Hz stimulation. In the adherent cells, the vitamin cancelled the upregulating effect of the 1,000 Hz vibration and downregulated *TCF3*, at comparable levels to the cells stimulated with vitamin D3 only (Fig. 4B; Table 3). However, in suspension cells, the *TCF3* upregulation was influenced by the 1,000 Hz vibrations, and the mRNA levels were comparable to the cells stimulated with the 1,000 Hz vibrations alone (Table 3; Fig. 3B).

The *TCF4* mRNA was downregulated in response to vitamin D3 stimulation in both suspension and adherent cells. In the combined treatment, *TCF4* was influenced by the vitamin only. The 1,000 Hz did not have any influence on the expression of this gene neither alone nor in combination with the vitamin (Figs. 3B and 4B). Similarly, the upregulation of *TCF7L2* mRNA was influenced only by the vitamin D3 in both adherent and suspension

cells, with the 1,000 Hz stimulation having no effect on the cells when applied alone or in combination with the vitamin (Figs. 3B and 4B).

The mRNA for *LEF-1* was downregulated in response to vitamin D3 stimulation. In the adherent cells little RNA was obtained for this gene, and no amplification was detected for the 1,000 Hz stimulation (Table 3). This needs to be investigated in the future to explain whether the lack of amplification was due to very low transcripts levels in total RNA, or because of some inhibitory effect that 1,000 Hz vibrations may have. In the suspension cells stimulated with the combined treatment, the 1,000 Hz weakened the downregulating effect of the vitamin D3, however the vitamin influenced the downregulation (Fig. 3B).

Regulation of genes encoding mechanosensors *PIEZO1*, *PKD1* and *PKD2*

The stimulation with the vitamin D3 resulted in upregulation of *PKD2* (*TRPP2*) mRNA in both adherent and suspension cells. The stimulation with vitamin D3 alone did not affect the expression of *PIEZO1* or *PKD1* (*TRPP1*) (Figs. 1A and 2A).

The 1,000 Hz vibrational stimulation resulted in upregulation of *PIEZO1* mRNA in both adherent and suspension cells. In the adherent cells, the stimulation downregulated *PKD2* mRNA. The vibrational stimulation did not affect *PKD1* expression (Figs. 1B and 2B).

The combined treatment resulted in the upregulation of *PKD2* mRNA in both adherent and suspension cells. *PIEZO1* and *PKD1* were not regulated in cells stimulated with the combined treatment (Figs. 1C and 2C).

The expression patterns of *PKD2* and *PIEZO1* in stimulated cells versus controls are shown in Tables 1 and 2, for suspension and adherent cells, respectively.

A comparison between treatments was performed for these genes encoding mechanosensors in the suspension (Fig. 3C) and adherent cells (Fig. 4C), and shown in Table 3.

The expression of *PKD2* was affected only by the vitamin D3, which also cancelled the downregulation effect of the 1,000 Hz in the adherent cells stimulated with the combined treatment (Table 3).

The *PIEZO1* upregulation occurred only in response to the stimulation with 1,000 Hz vibrations, but in the combined treatment the vitamin cancelled the upregulating effect of the vibrational stimulation (Table 3). The expression of *PKD1* mRNA was not affected by any of the stimulation methods (Table 3).

A comparison of *PIEZO1* expression between adherent and suspension cells stimulated with 1,000 Hz vibrations and the combined treatment was performed (Fig. S3). The 1,000 Hz vibrations upregulated *PIEZO1* stronger in adherent cells, than in the suspension cells (Fig. S3A). No difference was observed between the adherent and suspension cells stimulated with the combined treatment (Fig. S3B).

DISCUSSION

TCF/LEF pathway and gene nomenclature

TCF/LEF pathway plays roles in monocyte and macrophage differentiation (Thiele et al., 2001). It must be mentioned that some confusion exists about the nomenclature of the

TCFs. The mammalian TCF/LEF family comprises of four nuclear factors designated TCF7, LEF1, TCF7L1, and TCF7L2, which are also known as TCF1, LEF1, TCF3, and TCF4, respectively ([Hrckulak et al., 2016](#)). Confusion also exists between the nomenclature of genes and the corresponding products. For example, a gene called *TCF3* (NCBI gene ID: 6929), also known as *E2A*, encodes a product that is different from TCF3 encoded from *TCF7L1* (NCBI Gene ID: 83439). Similarly, *TCF4* (NCBI gene ID: 6925), encodes for TCF4 which is a different protein from the TCF4 encoded from *TCF7L2* (NCBI Gene ID: 6934). In this experiment, the mRNA investigated belongs to genes *TCF3* (*E2A*), *TCF4* (*E2-2*), *TCF7L2* and *LEF-1*, with the last two investigated in the context of WNT canonical pathways in monocyte-derived macrophages ([Malsin et al., 2019](#)). The pathways which involve *TCF3* and *TCF4* gene products can be complex and are not elucidated in context of monocyte to macrophage differentiation.

Vitamin D3 induced macrophage differentiation and downregulated HLA-DRA

Vitamin D3 has shown to target multiple monocyte genes and promote monocyte differentiation into macrophages ([Nurminen, Seuter & Carlberg, 2019](#)). Similarly, this study demonstrated that vitamin D3 stimulation induced differentiation of THP-1 monocytes into macrophages, when looking at transcriptional regulation of *CD14*, *CD36* and transcription factors *TCF7L2* (encoding TCF4) and *LEF-1* ([Tables 1 and 2](#)).

The stimulation with vitamin D3 resulted in upregulation of the *CD14* and *CD36* mRNA in both adherent and suspension cells. This pattern of regulation for these two genes was expected to occur during macrophage differentiation from monocytes ([Zhang et al., 1994](#); [Maeß, Sendelbach & Lorkowski, 2010](#)). The *CD14* is an important marker of the THP-1 differentiation into macrophages which upregulates strongly upon vitamin D3 stimulation ([Schwende et al., 1996](#); [Gocek et al., 2012](#)), as was also observed in this study. Furthermore, *CD14* and *CD36* are primary target genes for vitamin D3 in THP-1 monocytes ([Nurminen, Seuter & Carlberg, 2019](#)). The *CD14* and *CD36* are proteins involved in macrophage functions. *CD14* cooperates with Toll-like receptor 4 (TLR4) to mediate the macrophage immune response to bacterial lipopolysaccharide (LPS) ([Zanoni et al., 2011](#)).

CD36 is a scavenger receptor which has been associated with M2 polarization and enhanced phagocytosis ([Pennathur et al., 2015](#); [Woo et al., 2016](#)).

It has been reported that M2 activation of bone marrow-derived macrophages with IL-4 has resulted in upregulation of *CD36* expression, whereas M1 activation with LPS and interferon- γ has resulted in downregulation of the receptor ([Pennathur et al., 2015](#)).

During kidney injury, *CD36* is an important phenotypic marker of profibrotic M2 macrophages and a key phagocytic receptor for the clearance of apoptotic cells ([Pennathur et al., 2015](#)). Similarly, during the resolution phase of stroke, *CD36* macrophages have a reparative role through phagocytosis ([Woo et al., 2016](#)).

Vitamin D3 downregulated the expression of *HLA-DRA* in differentiating THP-1 cells ([Tables 1 and 2](#)). *HLA-DR* has been described as an M1 marker, which is upregulated in THP-1 and monocyte - derived macrophages stimulated with IFN γ /LPS, whereas its expression is very low with IL-4/IL-13 stimulation ([Yang et al., 2016](#)). The decreased

HLA-DR expression in monocytes has also been associated with anti-inflammatory states or immunosuppression. HLA-DR expression is decreased in all monocyte subsets upon IL-10 exposure *in vitro* and during septic shock (Monneret *et al.*, 2004; Lee *et al.*, 2017), whereas monocytes that have diminished or no HLA-DR expression, called CD14⁺HLA-DR^{lo/neg} monocytes, have emerged as important mediators of tumor-induced immunosuppression (Mengos, Gastineau & Gustafson, 2019).

Downregulation of the HLA-DR protein has been observed in primary monocytes treated with vitamin D3 (Tokuda & Levy, 1996), as well as in dendritic cells (Ferreira *et al.*, 2015). In dendritic cells, the downregulation of HLA-DR has been suggested to be part of tolerance processes induced by vitamin D3 signaling (Ferreira *et al.*, 2015).

The upregulation of *CD36* and downregulation of *HLA-DRA* mRNA by vitamin D3 in day 3-differentiating THP-1 macrophages, could indicate predisposition for M2 polarization.

In addition, vitamin D3 stimulation upregulated the mRNA of *HLA-DMB* in suspension cells. This molecule is important for antigen loading of the MHC class II by removal of CLIP from HLA-DR (Riberdy *et al.*, 1992; Sloan *et al.*, 1995). In one study, HIV-infected THP-1 monocytes had loss of mRNA for *HLA-DR*, but the mRNAs for *HLA-DM* continued to be transcribed, showing that genes may have non-corresponding expression patterns (Shao & Sperber, 2002), similar to what was observed in this study.

This study also identified an inverse relationship between *TCF7L2* and *LEF-1* mRNA regulation during vitamin D3-induced macrophage differentiation. The *TCF7L2* (encoding TCF4) in combination with β -catenin forms a complex that regulates expression of genes in monocytes and it is thus involved in the differentiation process (Thiele *et al.*, 2001; Tickenbrock, 2006; Malsin *et al.*, 2019), whereas LEF-1 facilitates nuclear localization of β -catenin and enhances proliferation in acute myeloid leukemia cells, including THP-1 cells (Morgan *et al.*, 2019). Therefore, the downregulation of *LEF-1* and the upregulation of *TCF7L2* could indicate decreased proliferation and increased differentiation as THP-1 monocytes become macrophages (Schwende *et al.*, 1996; Thiele *et al.*, 2001; Morgan *et al.*, 2019). The inverse relationship of *TCF7L2* and *LEF-1* has also been related to shifts in differentiation and proliferation states in other cancer cells (Kriegel *et al.*, 2010; Eichhoff, *et al.*, 2011). This pattern of regulation for these two genes can be signature of THP-1 monocyte to macrophage differentiation.

Another transcription factor downregulated in adherent cells in response to the vitamin D3 was *NFAT2*. The NFATs are important transcription factors for production of proinflammatory cytokines in T and B cells (Macian, 2005), but their roles are not only limited to the adaptive immune cells. It has been showed that the NFATs are required for Toll-like receptor (TLR)-initiated innate immune responses in bone marrow-derived macrophages (Minematsu *et al.*, 2011). In THP-1 monocytes *in vitro*, the *NFAT2* has shown to inhibit the release of high mobility protein box-1 (HMGB1) (Zhao *et al.*, 2016), a proinflammatory protein with roles in inflammation and autoimmunity (Magna & Pisetsky, 2014). The suppression of *NFAT2* expression by siRNA has resulted in increased HMGB1 in the supernatant of cells (Zhao *et al.*, 2016). In T cells, 1,25-dihydroxyvitamin D3 and its receptor complex (VDR-RXR) have shown to inhibit NFAT activity

(Wöbke, Sorg & Steinhilber, 2014), but its effect on monocytes and *NFAT2* mRNA are not known. In this study, the downregulation of *NFAT2* mRNA in the adherent cells, which are considered to be in a more advanced stage of differentiation than the suspension cells (Tsuchiya et al., 1982; Schwende et al., 1996) (Fig. S2), could be related to the production of proinflammatory proteins after the maturation of the monocytes into macrophages.

The vitamin D3 stimulation also downregulated *TCF3* (encoding E2A) in adherent cells, and *TCF4* (encoding E2-2) in both cell types compared to the respective controls. The roles of the products of these genes are not known in monocyte biology and macrophage differentiation, but as demonstrated in this study they are regulatable upon vitamin D3 stimulation.

The stimulation with vitamin D3 had no effect on the regulation of *PIEZO1* or *PKD1*, but it upregulated *PKD2* (*TRPP2*) mRNA in both suspension and adhesion cells. The roles of polycystin 2 (product of *PKD2*) are not known in THP-1 monocytes, but the results of this study suggest that the *PKD2* mRNA upregulation can be signature of vitamin D3-induced differentiation.

Monocyte responses to 1,000 Hz vibrational stimulation

The THP-1 monocytes are responsive to mechanical stressors. Biomechanical strain on THP-1 cells can induce expression of the class A scavenger receptor, degradation of extracellular matrix, monocyte differentiation, and promotion of atherosclerosis (Yamamoto, Ikeda & Shimada, 2003). In addition, DNA microarray analysis has shown that cyclic mechanical strain in THP-1 cells induces expression of genes, some encoding for inflammatory markers such as IL-8 and IEX-1 (Yamamoto, Ikeda & Shimada, 2003). Furthermore, upon differentiation, THP-1 cells become adherent (Tsuchiya et al., 1982; Schwende et al., 1996), which may result in altered mechanosensitivity. This study used 1,000 Hz vibrations as artificially applied mechanical stimulation, in order to study the mechanosensitivity of THP-1 monocytes and assess if it could affect macrophage differentiation.

The vibrational 1,000 Hz stimulation resulted in upregulation of *PIEZO1* transcripts in both suspension (Table 1) and adhesion cells (Table 2). *PIEZO1* channels are considered professional mechanosensory proteins, capable of sensing and converting mechanical stimuli (Zhong et al., 2018). Little is known about the mechanosensory roles of these channels in monocytes and macrophages. RNA expression analysis presented in cell atlas shows *PIEZO1* expressed in monocytes and macrophages, as well as in THP-1 cells (Human Protein Atlas, Cell Type RNA, Piezo1). *PIEZO2* expression has not been detected in blood cells, including monocytes, whereas its expression in THP-1 cells is negligible (Human Protein Atlas, Cell Type RNA, Piezo2).

In monocytes, *PIEZO1* has shown to signal in response to cyclical hydrostatic pressure, resulting in HIF1 α stabilization and secretion of molecules, such as endothelin-1 (EDN1), and neutrophil chemoattractant CXCL2 (Solis et al., 2019). The *PIEZO1* signaling to the cyclical pressure has induced inflammation and infiltration of monocytes, which recruit neutrophils in order to clear pulmonary *Pseudomonas aeruginosa* infection via EDN1 (Solis et al., 2019). In this study, we demonstrated that THP-1 cells upregulate *PIEZO1* mRNA

in response to 1,000 Hz vibrational stimulation, when applied in isolation. However, the biological significance of such regulation remains to be elucidated. In addition, the *PIEZO1* mRNA upregulation in response to the 1,000 Hz vibrations was stronger in the adherent cells that were in contact with the vibrating surface, compared to the floating suspension cells (Fig. S3A). This could indicate potential involvement of mechanotransduction for the regulation of *PIEZO1* expression in 1,000 Hz vibrated THP-1 monocytes.

The 1,000 Hz stimulation also caused *HLA-DRA* downregulation in adherent cells like vitamin D3, but when combined with the vitamin it did not show any synergetic effect (Table 3). Another gene which was upregulated during the stimulation with 1,000 Hz vibrations, was *TCF3*. This gene was upregulated in both suspension and adherent cells (Table 1; Table 2), but the role of this gene and its products are not known in monocytes. The 1,000 Hz vibrations downregulated the *NFAT2* mRNA at the same levels as the vitamin D3 in adherent cells (Table 2), and just like the vitamin it did not regulate this gene in suspension cells (Table 1).

The vibrational stimulation had no effect on the regulation of other transcription factors such as *TCF4*, *TCF7L2* and *LEF-1*, which were influenced by the vitamin D3 only (Tables 1–3).

The upregulation of *PIEZO1* and *TCF3* upon the application of the 1,000 Hz stimulation was interesting, but it was not associated with macrophage differentiation, because there was no transcriptional regulation for genes such as *CD14*, *TCF7L2* and *LEF-1* which would indicate transition from monocytes to macrophages. The *CD36* was upregulated for the 1,000 Hz stimulation in suspension cells. However, in adherent cells, the *CD36* mRNA levels were comparable to the unstimulated adherent controls.

The effects of the combined treatment on gene expression and comparison to vitamin D3 and 1,000 Hz vibrations

The combined treatment induced macrophages differentiation, but the process was influenced mostly by the vitamin D3 (Table 3).

The upregulation of *CD14* in suspension and adherent cells undergoing the combined treatment was comparable to cells stimulated with vitamin D3 only (Table 3). The *CD36* mRNA was upregulated in the adherent cells at comparable level to the cells stimulated with vitamin D3 only. However, in the suspension cells the 1,000 Hz had slightly weakened the upregulation of *CD36* by the vitamin D3. The combination of both stimuli resulted in lower mRNA expression than the stimulation with the vitamin, but higher than the stimulation with the 1,000 Hz, hence it could be said that the 1,000 Hz weakened the upregulating effect of the vitamin (Table 3).

Even though when applied in isolation the 1,000 Hz stimulation caused *HLA-DRA* downregulation in adherent cells at similar levels to vitamin D3, in the combined stimulation it did not show any synergetic effect. The downregulation of *HLA-DRA* in suspension cells undergoing the combined treatment was comparable to the cells stimulated with the vitamin D3 only, showing that in the combined treatment this gene was influenced only by the vitamin (Table 3).

In the suspension cells undergoing the combined treatment, the 1,000 Hz weakened the upregulation of *HLA-DMB* by the vitamin. When applied in isolation the 1,000 Hz vibrations downregulated *HLA-DMB*, however, in the combined treatment the vitamin overshadowed the effect of the vibrational stimulus and caused upregulation (Table 3).

The combined treatment downregulated *NFAT2* at comparable levels to both the vitamin D3 and 1,000 Hz treatments when applied alone in the adherent cells. However, in the suspension cells, the 1,000 Hz and the vitamin D3 may have synergistically caused the downregulation of *NFAT2* in suspension cells, because the vitamin and the 1,000 Hz did not regulate this gene when applied in isolation.

The regulation of *TCF4* and *TCF7L2* in the cells stimulated with the combined treatment was comparable to the cells stimulated with vitamin D3, and the 1,000 Hz stimulation had no effect on these genes in the combined treatment (Table 3), similar to when it was applied in isolation (Tables 1 and 2). The 1,000 Hz vibrations however, weakened the downregulating effect that the vitamin had on the mRNA encoding *LEF-1* in the suspension cell. In the adherent cells, the *LEF-1* mRNA in stimulated with the combined treatment was comparable to the unstimulated controls, but since the mRNA for this gene was not detected in cell stimulated with vibrations only, comparison could not take place (Table 3).

In the presence of the vitamin D3, the effect of 1,000 Hz stimulation on the regulation of *PIEZO1* was cancelled in both adherent and suspension cells. Furthermore, in adherent cells, the vitamin D3 cancelled the upregulating effect of 1,000 Hz on the *TCF3* and downregulated the gene (Tables 2 and 3). However, in suspension cells the 1,000 Hz stimulation continued to upregulate *TCF3* even in the presence of the vitamin (Table 3). This was the only case in which the effects of 1,000 Hz strongly influenced the expression pattern of a gene in the presence of the vitamin.

CONCLUSIONS

This study demonstrated that the stimulation with 50nM vitamin D3 for 3 days drives THP-1 macrophage differentiation, as was determined by upregulation of *CD14*, *CD36* and *TCF7L2*, and downregulation of *LEF-1*. The differentiation induced by vitamin D3 was accompanied by downregulation of *HLA-DRA* and upregulation of *PKD2* mRNA. Other genes that were regulated during vitamin D3-induced macrophage differentiation included *TCF3* and *TCF4* in both suspension and adherent cells, and *NFAT2* in adherent cells. The upregulation of the mechanosensitive non-selective cation channel *PKD2* mRNA could suggest a role during THP-1 macrophage differentiation, whereas the upregulation of *CD36* and downregulation of *HLA-DRA* mRNA could be indicative of predisposition for M2 polarization.

The vibrational stimulation which was used for the mechanical stimulation of cells did not induce the macrophage differentiation process because there was no transcriptional regulation of *CD14* and *TCF/LEF* transcription factors. However, the 1,000 Hz vibrations influenced upregulation of *PIEZO1* and *TCF3* in both adherent and suspension cells. Furthermore, in adherent cells, the vibrational stimulation downregulated *NFAT2* and *HLA-DRA* at comparable levels to the vitamin D3 stimulated adherent cells. This indicated

that while the 1,000 Hz vibrations did not induce differentiation, they induced regulation of genes in the THP-1 cells. However, the biological importance of such response remains to be elucidated.

In the combined treatment, the 1,000 Hz vibrations interfered with the regulation of some genes by the vitamin D3 but without changing their regulation pattern. The only exception was *TCF3* in suspension cells stimulated with the combined treatment, which was upregulated by the 1,000 Hz vibrations against the downregulating influence of the vitamin D3. The biological importance of such interference remains to be elucidated. However, the mRNA regulation patterns of the other genes of interest in the combined treatment were in response to vitamin D3 stimulation.

Furthermore, the influence of the 1,000 Hz stimulus in the presence of the vitamin D3 was cancelled (e.g., for *PIEZO1* in both cell types), overshadowed (e.g., for *CD36* in suspension cells), or cancelled and reversed (e.g., *PKD2* in adherent cells). This can have implication for the medicinal application of the 1,000 Hz (nano-scale amplitude) vibrations, because in inflamed tissues rich in chemical signals such as cytokines and chemokines, the cells may lose the ability to sense and respond to such mechanical stimulus.

Further work is necessary to assess the reproducibility of the observations of this study, especially in response to the 1,000 Hz vibrational stimulation. This study was limited by the technology, which was not provided for repeated runs and further work. Increased replicates, expanded time-points, assessment of protein expression, and use of primary monocytes to compare to THP-1 cell responses, are recommended for future work from the authors of this report. In addition, the effects of different frequencies and of vibrations applied in cyclical short-term patterns remain to be studied, in order to expand our understanding of THP-1 cell responses towards the vibrational stimulation.

Overall, this study presents experimental results indicating that the vibrational mechanical forces can be sensed by THP-1 monocytes, but that the chemical ligands such as vitamin D3 remain superior for the induction of macrophage differentiation.

ACKNOWLEDGEMENTS

The authors would also like to thank Dr Paul Campsie and Prof Stuart Reid for providing a “Nanokick” bioreactor for the vibrational stimulation of the cells, and for access to the interferometer for the assessment of bioreactor’s stability.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This work was supported by Alopecia UK and the University of West of Scotland. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
Alopecia UK and the University of West of Scotland.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Theodoros Simakou conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Robin Freeburn and Fiona Henriquez conceived and designed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

The raw fold change values are available in the [Supplemental Files](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.11773#supplemental-information>.

REFERENCES

- Bosshart H, Heinzelmann M. 2016.** THP-1 cells as a model for human monocytes. *Annals of Translational Medicine* **4(21)**:438–438.
- Chanput W, Mes JJ, Wichers HJ. 2014.** THP-1 cell line: an in vitro cell model for immune modulation approach. *International Immunopharmacology* **23(1)**:37–45.
- Pemberton GD, Childs P, Reid S, Nikukar H, Tsimbouri PM, Gadegaard N, Curtis AS, Dalby MJ. 2015.** Nanoscale stimulation of osteoblastogenesis from mesenchymal stem cells: nanotopography and nanokicking. *Nanomedicine* **10(4)**:547–560 DOI [10.2217/nnm.14.134](https://doi.org/10.2217/nnm.14.134).
- Dagna L, Pritchett JC, Lusso P. 2013.** Immunomodulation and immunosuppression by human herpesvirus 6A and 6B. *Future Virology* **8(3)**:273–287.
- Dalagiorgou G, Piperi C, Georgopoulou U, Adamopoulos C, Basdra EK, Papavassiliou AG. 2013.** Mechanical stimulation of polycystin-1 induces human osteoblastic gene expression via potentiation of the calcineurin/NFAT signaling axis. *Cellular and Molecular Life Sciences* **70(1)**:167–180 DOI [10.1007/s00018-012-1164-5](https://doi.org/10.1007/s00018-012-1164-5).
- Eichhoff OM, Weeraratna A, Zipser MC, Denat L, Widmer DS, Xu M, Kriegel L, Kirchner T, Larue L, Dummer R, Hoek KS. 2011.** Differential LEF1 and TCF4 expression is involved in melanoma cell phenotype switching. *Pigment Cell & Melanoma Research* **24(4)**:631–642 DOI [10.1111/j.1755-148X.2011.00871.x](https://doi.org/10.1111/j.1755-148X.2011.00871.x).
- Ferreira GB, Vanherwegen A-S, Eelen G, Gutiérrez ACF, Van Lommel L, Marchal K, Verlinden L, Verstuyf A, Nogueira T, Georgiadou M, Schuit F, Eizirik DL, Gysemans C, Carmeliet P, Overbergh L, Mathieu C. 2015.** Vitamin D3 induces tolerance in human dendritic cells by activation of intracellular metabolic pathways. *Cell Reports* **10(5)**:711–725.

- Ferrier GM, McEvoy A, Evans CE, Andrew JG. 2000.** The effect of cyclic pressure on human monocyte-derived macrophages in vitro. *The Journal of Bone and Joint Surgery. British Volume* **82(5)**:755–759.
- Fukuda S, Mitsuoka H, Schmid-Schönbein GW. 2004.** Leukocyte fluid shear response in the presence of glucocorticoid. *Journal of Leukocyte Biology* **75(4)**:664–670 DOI [10.1189/jlb.1003464](https://doi.org/10.1189/jlb.1003464).
- Gocek E, Bauraska H, Marchwicka A, Marcinkowska E. 2012.** Regulation of leukemic cell differentiation through the vitamin D receptor at the levels of intracellular signal transduction, gene transcription, and protein trafficking and stability. *Leukemia Research and Treatment* **2012**:1–11.
- Gordon S. 2007.** The macrophage: past, present and future. *European Journal of Immunology* **37(S1)**:S9–S17 DOI [10.1002/eji.200737638](https://doi.org/10.1002/eji.200737638).
- Hewison M. 2010.** Vitamin D and the immune system: new perspectives on an old theme. *Endocrinology and Metabolism Clinics of North America* **39(2)**:365–379.
- Hrckulak D, Kolar M, Strnad H, Korinek V. 2016.** TCF/LEF Transcription Factors: an Update from the Internet Resources. *Cancer* **8(7)**:70.
- Haussler MR, Whitfield GK, Kaneko I, Haussler CA, Hsieh D, Hsieh J-C, Jurutka PW. 2013.** Molecular mechanisms of vitamin D action. *Calcified Tissue International* **92(2)**:77–98 DOI [10.1007/s00223-012-9619-0](https://doi.org/10.1007/s00223-012-9619-0).
- Kim J-K, Shin YJ, Ha LJ, Kim D-H, Kim D-H. 2019.** Unraveling the mechanobiology of the immune system. *Advanced Healthcare Materials* **8(4)**:e1801332 DOI [10.1002/adhm.201801332](https://doi.org/10.1002/adhm.201801332).
- Kriegl L, Horst D, Reiche JA, Engel J, Kirchner T, Jung A. 2010.** LEF-1 and TCF4 expression correlate inversely with survival in colorectal cancer. *Journal of Translational Medicine* **8(1)**:123 DOI [10.1186/1479-5876-8-123](https://doi.org/10.1186/1479-5876-8-123).
- Lavin Y, Mortha A, Rahman A, Merad M. 2015.** Regulation of macrophage development and function in peripheral tissues. *Nature Reviews Immunology* **15(12)**:731–744.
- Lee J, Tam H, Adler L, Iltad-Minnihan A, Macaubas C, Mellins ED. 2017.** The MHC class II antigen presentation pathway in human monocytes differs by subset and is regulated by cytokines. *PLOS ONE* **12(8)**:e0183594 DOI [10.1371/journal.pone.0183594](https://doi.org/10.1371/journal.pone.0183594).
- Lin X, Meng X, Song Z. 2019.** Vitamin D and alopecia areata: possible roles in pathogenesis and potential implications for therapy. *American Journal of Translational Research* **11(9)**:5285–5300.
- Livak KJ, Schmittgen TD. 2001.** Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* **25(4)**:402–408.
- Macian F. 2005.** NFAT proteins: key regulators of T-cell development and function. *Nature Reviews Immunology* **5(6)**:472–484.
- Maeß MB, Sendelbach S, Lorkowski S. 2010.** Selection of reliable reference genes during THP-1 monocyte differentiation into macrophages. *BMC Molecular Biology* **11(1)**:90 DOI [10.1186/1471-2199-11-90](https://doi.org/10.1186/1471-2199-11-90).
- Magna M, Pisetsky DS. 2014.** The role of HMGB1 in the pathogenesis of inflammatory and autoimmune diseases. *Molecular Medicine* **20(1)**:138–146 DOI [10.2119/molmed.2013.00164](https://doi.org/10.2119/molmed.2013.00164).

- Malsin ES, Kim S, Lam AP, Gottardi CJ. 2019.** Macrophages as a Source and Recipient of Wnt Signals. *Frontiers in Immunology* **10**:1813 DOI [10.3389/fimmu.2019.01813](https://doi.org/10.3389/fimmu.2019.01813).
- McWhorter FY, Davis CT, Liu WF. 2015.** Physical and mechanical regulation of macrophage phenotype and function. *Cellular and Molecular Life Sciences* **72**(7):1303–1316 DOI [10.1007/s00018-014-1796-8](https://doi.org/10.1007/s00018-014-1796-8).
- Mengos AE, Gastineau DA, Gustafson MP. 2019.** The CD14+HLA-DRlo/neg monocyte: an immunosuppressive phenotype that restrains responses to cancer immunotherapy. *Frontiers in Immunology* **10**:1147. Available at <https://www.frontiersin.org/article/10.3389/fimmu.2019.01147/full>.
- Mennens SFB, van den Dries K, Cambi A. 2017.** Role for mechanotransduction in macrophage and dendritic cell immunobiology. *Results and problems in cell differentiation* **62**:209–242 DOI [10.1007/978-3-319-54090-0_9](https://doi.org/10.1007/978-3-319-54090-0_9).
- Mevoy A, Jeyam M, Ferrier G, Evans C, Andrew J. 2002.** Synergistic effect of particles and cyclic pressure on cytokine production in human monocyte/macrophages: proposed role in periprosthetic osteolysis. *Bone* **30**(1):171–177.
- Minematsu H, Shin MJ, Celil Aydemir AB, Kim K-O, Nizami SA, Chung G-J, Lee FY-I. 2011.** Nuclear presence of nuclear factor of activated T cells (NFAT) c3 and c4 is required for Toll-like receptor-activated innate inflammatory response of monocytes/macrophages. *Cellular Signalling* **23**(11):1785–1793.
- Monneret G, Finck M-E, Venet F, Debard A-L, Bohé J, Bienvenu J, Lepape A. 2004.** The anti-inflammatory response dominates after septic shock: association of low monocyte HLA-DR expression and high interleukin-10 concentration. *Immunology Letters* **95**(2):193–198.
- Morgan RG, Ridsdale J, Payne M, Heesom KJ, Wilson MC, Davidson A, Greenhough A, Davies S, Williams AC, Blair A, Waterman ML, Tonks A, Darley RL. 2019.** LEF-1 drives aberrant β -catenin nuclear localization in myeloid leukemia cells. **104**(7):1365–1377 DOI [10.3324/haematol.2018.202846](https://doi.org/10.3324/haematol.2018.202846).
- Mosser DM, Edwards JP. 2008.** Exploring the full spectrum of macrophage activation. *Nature Reviews Immunology* **8**(12):958–969.
- Neme A, Seuter S, Carlberg C. 2016.** Vitamin D-dependent chromatin association of CTCF in human monocytes. *Biochimica Et Biophysica Acta (BBA) - Gene Regulatory Mechanisms* **1859**(11):1380–1388.
- Nikukar H, Reid S, Tsimbouri PM, Riehle MO, Curtis ASG, Dalby MJ. 2013.** Osteogenesis of Mesenchymal Stem Cells by Nanoscale Mechanotransduction. *ACS Nano* **7**(3):2758–2767 DOI [10.1021/nn400202j](https://doi.org/10.1021/nn400202j).
- Nurminen V, Neme A, Ryyänen J, Heikkinen S, Seuter S, Carlberg C. 2015.** The transcriptional regulator BCL6 participates in the secondary gene regulatory response to vitamin D. *Biochimica Et Biophysica Acta (BBA) - Gene Regulatory Mechanisms* **1849**(3):300–308.
- Nurminen V, Neme A, Seuter S, Carlberg C. 2019.** Modulation of vitamin D signaling by the pioneer factor CEBPA. *Biochimica Et Biophysica Acta (BBA) - Gene Regulatory Mechanisms* **1862**(1):96–106.

- Nurminen V, Seuter S, Carlberg C. 2019. Primary vitamin D target genes of human monocytes. *Frontiers in Physiology* 10:194 DOI 10.3389/fphys.2019.00194.
- Patel NR, Bole M, Chen C, Hardin CC, Kho AT, Mih J, Deng L, Butler J, Tschumperlin D, Fredberg JJ, Krishnan R, Koziel H. 2012. Cell elasticity determines macrophage function. *PLOS ONE* 7(9):e41024 DOI 10.1371/journal.pone.0041024.
- Pennathur S, Pasichnyk K, Bahrami NM, Zeng L, Febbraio M, Yamaguchi I, Okamura DM. 2015. The macrophage phagocytic receptor CD36 Promotes fibrogenic pathways on removal of apoptotic cells during chronic kidney injury. *The American Journal of Pathology* 185(8):2232–2245.
- Phillip Koeffler H, Reichel H, Bishop JE, Norman AW. 1985. γ -interferon stimulates production of 1, 25-dihydroxyvitamin D₃ by normal human macrophages. *Biochemical and Biophysical Research Communications* 127(2):596–603.
- Riberdy JM, Newcomb JR, Surman MJ, Barbosat JA, Cresswell P. 1992. HLA-DR molecules from an antigen-processing mutant cell line are associated with invariant chain peptides. *Nature* 360(6403):474–477.
- Robertson SN, Campsie P, Childs PG, Madsen F, Donnelly H, Henriquez FL, Mackay WG, Salmerón-Sánchez M, Tsimbouri MP, Williams C, Dalby MJ, Reid S. 2018. Control of cell behaviour through nanovibrational stimulation: nanokicking. *Philosophical Transactions of the Royal Society a: Mathematical, Physical and Engineering* 376(2120):20170290 DOI 10.1098/rsta.2017.0290.
- Schwende H, Fitzke E, Ambs P, Dieter P. 1996. Differences in the state of differentiation of THP-1 cells induced by phorbol ester and 1, 25-dihydroxyvitamin D₃. *Journal of Leukocyte Biology* 59(4):555–561 DOI 10.1002/jlb.59.4.555.
- Seuter S, Neme A, Carlberg C. 2016. Epigenome-wide effects of vitamin D and their impact on the transcriptome of human monocytes involve CTCF. *Nucleic Acids Research* 44(9):4090–4104 DOI 10.1093/nar/gkv1519.
- Seuter S, Neme A, Carlberg C. 2017. Epigenomic PU.1-VDR crosstalk modulates vitamin D signaling. *Biochimica Et Biophysica Acta (BBA) - Gene Regulatory Mechanisms* 1860(4):405–415.
- Seuter S, Neme A, Carlberg C. 2018. ETS transcription factor family member GABPA contributes to vitamin D receptor target gene regulation. *The Journal of steroid biochemistry and molecular biology* 177:46–52 DOI 10.1016/j.jsbmb.2017.08.006.
- Shao L, Sperber K. 2002. Impaired regulation of HLA-DR expression in human immunodeficiency virus-infected monocytes. *Clinical and Vaccine Immunology* 9(4):739–746.
- Shi C, Pamer EG. 2011. Monocyte recruitment during infection and inflammation. *Nature Reviews Immunology* 11(11):762–774.
- Sloan VS, Cameron P, Porter G, Gammon M, Amaya M, Mellins E, Zaller DM. 1995. Mediation by HLA-DM of dissociation of peptides from HLA-DR. *Nature* 375(6534):802–806.
- Solis AG, Bielecki P, Steach HR, Sharma L, Harman CCD, Yun S, De Zoete MR, Warnock JN, To SDF, York AG, Mack M, Schwartz MA, Dela Cruz CS, Palm

- NW, Jackson R, Flavell RA. 2019. Mechanosensation of cyclical force by PIEZO1 is essential for innate immunity. *Nature* 573(7772):69–74.
- Thiele A, Wasner M, Müller C, Engeland K, Hauschildt S. 2001. Regulation and possible function of β -catenin in human monocytes. *The Journal of Immunology* 167(12):6786–6793 DOI 10.4049/jimmunol.167.12.6786.
- Tickenbrock L. 2006. Wnt signaling regulates transendothelial migration of monocytes. *Journal of Leukocyte Biology* 79(6):1306–1313 DOI 10.1189/jlb.0905539.
- Tokuda N, Levy RB. 1996. 1, 25-Dihydroxyvitamin D₃ stimulates phagocytosis but suppresses HLA-DR and CD13 antigen expression in human mononuclear phagocytes. *Experimental Biology and Medicine* 211(3):244–250 DOI 10.3181/00379727-211-43967.
- Tsimbouri P. 2015. Adult stem cell responses to nanostimuli. *Journal of Functional Biomaterials* 6(3):598–622.
- Tsuchiya S, Gota Y, Okumura H, Nakae S, Konno T, Tada K, Kobayashi Y. 1982. Induction of Maturation in Cultured Human Monocytic Leukemia Cells by a Phorbol Diester. *Cancer Research* 42(4):1530–1536.
- Ulbricht T, Alzrigat M, Horch A, Reuter N, Mikecz Avon, Steimle V, Schmitt E, Krämer OH, Stamminger T, Hemmerich P. 2012. PML promotes MHC class II gene expression by stabilizing the class II transactivator. *The Journal of Cell Biology* 199(1):49–63.
- Wöbke TK, Sorg BL, Steinhilber D. 2014. Vitamin D in inflammatory diseases. *Frontiers in Physiology* 5:244 DOI 10.3389/fphys.2014.00244.
- Woo M-S, Yang J, Beltran C, Cho S. 2016. Cell surface CD36 protein in monocyte/macrophage contributes to phagocytosis during the resolution phase of ischemic stroke in mice. *Journal of Biological Chemistry* 291(45):23654–23661 DOI 10.1074/jbc.M116.750018.
- Yamamoto K, Ikeda U, Shimada K. 2003. Role of mechanical stress in monocytes / macrophages: implications for Atherosclerosis. *Current Vascular Pharmacology* 1(3):315–319.
- Yang C-Y, Leung PSC, Adamopoulos IE, Gershwin ME. 2013. The implication of vitamin D and autoimmunity: a comprehensive review. *Clinical Reviews in Allergy & Immunology* 45(2):217–226 DOI 10.1007/s12016-013-8361-3.
- Yang M, Ma B, Shao H, Clark AM, Wells A. 2016. Macrophage phenotypic subtypes diametrically regulate epithelial-mesenchymal plasticity in breast cancer cells. *BMC Cancer* 16(1):419 DOI 10.1186/s12885-016-2411-1.
- Zanoni I, Ostuni R, Marek LR, Barresi S, Barbalat R, Barton GM, Granucci F, Kagan JC. 2011. CD14 Controls the LPS-Induced Endocytosis of Toll-like Receptor 4. *Cell* 147(4):868–880.
- Zhang DE, Hetherington CJ, Gonzalez DA, Chen HM, Tenen DG. 1994. Regulation of CD14 expression during monocytic differentiation induced with 1 alpha, 25-dihydroxyvitamin D₃. *Journal of Immunology* 153(7):32766–32784.

Zhao QL, W jH, H L. 2016. Role of nuclear factor of activated T cells-2 in high mobility protein box-1 release in human monocytic THP-1 cells in vitro. *Journal of Southern Medical University (Chinese)* **36(1)**:8–12.

Zhong M, Komarova Y, Rehman J, Malik AB. 2018. Mechanosensing Piezo channels in tissue homeostasis including their role in lungs. *Pulmonary Circulation* **8(2)**:204589401876739 DOI [10.1177/2045894018767393](https://doi.org/10.1177/2045894018767393).