



UWS Academic Portal

The validity and reliability of a novel mobile app to measure agility performance in the physically active youth population

Escobar Álvarez, Juan A.; Carrasco-Zahinos, Rocío; Olivares, Pedro R.; Feu, Sebastián; Ramírez-Velez, Robinson; Pérez Sousa, Miguel Ángel

Published in:
European Journal of Human Movement

DOI:
[10.21134/eurjhm.2020.45.9](https://doi.org/10.21134/eurjhm.2020.45.9)

Published: 31/12/2020

Document Version
Publisher's PDF, also known as Version of record

[Link to publication on the UWS Academic Portal](#)

Citation for published version (APA):
Escobar Álvarez, J. A., Carrasco-Zahinos, R., Olivares, P. R., Feu, S., Ramírez-Velez, R., & Pérez Sousa, M. Á. (2020). The validity and reliability of a novel mobile app to measure agility performance in the physically active youth population. *European Journal of Human Movement*, 45, [9]. <https://doi.org/10.21134/eurjhm.2020.45.9>

General rights

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact pure@uws.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Original research

The validity and reliability of a novel mobile app to measure agility performance in the physically active youth population

Juan A. Escobar-Alvarez¹, Rocío Carrasco-Zahinos², Pedro R. Olivares³, Sebastián Feu⁴,
Robinson Ramírez-Vélez⁵, Miguel Angel Pérez-Sousa^{6*}

1. Faculty of Higher Education, South Essex College University Centre, Southend on the Sea, UK
2. Facultad de Educación, Universidad de Extremadura, 06006 Badajoz, Spain
3. Facultad de Educación, Psicología y Ciencias del Deporte, Universidad de Huelva, Spain, Instituto de Actividad Física y Salud, Universidad Autónoma de Chile, Chile
4. Facultad de Educación, Universidad de Extremadura, 06006 Badajoz, Spain
5. Navarrabiomed-Universidad Pública de Navarra (UPNA), Complejo Hospitalario de Navarra (CHN), Instituto de Investigación Sanitaria de Navarra (IdiSNA), Pamplona, Navarra, Spain; CIBER of Frailty and Healthy Aging (CIBERFES), Instituto de Salud Carlos III, Pamplona, Navarra, Spain
6. Faculty of Sport Sciences, University of Extremadura, Cáceres, Spain; Epidemiology of Physical Activity and Fitness Across Lifespan research group, University of Seville, Seville, Spain

*Correspondence: (Miguel A. Pérez-Sousa) perezsousa@gmail.com

Received: 24/08/2020; Accepted: 16/12/2020; Published: 31/12/2020

Abstract: Agility is a key component of physical fitness in adolescents. However, the measurement of this variable is usually complex, requiring high cost instruments and complex software. To test the validity and reliability of a novel iPhone app (Lap Tracker Auto-timer) to measure agility performance among adolescents. Twenty-four physically active adolescents (15.7 ± 2.3 years old) participated in two testing sessions (separated by 7 days). They performed three 4×10 m agility test trials measured by Photocell or the iPhone app. The correlation analysis revealed high validity ($r = .92$; 95% confidence interval [CI] = $.88 - .95$), with a standard error of the estimate of 0.56 s ($p < 0.001$). The coefficient of variation (CV; 0.09) and intraclass correlation coefficient (ICC; $.93$; 95% CI = $.85 - .97$) showed an acceptable reliability. This study demonstrated that the iPhone App Lap Tracker Auto-timer could be a valid, reliable and low-cost tool to evaluate agility performance in adolescents. However, more studies are required to guarantee the utility of this app.

Keywords: app, valid, reliable, performance, agility

1. Introduction

Agility can be defined as the ability to perform a rapid whole-body movement given a change in velocity or direction in response to a stimulus (Sheppard & Young, 2006). This ability is integrated in athletic performance, especially in team sports,

where continuous accelerations, decelerations, and change of direction (COD) or speed are required during a match. For example, agility is suggested to be a key performance indicator in soccer (Lloyd et al. 2015), basketball (Scanlan, Tucker, and Dalbo 2014) and rugby (Freitas et al. 2018). In fact, agility is one of the components of health-related physical fitness, and several studies



have shown that agility contributes to better health (Ortega et al. 2008) and quality of life (Perez-Sousa et al. 2019).

Agility is widely examined using several protocols (Ruiz et al., 2011; Sheppard et al., 2014). The aforementioned evaluations are characterised by having to complete a trajectory of previously planned directional changes as fast as possible. These timed events can be measured by several tools, including stopwatches (Ruiz et al. 2011), radar guns (Samozino et al. 2016) and mobile apps. However, the gold standard method within this field are the photoelectric timer gates devices (Samozino et al. 2016; Vicente-Rodríguez et al. 2011). Most of these tests provide more accurate and reliable (intraclass correlation coefficient [ICC] > .70) information when they are performed with high-cost devices that are of limited use under laboratory conditions. In addition, these instruments usually require specific and complex software to interpret the obtained data.

In the past 5–10 years, there has been increasing research on the use of mobile phones (including smartphones) for measuring athletic performance (Peart, Balsalobre-Fernández, and Shaw 2017). The use of this wearable and low-cost technology facilitates the analysis of human movement for several professionals in this field (physical education teachers, coaches, physicians, physiotherapists, researchers, etc.). Mobile applications also offer a potential solution to provide support for sports settings in general because mobile phones are commonly accessible (Muntaner-Mas et al. 2019).

Although not all mobile apps have been subjected to empirical testing, previous research have analysed the validity and reliability of commercial apps for the measurement of physical parameters, including vertical jump in different populations (Balsalobre-Fernández et al. 2015; Cruvinel-Cabral et al. 2018), sprinting ability (Romero-Franco et al. 2017; Stanton et al. 2016) and more recently COD (Balsalobre-

Fernández et al. 2019). Therefore, further studies are warranted to determine the utility of commercial apps for objectively and remotely assess agility performance. Thus, this study aimed to determine the validity and reliability of an application for iOS (Lap Tracker Auto-timer) to measure agility performance in physically active adolescents.

2. Materials and Methods

Subjects

Twenty-four physically active participants volunteered for this study (18 males and 4 females). Participants' characteristics were: 14–18 years old (15.7 ± 2.3 years), 163.9 ± 6.4 cm tall and 64.4 ± 6.5 kg. Criteria to be included were: i) did not suffer from any disease or injury that would limit their performance; ii) between 14 and 18 years; iii) provided informed consent. Also a question of physical activity was conducted as follow: (i) "Have you regularly exercised, such as jogging or dancing, or performed rigorous physical activity as football, hockey...etc. at least three times a week?". Participants were considered physically active if they responded affirmatively. Additionally, written informed consent was obtained from their parents or legal tutors after being informed of the benefits and risks of the research. This study was approved by the ethics board at South Essex College (ID 3526AL) and performed in agreement with the Declaration of Helsinki.

Methodology

Kit Racetime2 Light Radio (Micro Gate) - Kit Racetime 2 (Microgate, Bolzano, Italy) are photocells that transmits data via radio. It is designed to measure total or intermediate times accurately and reliably. Previous studies have considered this device to be the 'gold standard' for measuring times and split times in different sports performance test, such as speed/agility, course ship, sprints, etc (Bond et al. 2017; García López et al. 2012). The photocells were aligned with the reflectors, following the manufacturer's instructions, during each participant's trial.

Lap Tracker Auto-timer app - Lap Tracker Auto-timer (version 4.0) is a motion-detection-based lap timer that can be used to track different athletes in distinct tests performance. The camera view has up to four adjustable detection zones (see Figure 1) that detect the movement when an object or person passes in front of the camera of the mobile device where it is installed. For this study, the app was used on an iPhone® 8 that was running iOS 12.2 (Apple Inc., USA), although it is also available for iPad and iPod. The app was configured to stop on the fourth round of the test, and the data were recorded on an Excel sheet for its subsequent analysis.

Testing protocol - All the participants were asked to meet in the morning of the study. Body mass was measured using a scale (Seca model 711, Hamburg, Germany), and height was estimated with an aluminium stadiometer (Seca model 220, Postfach, Germany), according to the recommended procedures in the body composition assessment protocol (Norton et al. 1996). Participants performed a standardised warm-up that comprised 8 min of moderate jogging and 5 min of dynamic stretching. At the end of the warm-up and after explaining the test, each subject performed two familiarisation during the testing day. No physical limitations or musculoskeletal injuries that could affect testing were reported.

The 4 x 10 m shuttle run test has been used for assessing speed/agility and coordination in previous studies (Artero et al. 2009; Moliner-Urdiales et al. 2011). It is especially recommended for children and adolescents (Ruiz et al. 2011). Two parallel lines were marked on a slip-proof floor (10 m apart) with a yellow-coloured spray, and three sponge blocks were placed behind the marked lines (one behind the starting line and two behind the opposite line). Participants were instructed to run back and forth as fast as possible and cross each line with



both feet. Each time the participants crossed any of the lines, they had to pick up (the first time) or exchange (second and third time) a sponge block. The app and the photocells were stopped when the adolescent crossed the end line and thus completed the total 40 m (4 x 10 m). The iPhone 8 was placed on a fixed tripod (same position as the photocells) and aligned with the beam of the photocell with a laser to ensure that both instruments automatically collected the data at the same time (Figure 2).




Figure 2. Representation of the 4 x 10 m shuttle run test. The camera phone was placed just below of Photocell and the detection zone configured to detect movement to the height of photocell receptor

The aforementioned protocol was performed three times (with 7 days between each trial). For the first trial, the times for the agility performance test were collected with both devices to observe the validity of the app in comparison with the photocells. For the second and third trials, only the app was used, in order to analyse the absolute and relative reliability of it. Participants always performed the task at the same time of day, a consideration that controlled for potential biological-contaminating variables (e.g., circadian rhythms) that can influence the results (Valdez-Ramírez et al. 2009).

Statistical Analysis

Descriptive analyses were performed for the dataset obtained with the photocells and the app. The Shapiro-Wilk test was used to examine whether each analysed variable was normally distributed. The course of normality and the capacity improvement process of Box-Cox (Nachtsheim et al. 2004) were also examined to ensure data uniformity and the ability to use parametric statistical tests.

The validity of the app was analysed using a Pearson correlation with the data obtained for each device. The chosen criterion to interpret the magnitude of the correlation (r) was: ≤ 0.1 = trivial, $> .1$ to $.3$ = small, $> .3$ to $.5$ = moderate, $> .5$ to $.7$ = large, $> .7$ to $.9$ = very large, $> .9$ to 1.0 = almost perfect (Hopkins et al., 2009).

The relative and absolute reliability, besides the magnitude of the change was analysed using Excel (with a datasheet provided by Hopkins) for validity and reliability studies (Hopkins, 2015). The relative reliability was assessed using ICC with 95% confidence interval (CI). An ICC $\geq .70$ was considered to be acceptable (Koo and Li 2016; Weir 2005). To examine the absolute reliability, we first calculated the change in the mean between sets and the typical error of the measure expressed as coefficient of variation (CV) (in per cent). A CV $< 5\%$ was set as the reliability criterion (Hopkins et al., 2009). Finally, the magnitude of the change between tests was calculated using Cohen's effect size (ES). The criterion for interpreting this magnitude was

< 0.2 = trivial, $0.2-0.6$ = small, $0.6-1.2$ = moderate and > 1.2 = large (Hopkins et al., 2009). Reliability was also assessed according to the Bland-Altman method. This analysis measures bias as estimated from mean differences, the 95% confidence interval for bias, the limits of agreement and ± 1.96 standard deviation (SD) of the difference. All data were analyzed using MedCalc Statistical Software version 18.2 (MedCalc Software BVBA, Ostend, Belgium). Figures were designed using GraphPad Prism 6.0 (GraphPad Software Inc., California, USA).

3. Results

Mean \pm standard deviation values for time performance (s) for the 4×10 m test with the photocells and the Lap Tracker Auto-timer app are shown in Table 1. Pearson correlation analysis is also shown in Table 1. There was a very large correlation ($r = .92$; $p < .001$) between the devices.

The ICC was .93, a value that indicated almost perfect reliability between test-retest, while the CV for the absolute reliability analysis was 0.09%. Finally, the magnitude of the test-retest change provided a Cohen's ES of 0.1, a value indicative of a trivial effect (Table 2).

Finally, the Bland-Altman plots are depicted in Figure 3. Graphical analysis showed a noticeable and similar concordance between the photocells and app for average time measurements, as shown by the low bias value (SD of bias = 0.30) and narrow confidence interval (range: -0.53 to 0.63%).

Table 1: Mean \pm standard deviation and correlation analysis of the 4×10 m shuttle run test performance (s) for the photocells and the app *Lap Tracker Auto-timer*.

Photocells	App	r	95% CI	SEE	R^2	P	Inference
11.2 \pm 0.6	11.3 \pm 0.7	.92	(.88, .95)	0.56	.86	< .001	Very large

Note: 95% Confidence interval. SEE : standard error estimate




Figure 3. Correlation plot between Photocell and the app Lap tracker Auto-timer for the measurement of the total time in 4x10 shuttle run test.

Table 2. Mean \pm standard deviation values for test-retest, intraclass correlation (ICC), coefficient of variability (CV) and effect size (ES).

Test	Retest	ICC	95% CI	%CV	95% CI	ES	95% CI	Inference
10.7 \pm 0.9	10.6 \pm 0.9	0.93	(0.85 – 0.97)	0.09	(0.0 – 1.4)	0.1	(0.69 – 0.49)	Trivial

Note: 95% Confidence interval.

4. Discussion

The purpose of this study was to examine the validity and reliability of the mobile app *Lap Tracker Auto-timer* for the measurement of agility performance. Our results showed that the app provided valid and reproducible data when compared with the photocells. The data obtained during the validity assessment process reflected a very large correlation ($r = .92$, $p < .001$) between the times obtained with both devices (Hopkins et al., 2009). Previous research have suggested that when seeking to replace an existing measurement instrument on the other, due to easy handling or lower costs, the correlation index must be greater than 0.8 (Lamprea and Gómez-Restrepo 2007), as was the case of this study. Regarding the reliability of the app, the results obtained for the test-retest display almost perfect reliability (ICC = .93) (Koo and Li 2016), with a coefficient of variation of 0.09 and a trivial ES (0.1) for the test-retest comparison (Hopkins et al., 2009). Therefore, the aforementioned results suggested that the app provides reliable data.

To the best of our knowledge, this study is the first to observe the validity and reliability of the app *Lap Tracker Auto-timer* for testing agility performance. This app differs from others for testing physical

parameters due to the use of automatic detection of movement. While other valid and reliable apps are based on video-analysis, they require importing and manipulating high-speed videos of the participant's performance by manual selection of the appropriated frame (Balsalobre-Fernández et al. 2015; Romero-Franco et al. 2017). In recent years, different smartphone apps have been developed to collect several types of vertical jumps (countermovement jump, squat jump, etc.) (Balsalobre-Fernández et al. 2015), sprint performance (Romero-Franco et al. 2017) and COD (Balsalobre-Fernández et al. 2019),

among others (Peart et al. 2017). A recent study by study by Balsalobre-Fernández et al (2019) reported good validity and reliability for a smartphone app for the measurement of COD performance, although they used the 5+5 test (Nimphius et al. 2018) while our results are based on the 4 x 10 m shuttle run test (Ruiz et al. 2011).

Considering the rise of smartphone technology for the measurement of sport performance and physical testing (Peart et al. 2017), this study provides relevant information about the accuracy of an app with a novel design. This app could be used by sport scientists, researchers, strength and conditioning coaches or even practitioners to measure performance without post-video-analysis. To optimise the use of this app and obtain more accurate data, we recommend: i) using a tripod, ii) the height of the device (iPhone, iPad or iPod) should be between 80 cm and 1 m (to ensure the mobile camera captures the area from the hip to the chest) and iii) the space where the trigger point is very sensitive to movement and therefore must be free of elements.

There are several limitations to this study that must be discussed. First, the sample size was small. Future studies involving larger samples are needed. Another limitation of this study was

experienced during data collection. The camera is sensitive to any movement within the image frame. In this way, if the app detects movement, the timer starts its account, so it is necessary to limit this image frame as much as possible. Furthermore, *Lap Tracker Auto-timer* can be used for some other tests that measure speed/agility as 5-0-5, Illinois (Sheppard & Young, 2006), Arrowhead, 5-10-5 and T-tests (Walker 2016), besides tests where the participants starts from a point to which they must return to finish it.

In conclusion, our findings suggest that the app *Lap Tracker Auto-timer* for iOS devices could be a valid, reliable and low-cost alternative to measure speed/agility in the 4 x 10 m test. In particular, we identified empirical validation that the *Lap Tracker Auto-timer* offers an acceptable remotely assess speed/agility as a complementary tool to traditional methods in the fitness and sports settings.

Funding: Non-declared

Acknowledgments: The authors would like to thank the participants for their involvement and interest in the present study.

Conflicts of Interest: The authors declare no affiliations with or involvement in any organisation or entity with any financial interest or non-financial interest in the subject matter discussed in this manuscript. No potential conflict of interest was reported by the authors.

References

- Artero, E. G., V. España-Romero, F. B. Ortega, D. Jiménez-Pavón, J. R. Ruiz, G. Vicente-Rodríguez, M. Bueno, A. Marcos, S. Gómez-Martínez, A. Urzanqui, M. González-Gross, L. A. Moreno, A. Gutiérrez, and M. J. Castillo. 2009. "Health-Related Fitness in Adolescents: Underweight, and Not Only Overweight, as an Influencing Factor. The AVENA Study." *Scandinavian Journal of Medicine & Science in Sports* 20(3):418–27.
- Balsalobre-Fernández, Carlos, Chris Bishop, José Vicente Beltrán-Garrido, Pau Cecilia-Gallego, Aleix Cuenca-Amigó, Daniel Romero-Rodríguez, and Marc Madruga-Parera. 2019. "The Validity and Reliability of a Novel App for the Measurement of Change of Direction Performance." *Journal of Sports Sciences* 1–5.
- Balsalobre-Fernández, Carlos, Mark Glaister, Richard Anthony Lockey, C. Balsalobre-Fernandez, Mark Glaister, and Richard Anthony Lockey. 2015. "The Validity and Reliability of an iPhone App for Measuring Vertical Jump Performance." *Journal of Sports Sciences* 33(15):1574–79.
- Bond, Colin W., Emily M. Willaert, Kyle E. Rudningen, and Benjamin C. Noonan. 2017. "Reliability of Three Timing Systems Used to Time Short on Ice-Skating Sprints in Ice Hockey Players." *Journal of Strength and Conditioning Research* 31(12):3279–86.
- Cruvinel-Cabral, Rejane Maria, Iransé Oliveira-Silva, André Ricarte Medeiros, João Gustavo Claudino, Pedro Jiménez-Reyes, and Daniel A. Boullosa. 2018. "The Validity and Reliability of the 'My Jump App' for Measuring Jump Height of the Elderly." *PeerJ*.
- Freitas, Tomás, Pedro Alcaraz, Chris Bishop, Julio Calleja-González, Ademir Arruda, Aristide Guerriero, Valter Reis, Lucas Pereira, and Irineu Loturco. 2018. "Change of Direction Deficit in National Team Rugby Union Players: Is There an Influence of Playing Position?" *Sports* 7(1):2.
- García López, J., Juan Carlos Morante Rábago, Ana Ogueta-Alday, Javier González Lázaro, José Antonio Rodríguez Marroyo, and José G. Villa Vicente. 2012. "El Uso de Fotocélulas de Haz Simple y Doble Para Medir La Velocidad En Carreras®. The Use of Single- and Dual-Beam Photocells to Measure the Sprint Time®." *RICYDE. Revista Internacional de Ciencias Del Deporte*. 8(30):324–33.
- Hopkins, W. G. 2015. "Spreadsheets for Analysis of Validity and Reliability." *Sportscience* 19:36–42.
- Hopkins, William G., Stephen W. Marshall, Alan M. Batterham, and Juri Hanin. 2009. "Progressive Statistics for Studies in Sports Medicine and Exercise Science." *Medicine and Science in Sports and Exercise* 41(1):3–12.
- Koo, Terry K., and Mae Y. Li. 2016. "A Guideline of Selecting and Reporting Intraclass Correlation Coefficients for Reliability Research." *Journal of Chiropractic Medicine* 15(2):155–63.
- Lamprea, J. A., and C. Gómez-Restrepo. 2007. "Validez En La Evaluación de Escalas." *Revista Colombiana de Psiquiatría* 36(2):340–48.
- Lloyd, Rhodri S., Jon L. Oliver, John M. Radnor, Benjamin C. Rhodes, Avery D. Faigenbaum, and Gregory D. Myer. 2015. "Relationships between Functional Movement Screen Scores, Maturation and Physical Performance in Young Soccer Players." *Journal of Sports*

- Sciences 33(1):11–19.
- Moliner-Urdiales, D., J. R. Ruiz, G. Vicente-Rodriguez, F. B. Ortega, J. P. Rey-Lopez, V. Espana-Romero, J. A. Casajus, D. Molnar, K. Widhalm, J. Dallongeville, M. Gonzalez-Gross, M. J. Castillo, M. Sjostrom, L. A. Moreno, and HELENA Study Group. 2011. "Associations of Muscular and Cardiorespiratory Fitness with Total and Central Body Fat in Adolescents: The HELENA Study." *British Journal of Sports Medicine* 45(2):101–8.
- Muntaner-Mas, Adrià, Antonio Martinez-Nicolas, Carl J. Lavie, Steven N. Blair, Robert Ross, Ross Arena, and Francisco B. Ortega. 2019. "A Systematic Review of Fitness Apps and Their Potential Clinical and Sports Utility for Objective and Remote Assessment of Cardiorespiratory Fitness." *Sports Medicine* 49(4):587–600.
- Nachtsheim, Christopher J., John Neter, William Li, and Michael H. Kutner. 2004. "Applied Linear Statistical Models." *Journal Of The Royal Statistical Society Series A General*.
- Nimphius, Sophia, Samuel J. Callaghan, Neil E. Bezodis, and Robert G. Lockie. 2018. "Change of Direction and Agility Tests: Challenging Our Current Measures of Performance." *Strength and Conditioning Journal*.
- Norton, K., N. Whittingham, J. Carter, D. Kerr, C. Gore, and M. Marfell-Jones. 1996. *Anthropometrica: A Textbook of Body Measurement for Sports and Health Courses*. Sydney: UNSW Press.
- Ortega, F. B., J. R. Ruiz, M. J. Castillo, and M. Sjöström. 2008. "Physical Fitness in Childhood and Adolescence: A Powerful Marker of Health." *International Journal of Obesity* 32(1):1–11.
- Peart, Daniel J., Carlos Balsalobre-Fernández, and Matthew P. Shaw. 2017. "The Use of Mobile Applications to Collect Data in Sport, Health and Exercise Science." *Journal of Strength and Conditioning Research*.
- Perez-Sousa, Miguel A., Pedro R. Olivares, Antonio Garcia-Hermoso, and Narcis Gusi. 2019. "Fitness as a Mediator of the Enhancement of Quality of Life after a 6-Months Exercise Program." *Research Quarterly for Exercise and Sport* 1–10.
- Romero-Franco, Natalia, Pedro Jiménez-Reyes, Adrián Castaño-Zambudio, Fernando Capelo-Ramírez, Juan José Rodríguez-Juan, Jorge González-Hernández, Francisco Javier Toscano-Bendala, Víctor Cuadrado-Peñafiel, and Carlos Balsalobre-Fernández. 2017. "Sprint Performance and Mechanical Outputs Computed with an iPhone App: Comparison with Existing Reference Methods." *European Journal of Sport Science* 17(4):386–92.
- Ruiz, J. R., J. Castro-Pinero, V. Espana-Romero, E. G. Artero, F. B. Ortega, M. M. Cuenca, D. Jimenez-Pavon, P. Chillón, M. J. Girela-Rejon, J. Mora, A. Gutierrez, J. Suni, M. Sjostrom, and M. J. Castillo. 2011. "Field-Based Fitness Assessment in Young People: The ALPHA Health-Related Fitness Test Battery for Children and Adolescents." *British Journal of Sports Medicine* 45(6):518–24.
- Samozino, P., G. Rabita, S. Dorel, J. Slawinski, N. Peyrot, E. Saez de Villarreal, and J. B. Morin. 2016. "A Simple Method for Measuring Power, Force, Velocity Properties, and Mechanical Effectiveness in Sprint Running." *Scandinavian Journal of Medicine & Science in Sports* 26(6):648–58.
- Scanlan, Aaron T., Patrick S. Tucker, and Vincent J. Dalbo. 2014. "A Comparison of Linear Speed, Closed-Skill Agility, and Open-Skill Agility Qualities Between Backcourt and Frontcourt Adult Semiprofessional Male Basketball Players." *Journal of Strength and Conditioning Research* 28(5):1319–27.
- Sheppard, J., and W. Young. 2006. "Agility Literature Review: Classifications, Training and Testing." *Journal of Sports Sciences* 24(9):919–32.
- Sheppard, Jeremy, J. Jay Dawes, Ian Jeffreys, Tania Spiteri, and Sophia Nimphius. 2014. "Broadening The View Of Agility: A Scientific Review Of The Literature." *ECU Publications Post 2013*.
- Stanton, Robert, Melanie Hayman, Nyree Humphris, Hanna Borgelt, Jordan Fox, Luke Del Vecchio, and Brendan Humphries. 2016. "Validity of a Smartphone-Based Application for Determining Sprinting Performance." *Journal of Sports Medicine (Hindawi Publishing Corporation)* 2016:7476820.
- Valdez-Ramírez, Pablo, Candelaria Ramírez-Tule, Aída García-García, and Javier Talamantes-López. 2009. "Ritmos Circadianos En La Eficiencia Para Responder En Una Prueba de Ejecución Continua." *Revista Mexicana de Análisis de La Conducta* 35(1):75–91.
- Vicente-Rodríguez, Germán, Juan P. Rey-López, Jonathan R. Ruíz, David Jiménez-Pavón, Patrick Bergman, Donatella Ciarapica, Jose M. Heredia, Denes Molnar, Angel Gutierrez, Luis A. Moreno, Francisco B. Ortega, and HELENA Study Group. 2011. "Interrater Reliability and Time Measurement Validity of Speed–Agility Field Tests in Adolescents." *Journal of Strength and Conditioning Research* 25(7):2059–63.

Walker, Owen. 2016. "Agilidad | Ciencia Para El Deporte." Science for Sport.
Weir, J. P. 2005. "Quantifying Test-Retest

Reliability Using the Intraclass Correlation Coefficient and the SEM." J Strength Cond Res 19:231–40.