



## UWS Academic Portal

### **Effect of supplementary physical training on vertical jump height in professional ballet dancers**

Escobar-Álvarez, Juan A.; Jiménez-Reyes, Pedro; Da Conceição, Filipe A.; Fuentes-García, Juan P.

*Published in:*  
International Journal of Sports Physiology and Performance

*DOI:*  
[10.1123/ijsp.2021-0513](https://doi.org/10.1123/ijsp.2021-0513)

Published: 31/08/2022

*Document Version*  
Peer reviewed version

[Link to publication on the UWS Academic Portal](#)

*Citation for published version (APA):*

Escobar-Álvarez, J. A., Jiménez-Reyes, P., Da Conceição, F. A., & Fuentes-García, J. P. (2022). Effect of supplementary physical training on vertical jump height in professional ballet dancers. *International Journal of Sports Physiology and Performance*, 17(8), 1257-1263. <https://doi.org/10.1123/ijsp.2021-0513>

**General rights**

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

If you believe that this document breaches copyright please contact [pure@uws.ac.uk](mailto:pure@uws.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

Escobar-Álvarez, J. A., Jiménez-Reyes, P., Da Conceição, F. A., & Fuentes-García, J. P. (2022). Effect of supplementary physical training on vertical jump height in professional ballet dancers. *International Journal of Sports Physiology and Performance*, 17(8), 1257–1263.

Accepted author manuscript version reprinted, by permission, from International Journal of Sports Physiology and Performance, 2022, <https://doi.org/10.1123/ijsp.2021-0513> © Human Kinetics, Inc.



## Effect of supplementary physical training on vertical jump height in professional ballet dancers

Journal:	<i>International Journal of Sports Physiology and Performance</i>
Manuscript ID	IJSPP.2021-0513.R3
Manuscript Type:	Original Investigation
Date Submitted by the Author:	28-Apr-2022
Complete List of Authors:	Escobar-Alvarez, Juan; University of the West of Scotland - Ayr Campus, School of Health and Life Sciences; UP CIFI2D, CIFI2D-Center of Research, Education, Innovation and Intervention in Sport. University of Porto; South Essex College, Faculty of Higher Education Jiménez-Reyes, Pedro; Universidad Rey Juan Carlos, Centre for Sport Studies, Rey Juan Carlos University, Madrid, Spain Conceição, Filipe; UP LABIOMEPE, LABIOMEPE, Porto Biomechanics Laboratory. University of Porto; UP CIFI2D, CIFI2D-Center of Research, Education, Innovation and Intervention in Sport, Faculty of Sports. University of Porto Fuentes, Juan P.; Universidad de Extremadura, Facultad de Ciencias del Deporte
Keywords:	Plyometric training, Vertical jump, Dance performance, strength training, Sauté in first position

SCHOLARONE™  
Manuscripts

1 **Effect of supplementary physical training on vertical jump height in professional**  
2 **ballet dancers**

3  
4  
5  
6 **Running head:** Plyometric vs Combined training in ballet dancers  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

For Peer Review

51 **Abstract**

52

53 Dancers require many specific dance skills of a ballistic nature. Therefore, the design of  
54 supplementary training to improve the strength of the lower limbs and jump height is a  
55 relevant area of research. The purpose of this study was (i) to compare the effect of  
56 plyometric training (PT) vs combined training (CTr) on countermovement jump (CMJ),  
57 squat jump (SJ) and *sauté* in first position (*sauté*) height; and (ii) to observe whether  
58 changes on CMJ and SJ are associated with changes in *sauté* in female and male dancers.  
59 Eighty-one classical professional ballet dancers (41 females and 40 males,  
60 age=22.9±3.7years, body mass-BM=59.7±8.6Kg, height=167.4±7.3cm) from two  
61 different dance companies were divided into a control group and 2 experimental groups:  
62 PT and CTr. All groups followed their common routine of training regarding classes and  
63 rehearsal practice, while the experimental groups added 2 sessions (1hour per session) for  
64 9 weeks of supplementary training. Significant increases (medium to large effect size) in  
65 CMJ, SJ and *sauté* height were found in the pre- vs post-test comparisons for both  
66 experimental groups. Significant, very large correlations were found between the  
67 magnitude of improvement in *sauté* and the magnitude of improvement in CMJ and SJ.  
68 Plyometric and combined training programs are effective ways to improve jumping  
69 ability in professional female and male ballet dancers. The improvement in CMJ and SJ  
70 has a good transference on *sauté* performance. These findings support the use of  
71 traditional training methods to improve jump height in specific and **non** specific ballet  
72 jumping ability.

73

74 **Key Words:** Performance, Exercise Fitness, Strength, Training

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102 **INTRODUCTION**

103

104 Ballet is an artistic discipline that requires both male and female dancers to perform many  
105 unilateral and bilateral jumps. Therefore, jumping ability has been identified as an  
106 important part of dance performance, giving dancers a longer time to implement technical  
107 skills during the flight stage of the leap<sup>1-4</sup>. A previous study has compared the number  
108 of jumps in female (5.29+4.97 jumps per minute) and male (4.68+4.98 jumps per minute)  
109 ballet dancers, reporting significant differences between them<sup>5</sup>. The prescription of  
110 supplementary training aimed to improve jump height<sup>2,6-9</sup>, however, is not as  
111 commonplace in ballet as it is in sports.

112 It has been suggested that in ballet, male dancers are more focused on supplementary  
113 weight training, while females undertake more cardiovascular exercise<sup>7</sup>. One of the  
114 reasons for this, has been the assumption that dancers are concerned (particularly in  
115 females) about increasing muscle mass and its impact on the aesthetic component of  
116 dance. However, a recent study with a mixed genre of professional dancers (19 ballet  
117 dancers included) has concluded that the dancers' perception is to not be afraid of muscle  
118 hypertrophy, which is a more prevalent preconception in dance teachers<sup>10</sup>.

119

120 Due to the physical demands and physiological responses observed in professional ballet  
121<sup>11</sup>, designing optimal training programs for dancers is a relevant area of research. Recent  
122 investigations have suggested that due to the high number of jumps, *pliés*, and lifts,  
123 managing training loads and recovery may be a focus for strategies seeking to optimize  
124 dancer health and wellbeing<sup>11</sup>. Therefore, ballet companies need to provide dancers with  
125 supplementary physical training<sup>11</sup>. The implementation of training principles (i.e.  
126 periodization and progressive overload) would also benefit dancers' health and wellbeing  
127<sup>11,12</sup>. Different methods of training such as plyometrics (PT)<sup>2</sup>, whole-body vibration<sup>13-</sup>  
128<sup>16</sup>, traditional strength training<sup>2,13</sup> and individualized training based on force-velocity  
129 profiling during jumping<sup>17</sup> have reported a positive effect on jumping ability in elite  
130 ballet and modern dancers<sup>18,19</sup>. However, most of the studies have been conducted with  
131 female participants. A recent systematic review of the kinetic and kinematic parameters  
132 in ballet jumping concluded that only 4 out of 29 articles investigated male ballet dancers,  
133 identifying a gap in the literature<sup>19</sup>.

134

135 Dance choreography routinely demands several types of jumps, including standard  
136 actions such as countermovement jumps (CMJ) or squat jumps (SJ) and ballet-specific  
137 jumps such as the *sauté* in first position (*sauté*)<sup>20</sup>. *Sauté* is one of the most common ballet  
138 jumps, which can be defined as a CMJ that is performed with an external rotation of the  
139 lower limbs and the heels together, known as turnout<sup>21</sup>. Investigations have reported the  
140 differences between CMJ and SJ in comparison with *sauté*<sup>22,23</sup>. However, to the best of  
141 our knowledge, investigations have not observed the effect of supplementary training to  
142 improve jump height (CMJ and SJ), and its effect on a specific dance skill like the *sauté*.

143

144 Some other specific dance jumps have been studied. Blanco and colleagues found large  
145 to very large associations between *grand jeté* leap height and CMJ and drop jump in ballet  
146 dancers with different skill levels, suggesting a relationship between ballet specific and  
147 non-specific jumps in this population<sup>24</sup>. This may support the use of traditional training

148 methods such as plyometrics or combined training (Ctr) <sup>25</sup> to improve ballet specific  
149 jumps. Moreover, recent findings suggest that supplementary training can have positive  
150 outcomes on ballet specific jumps <sup>26</sup>. As a consequence, performance in specific dance  
151 jumps that are vertically orientated and with an external rotation of the lower limbs, such  
152 as *sauté*, may be enhanced through training programs based on improving CMJ or SJ.

153

154 The current literature reveals a lack of studies observing the effect of vertical jump  
155 training (CMJ and SJ) on specific dance skills based on leaping. In addition to this,  
156 investigations observing the effect of supplementary physical training on male dancers  
157 are very limited <sup>19</sup>. Therefore, the purpose of this study was (i) to compare the effect of  
158 two common methods of supplementary training (PT, CTr) on CMJ, SJ and *sauté* height;  
159 and (ii) to observe whether changes in CMJ and SJ were associated with changes in *sauté*  
160 after nine weeks of physical supplementary training in female and male dancers.

## 161 **METHODS**

### 162 **Subjects**

163 Eighty-one (41 females-F and 40 males-M) professional classical ballet dancers from 2  
164 different dance companies (38 and 43 dancers each) participated in this study  
165 (age=22.9±3.7years, body mass-BM=59.7±8.6Kg, height=167.4±7.3cm). All the  
166 participants had more than 12 years' experience (training volume of 13±3.5 hours per  
167 week) and were informed of the benefits and risks of the investigation through a  
168 structured consent form and a Physical Activity Readiness Questionnaire (PAR-Q). This  
169 study was approved by the Research Ethics Board of *removed for peer-reviewed* in  
170 agreement with the Declaration of Helsinki.

171

### 172 **Procedures**

173 All subjects were asked to meet for two sessions one week before any training  
174 intervention (Pre-test week). Participants performed the same standardized warm-up both  
175 days, consisting of 5 minutes' jogging, dynamic stretching, a range of movement  
176 exercises based on specific dance actions, and preparatory jumps in countermovement,  
177 squat, and *sauté* in first position (unilateral and bilateral jumps). Body mass (BM) in kg  
178 and height (cm) were measured using a Tanita SC-330 (TANITA Corporation, Itabashi-  
179 ku Tokyo, Japan), and an aluminium stadiometer (Seca 713 model, Postfach, Germany)  
180 respectively. Knee flexion was standardised to 90 degrees during the CMJ and SJ. CMJ  
181 height (cm), SJ height (cm) and *sauté* height (cm) were measured, selecting the best  
182 performance of 3 trials (2 min intra-set and 5 min for inter-set recovery) using a contact  
183 platform size A-1 (Chronojump Boscossystem®) <sup>27</sup>. The Chronojump contact platform  
184 calculates the jump height based on flight time. Data were inputted into Microsoft excel  
185 prior to statistical analysis.

186

187 First testing Session (Tuesday): The first session was planned to collect participants' BM,  
188 height and CMJ. All the participants performed the CMJ barefoot, and two observers (one  
189 on either side of the subject) provided feedback about the starting height of 90° (knee  
190 flexion) for the squat during jump performance. If one of the observers disagreed with  
191 the 90° (knee flexion) squat, the trial was invalidated and was repeated.

192

193 Second testing session (Friday): The second session was aimed to collect SJ height and  
194 *sauté* height. The methods for SJ were consistent with that of the CMJ and the participants  
195 were informed to hold for 3 seconds the squat position previous jumping.

196 All the participants performed the jumps barefoot. Two observers (one on either side of  
197 the subject) provided feedback about the starting height of 90° (knee flexion) squat during  
198 SJ performance. If one of the observers disagreed with the 90° (knee flexion) squat, the  
199 trial was invalidated and was repeated.

200

201 One week after the first testing session (Tuesday), participants were divided into a control  
202 group (CG) (14 females and 12 males) and 2 different experimental groups,  
203 corresponding to PT (15 females and 11 males) and CTr (12 females and 17 males). All  
204 the groups performed their usual training routine with regard to classes and rehearsal  
205 practice (34.2±4 hours). The experimental groups (PT and CTr) added 2 sessions (1 hour  
206 per session) of supplementary training per week, for 9 weeks (Tuesdays and Fridays). It  
207 was decided to have at least 48 hours between complementary training sessions for  
208 fatigue management, since these sessions were added in the early stage of the season. All  
209 the sessions were supervised by certified strength and conditioning (S&C) coaches. The  
210 training programs implemented for the PT and CTr groups can be seen in Table 1.  
211 Recovery periods of 2 and 5 minutes were set for intra and inter-set exercises,  
212 respectively. Due to the height and jumping differences between female and male  
213 dancers, the box heights were set at 40cm (F-40) and 50cm (M50) respectively.  
214 Participants were asked to meet the week after the end of the training intervention for the  
215 post-test measurements, following the same procedures used in the pre-test week.

216

217

**Table 1 here**

## 218 **Statistical Analysis**

219 All statistical analyses were conducted using the Statistical Package for Social Sciences  
220 (IBM SPSS version 27.0; SPSS, Chicago, IL, USA) and presented as means ± SD.  
221 Normal distribution for the study variables was observed with the Shapiro-Wilk Test.  
222 Two (one per each sex) two-way analysis of variance (ANOVA) (3group x 2time) with  
223 Bonferroni adjustment and level of significance set at  $p \leq 0.05$  was used for inter- and  
224 intra-group comparisons within sex. The magnitudes of change, within and between  
225 group comparisons, were calculated using Cohen's effect size (ES)<sup>28</sup>. The criterion for  
226 interpreting these magnitudes was  $< 0.2$  = trivial,  $0.2 - 0.6$  = small,  $0.6 - 1.2$  = moderate,  
227  $1.2 - 2$  = large and  $> 2.0$  = very large<sup>28</sup>. Within-session reliability for the pre-test session  
228 for CMJ, SJ and *sauté* was assessed using the intraclass correlation coefficient (ICC) and  
229 the coefficient of variation (CV), with the corresponding 95% confidence interval (CI).  
230 Acceptable reliability was determined as a  $CV < 10\%$  and  $ICC > 0.70$ <sup>29</sup>.

231

232 The associations between the magnitude of improvement in CMJ ( $\Delta\%$ -CMJ) and SJ ( $\Delta\%$ -  
233 SJ) height with the magnitude of improvement in *sauté* height ( $\Delta\%$ - *Sauté*) were analyzed  
234 using a Pearson correlation (r) and the coefficient of determination ( $R^2$ ), the level of  
235 significance was set at  $p \leq 0.05$ . The chosen criteria to interpret the magnitude of r were:  
236  $\leq 0.1$  = trivial,  $> 0.1 - 0.3$  = small,  $> 0.3 - 0.5$  = moderate,  $> 0.5 - 0.7$  = large,  $> 0.7 - 0.9$   
237 = very large,  $> 0.9 - 1.0$  = almost perfect<sup>28</sup>. The magnitudes of improvement for the  
238 variables previously mentioned were calculated using the formula:



239

240 
$$\left( \frac{\text{Post} - \text{Test Value}}{\text{Pre} - \text{Test Value}} - 1 \right) * 100.$$

241

242

243 **RESULTS**

244

245 Acceptable reliability **between the 3 trials for each jump**<sup>29</sup> was observed in the pre-test.  
 246 CMJ height (Female Pre-ICC=0.95, CV=7.4%, C.I.=30.7-31.4 **cm**; Male Pre-ICC=0.92,  
 247 CV=5.3%, C.I.=49.6-51.7 **cm**), SJ height (Female Pre-ICC=0.91, CV=9%, C.I.=26.5-29  
 248 **cm**; Male Pre-ICC=0.91, CV=6.7%, C.I.= 47.2-47.9 **cm**) and *sauté* height (Female Pre-  
 249 ICC=0.92, CV=9.1%, C.I.= 29.1-30.5 **cm**; Male Pre-ICC=0.94, CV=6.9%, C.I.= 48.8-  
 250 49.5 **cm**). The data for the CMJ, SJ and *sauté* performance in the pre and post comparisons  
 251 after 9 weeks of intervention for female and male dancers are provided in Table 2.

252

253

253 **Table 2 here**

254

255 The two-way ANOVA (3group x 2time) for the pre-test, reported no significant  
 256 differences between groups, for CMJ (Female F=0.08, p=0.92 and Male F=0.38, p=0.69),  
 257 SJ (Female F=1.09, p=0.35 and Male F=1.22, p=0.31) and *sauté* (Female F=0.47, p=0.63  
 258 and Male F=0.01, p=0.99) height in female and male dancers. The post-test analysis  
 259 showed significant differences for CMJ (Female F=18.4, p≤0.001, Male F=11.2, p≤0.01),  
 260 SJ (Female F=34.2, p≤0.001, Male F=3.54, p=0.04) and *sauté* (Female F=14.9, p≤0.001,  
 261 Male F=5.16, p=0.01) in both experimental groups (PT and CTr) against the CG for  
 262 female and male participants, as the post-hoc revealed.

263

264 Associations between the magnitude of improvement in CMJ height ( $\Delta$ -CMJ) and SJ  
 265 ( $\Delta$ %-SJ) height compared with the magnitude of improvement in *sauté* height ( $\Delta$ %-  
 266 *Sauté*) for each group can be observed in figure 1 and table 3.

267

268

268 **Figure 1 here**

269

270

270 **Table 3 here**

271

272 **DISCUSSION**

273

274 The aim of this study was (i) to compare the effects of two common methods of  
 275 supplementary training (PT, CTr) on CMJ, SJ and *sauté* height; and (ii) to observe  
 276 whether changes in CMJ and SJ were associated with changes on *sauté* after 9 weeks of  
 277 physical supplementary training in female and male dancers. Both methods significantly  
 278 improved CMJ, SJ and *sauté* in male and female dancers. In addition to this, our results  
 279 suggest that the improvement in CMJ and SJ has a positive transference to specific dance  
 280 skills based on jumping ability as measured during a *sauté*.

281

282 To our knowledge, this is the first study to observe the effect of supplementary training  
 283 to improve vertical jumping ability (CMJ and SJ), and its effect on a specific dance skill  
 284 such as *sauté*. Our findings suggest that irrespective of the sex of ballet dancers, PT and  
 285 CTr supplementary training are beneficial ways to improve jumping ability<sup>2,13</sup>. These  
 286 results align with those of Brown and colleagues for female dancers<sup>2</sup>. However, their  
 287 findings reported better results for CMJ using PT in comparison to traditional weight

288 training, while in our study the CTr showed a higher ES than PT. The reason for this may  
289 be the different strength training programs that were applied in each study. The exercises  
290 included by Brown and colleagues in their strength training group did not include loaded  
291 jumps. This exercise has reported large improvements in vertical jump, besides being a  
292 relevant training stimulus for the optimization of the stretch-shorten cycle<sup>30</sup>. Therefore,  
293 the loaded CMJ and SJ (with one or both limbs) included in our study, based on a  
294 percentage of the participants' BM, may have caused a specific adaptation in jump height  
295 performance.

296

297 Our results show that male dancers jump higher than females (~20cm either in CMJ, SJ  
298 and *sauté*), as has been previously suggested in other studies<sup>31</sup>. The CMJ showed the  
299 highest value for jump height, followed by *sauté*, as previous investigators have reported  
300<sup>22,23</sup>. The lack of involvement of the stretch-shortening cycle during the SJ provides an  
301 explanation for the lower values for this type of jump<sup>32</sup>. Interestingly, the CG for female  
302 dancers shows a significant decrease of ~1.3cm in CMJ (E.S.=0.52 Small) and ~1.2cm in  
303 SJ (E.S.=0.46 Small). Similar results can be seen in the dance science literature<sup>13,14,16</sup>,  
304 and may be related to the insufficient number of supplementary training sessions provided  
305 for dancers<sup>2,6-9</sup>, especially female participants<sup>31</sup>. Similarly, an analysis of the force-  
306 velocity profile during jumping, conducted by Escobar and colleagues, found that female  
307 dancers have a force deficit. This analysis of the mechanical variables during jumping  
308 suggests that dance training develops an insufficient amount of force to reach the optimal  
309 performance in CMJ<sup>17</sup>. On the other hand, our results also support the findings reported  
310 by Brown and colleagues, reporting plyometric training as an effective methodology to  
311 improve vertical jump height in female collegiate dancers<sup>2</sup>. The involvement of the lower  
312 limbs in quick eccentric and concentric actions enhance the capacity to use the stored  
313 energy in the muscle tendons<sup>33</sup>. All the above supports the suggestion that professional  
314 ballet dancers must implement supplementary training programs to develop force  
315 capabilities and improve vertical jump ability<sup>2,6-9,17</sup>. In contrast with the female dancers,  
316 the CG for male participants does not show a significant decrease in CMJ, SJ and *sauté*.  
317 This may be explained by the different choreographic demands and training programs for  
318 female and male dancers, as has been previously reported by some authors<sup>7</sup>.

319

320 One of the novel aspects of this investigation was observing the effect of improving CMJ  
321 and SJ height on *sauté*. The significant change found in *sauté* height in male and female  
322 participants in both experimental groups (see table 2), suggests that supplementary  
323 training has a positive transference to this specific dance skill. Moreover, the  
324 improvements in CMJ and SJ are strongly associated with improvements in *sauté* jump  
325 height. Table 3 shows a very large association between the improvement in CMJ and SJ  
326 and improvement in *sauté*. Interestingly, the r values for CMJ compared with the *sauté*  
327 are higher than those for SJ for each group regardless of the participants' sex, especially  
328 for the CTr group. This may be related to the adaptations caused by heavy strength training.  
329 The findings of Cormie and colleagues suggest that maximal strength plays an important  
330 role in the production of maximal power and the enhancement of athletic performance  
331 (i.e., jumping)<sup>30</sup>. Therefore, the combination of exercises with loads using 80% of 1RM,  
332 as well as loaded jumps using a percentage of the participants' BM, may have caused  
333 more optimal adaptations in both specific and non-specific dance jumps (*sauté* and CMJ,  
334 respectively).

335

336 Despite the kinetic and kinematic differences between CMJ, SJ and *sauté*<sup>22</sup>, our study  
337 supports the use of supplementary training based on plyometrics and strength training to  
338 improve *sauté* performance. In our study, the CTr group for male and female dancers  
339 showed a higher ES than the PT group in the pre- vs post-test comparison. Although the  
340 turnout position is not the conventional resistance training technique, used for exercises  
341 with loads such as back squats or loaded jumps, this study confirms the positive effect of  
342 lifting weights in a standard position on *sauté*. Moreover, our findings suggest that non-  
343 ballet specific testing (CMJ and SJ height) are appropriate for ballet dancers if only jump  
344 height is of interest.

345

346 One strength of our study lies in including professional ballet dancers. This is especially  
347 relevant regarding the male dancers, due to the limited number of studies observing this  
348 population<sup>19</sup>. Also, this study suggests that as little as two sessions (1 hour each) of  
349 supplementary training such as PT or CTr provides a significant improvement on *sauté*.  
350 However, our investigation also has limitations that must be considered. The transference  
351 of improved CMJ and SJ to a specific jump action such as *sauté* was observed from the  
352 perspective of performance. However, it may be interesting to observe the impact of this  
353 improvement on subjective or qualitative dance performance<sup>1,2,13,34</sup>. Although the  
354 validity and reliability of the contact mat used in our study have been reported<sup>27</sup>, the use  
355 of a force platform would be more optimal, due to being considered the gold standard to  
356 measure jump height. In addition to the above, it is important to point out that progressive  
357 overload in volume or intensity was not applied to the experimental groups, even so, large  
358 changes were observed in the height of the CMJ. The prescription of exercises based on  
359 1RM were not adjusted within the 9 weeks of intervention, and the groups were self-  
360 selected instead of randomized.

361

## 362 PRACTICAL APPLICATIONS

363

364 The findings of this investigation may be relevant for classical dancers and S&C coaches,  
365 due to the large number of vertical jumps that are part of ballet choreographies<sup>5,34</sup>. This  
366 study supports the use of combined and plyometrics training programs as effective  
367 methods for improving specific (*sauté*) and non-specific dance jumps (CMJ and SJ).

368

## 369 Conclusion

370

371 Plyometric and combined training programs are effective ways to improve jumping  
372 ability in female and male ballet dancers. The improvement in CMJ and SJ has a positive  
373 transference to an essential dance skill such as *sauté*.

374

## 375 ACKNOWLEDGMENTS

376

377 The authors thank all the dancers, teachers, and S&C coaches who were part of this  
378 investigation.

379

380 This study was made possible thanks to the removed for peer-reviewed

381

382

383

384

385

386

## REFERENCES

387

388

1. Angioi M, Metsios GS, Twitchett E, Koutedakis Y, Wyon M. Association between selected physical fitness parameters and esthetic competence in contemporary dancers. *J Dance Med Sci.* 2009;13(4):115-123.

389

2. Brown AC, Wells TJ, Schade ML, Smith DL, Fehling PC. Effects of Plyometric Training Versus Traditional Weight Training on Strength, Power, and Aesthetic Jumping Ability in Female Collegiate Dancers. *J Danc Med Sci.* 2007.

391

3. Rafferty S. Considerations for integrating fitness into dance training. *J Dance Med Sci.* 2010;14(2):45-49.

392

4. Hewett TE, Stroupe AL, Nance TA, Noyes FR. Plyometric training in female athletes: Decreased impact forces and increased hamstring torques. *Am J Sports Med.* 1996. doi:10.1177/036354659602400611

393

5. Wyon MA, Twitchett E, Angioi M, Clarke F, Metsios G, Koutedakis Y. Time Motion and Video Analysis of Classical Ballet and Contemporary Dance Performance. *Int J Sports Med.* 2011;32(11):851-855. doi:10.1055/s-0031-1279718

394

6. Wyon MA, Deighan MA, Nevill AM, et al. The cardiorespiratory, anthropometric, and performance characteristics of an international/national touring ballet company. *J Strength Cond Res.* 2007. doi:10.1519/R-19405.1

395

7. Wyon M, Allen N, Angioi M, Nevill A, Twitchett E. Anthropometric Factors Affecting Vertical Jump Height in Ballet Dancers. *J Danc Med Sci.* 2006;10:106-110.

396

8. Twitchett EA, Angioi M, Koutedakis Y, Wyon M. Do increases in selected fitness parameters affect the aesthetic aspects of classical ballet performance? *Med Probl Perform Art.* 2011.

397

9. Stalder MA, Noble BJ, Wilkinson JG. The effects of supplemental weight training for ballet dancers. *J Appl Sport Sci Res.* 1990.

398

10. Farmer C, Brouner J. Perceptions of Strength Training in Dance. *J Danc Med Sci.* 2021;25(3). doi:10.12678/1089-313X.091521a

399

11. Shaw JW, Mattiussi AM, Brown DD, Springham M, Pedlar CR, Tallent J. The activity demands and physiological responses observed in professional ballet: A systematic review. *J Sport Exerc Sci.* 2021;5(4). doi:10.36905/jses.2021.04.04

400

12. Shaw JW, Mattiussi AM, Brown DD, et al. *Dance Exposure, Individual Characteristics, and Injury Risk over Five Seasons in a Professional Ballet Company Short Title: Injury Risk Factors in Professional Ballet.*

401

13. Angioi PhD M, Metsios PhD G, Twitchett PhD EA, Koutedakis PhD Y, Wyon PhD M. Effects of Supplemental Training on Fitness and Aesthetic Competence Parameters in Contemporary Dance: A Randomised Controlled Trial. *Med Probl Perform Art.* 2012.

402

14. Wyon M, Guinan D, Hawkey A. Whole-body vibration training increases vertical jump height in a dance population. *J Strength Cond Res.* 2010. doi:10.1519/JSC.0b013e3181c7c640

403

15. Annino G, Padua E, Castagna C, et al. Effect of whole body vibration training on lower limb performance in selected high-level ballet students. *J Strength Cond Res.* 2007. doi:10.1519/R-18595.1

404

16. Marshall LC, Wyon MA. The effect of whole-body vibration on jump height and active range of movement in female dancers. *J Strength Cond Res.* 2012. doi:10.1519/JSC.0b013e31822a5ce8

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

- 435 17. Escobar Álvarez JA, Fuentes García JP, Da Conceição FA, Jiménez-Reyes P.  
436 Individualized Training Based on Force–Velocity Profiling During Jumping in  
437 Ballet Dancers. *Int J Sports Physiol Perform*. 2019. doi:10.1123/ijsp.2019-0492
- 438 18. Girard J, Koenig K, Village D. The effect of strength and plyometric training on  
439 functional dance performance in elite ballet and modern dancers. *Phys Ther Rev*.  
440 2015. doi:10.1179/1743288X15Y.0000000017
- 441 19. Mattiussi A, Shaw JW, Brown DD, et al. Jumping in Ballet: A Systematic  
442 Review of Kinetic and Kinematic Parameters. *Med Probl Perform Art*.  
443 2021;36(2). doi:10.21091/mppa.2021.2011
- 444 20. Karim AY, Ashmore C, Burriss S, Roddey T, Ortiz A, Olson S. Assessment of  
445 Countermovement Jump Performance in First Position (Sauté) in Healthy Adults.  
446 *Sports Health*. 2019. doi:10.1177/1941738118824569
- 447 21. Grant G. *Technical Manual and Dictionary of Classic Ballet*.; 2008.
- 448 22. Imura A, Iino Y. Comparison of lower limb kinetics during vertical jumps in  
449 turnout and neutral foot positions by classical ballet dancers. *Sport Biomech*.  
450 2017;16(1). doi:10.1080/14763141.2016.1205122
- 451 23. Ravn S, Voigt M, Simonsen EB, Alkjær T, Bojsen-Møller F, Klausen K. Choice  
452 of jumping strategy in two standard jumps, squat and countermovement jump -  
453 Effect of training background or inherited preference? *Scand J Med Sci Sport*.  
454 1999;9(4):201-208. doi:10.1111/j.1600-0838.1999.tb00234.x
- 455 24. Blanco P, Nimphius S, Seitz LB, Spiteri T, Haff GG. Countermovement Jump  
456 and Drop Jump Performances Are Related to Grand Jeté Leap Performance in  
457 Dancers With Different Skill Levels. *J Strength Cond Res*. 2019.  
458 doi:10.1519/jsc.0000000000003315
- 459 25. Lloyd RS, Radnor JM, De Ste Croix MBA, Cronin JB, Oliver JL. Changes in  
460 Sprint and Jump Performances After Traditional, Plyometric, and Combined  
461 Resistance Training in Male Youth Pre- and Post-Peak Height Velocity. *J*  
462 *Strength Cond Res*. 2016;30(5):1239-1247. doi:10.1519/JSC.0000000000001216
- 463 26. Rice PE, Nishikawa K, Nimphius S. Isolated Joint Block Progression Training  
464 Improves Leaping Performance in Dancers. *Front Sport Act Living*. 2021;3.  
465 doi:10.3389/fspor.2021.779824
- 466 27. Pueo B, Penichet-Tomas A, Jimenez-Olmedo J. Reliability and validity of the  
467 Chronojump open-source jump mat system. *Biol Sport*. 2020;37(3).  
468 doi:10.5114/biolspor.2020.95636
- 469 28. Cohen J. *Statistical Power Analysis of the Behavioral Sciences*.; 1988.  
470 doi:10.1234/12345678
- 471 29. Hopkins WG, Marshall SW, Batterham AM, Hanin J. *Progressive Statistics for*  
472 *Studies in Sports Medicine and Exercise Science*. Vol 41.; 2009:3-12.  
473 doi:10.1249/MSS.0b013e31818cb278
- 474 30. Cormie P, McGuigan MR, Newton RU. Adaptations in athletic performance after  
475 ballistic power versus strength training. *Med Sci Sports Exerc*. 2010.  
476 doi:10.1249/MSS.0b013e3181d2013a
- 477 31. Wyon MMA, Allen N, Angioi MM, Nevill A, Twitchett E. Anthropometric  
478 Factors Affecting Vertical Jump Height in Ballet Dancers. *J Danc Med Sci*.  
479 2006;10(3-4):106-110.
- 480 32. KOMI P V. Physiological and Biomechanical Correlates of Muscle Function:  
481 Effects of Muscle Structure and Stretch—Shortening Cycle on Force and Speed.  
482 *Exerc Sport Sci Rev*. 1984;12(1). [https://journals.lww.com/acsm-](https://journals.lww.com/acsm-essr/Fulltext/1984/01000/Physiological_and_Biomechanical_Correlates_of.6.aspx)  
483 [essr/Fulltext/1984/01000/Physiological\\_and\\_Biomechanical\\_Correlates\\_of.6.aspx](https://journals.lww.com/acsm-essr/Fulltext/1984/01000/Physiological_and_Biomechanical_Correlates_of.6.aspx)  
484 x.



- 485 33. Baechle T, Earle R, Wathen D. Essentials of Strength and Conditioning. *Hum*  
486 *Kinet Publ.* 2000. doi:10.1080/14763140208522788
- 487 34. Dowse RA, McGuigan MR, Harrison C. Effects of a Resistance Training  
488 Intervention on Strength, Power, and Performance in Adolescent Dancers. *J*  
489 *Strength Cond Res.* 2017. doi:10.1519/JSC.0000000000002288  
490

For Peer Review

<b>PT</b>	<b>1st Session (Tuesdays)</b>		<b>2nd Session (Fridays)</b>	
	<b>Exercises</b>	<b>Sets/repetitions</b>	<b>Exercises</b>	<b>Sets/repetitions</b>
	Horizontal SJ	3x10	Single leg SJ	3x10 each leg
	Single leg CMJ	3x10 each leg	Sidestep up + Drop SJ	3x10 each leg (F40-M50cm)
	Box SJ	3x10 (F40-M50cm)	Box CMJ	3x10 (F40-M50cm)
	Step up + Drop CMJ	3x10 each leg (F40-M50cm)	Drop CMJ	3x10 (F40-M50cm)
<b>CTr</b>	<b>1st Session (Mondays)</b>		<b>2nd Session (Thursdays)</b>	
	<b>Exercises</b>	<b>Sets/repetitions/Training load</b>	<b>Exercises</b>	<b>Sets/repetitions/Training load</b>
	Leg press	3x8 (80%1RM)	Back-Squat	3x8 (80%1RM)
	Deadlift	3x8 (80%1RM)	CMJ	3x8 (80% BM)
	Leg Curl	3x8 (80%1RM)	Leg extension	3x8 (80%1RM)
	SJ	3x6 (80%BM)	Single leg CMJ	3x6 (10% BM)

**Table 1:** Training programs followed by both intervention groups; Plyometric training (PT) and Combined training (CTr). Box height in females set at 40cm (F-40) and 50cm in males (M50).

	Variables	Pre	Post	P	95%C.I.	E.S.
CG ♀	CMJ (cm)	30.9±2.6	29.6±2.3	<b>0.04*</b>	-1.28±0.22	0.52 (S)
	S.J. (cm)	27.2±2.7	26±2.5	<b>0.05*</b>	-1.21±0.28	0.46 (S)
	<i>Sauté</i> (cm)	29.5±2.6	28 ±2.4	0.1	-1.35±0.15	--
PT ♀	CMJ (cm)	31.2±2	33.4±2.5	<b>0.03*</b>	0.21±1.72	0.97 (M)
	S.J. (cm)	28±2.3	30.3±2.6	<b>0.03*</b>	0.16±1.67	0.91 (M)
	<i>Sauté</i> (cm)	30.1±2.9	31.6±2.3	<b>0.03*</b>	0.12±1.34	0.61 (M)
CTr ♀	CMJ (cm)	31±2.3	34.7±2.9	<b>0.01*</b>	0.52±2.3	1.41 (L)
	S.J. (cm)	28.1±2.4	31.2±2.6	<b>0.01*</b>	0.36±2.11	1.23 (L)
	<i>Sauté</i> (cm)	29.8±2.7	31.9±2.5	<b>0.02*</b>	0.02±1.63	0.80 (M)
CG ♂	CMJ (cm)	50.2±3.1	49.9±3.6	0.7	-0.90±0.71	--
	S.J. (cm)	47.4±3.4	47.3±3.5	0.8	-0.80±0.77	--
	<i>Sauté</i> (cm)	49.1±3.3	48.4±3.6	0.5	-1±0.60	--
PT ♂	CMJ (cm)	51±2.7	53.6±3.4	<b>0.03*</b>	0.02±1.70	0.84 (M)
	S.J. (cm)	47.7±3.3	50.1±3.5	<b>0.04*</b>	0.15±1.56	0.70 (M)
	<i>Sauté</i> (cm)	49±3.7	52.1±3.4	<b>0.02*</b>	0.00±1.74	0.87 (M)
CTr ♂	CMJ (cm)	50.8±2.2	54.3±3.1	<b>0.01*</b>	0.56±2.04	1.30 (L)
	S.J. (cm)	47.6±3	50.9±3.4	<b>0.02*</b>	0.31±1.74	1.02 (M)
	<i>Sauté</i> (cm)	49.3±3.3	52.3±3.2	<b>0.02*</b>	0.21±1.63	0.92 (M)

**Table 2:** Mean ± S.D., p-values ( $p \leq 0.05$ ), 95% confidence intervals (95% C.I.) and effect size (Small-S, Medium-M, Large-L and very large-VL) for jumping performance in female (♀) and male (♂) dancers in control group (C.G.), plyometric group (PT) and combined training group (CTr). The presented variables are defined as countermovement jump (CMJ), squat jump (S.J.) and *sauté* in first position (*sauté*). Statistically significant differences are denoted in bold an \*.



Group		$\Delta$ - <i>Sauté</i>	R <sup>2</sup>
PT ♀	$\Delta$ -CMJ	0.82 (VL)*	0.67*
	$\Delta$ -SJ	0.79 (VL)*	0.62*
CTr ♀	$\Delta$ -CMJ	0.88 (VL)*	0.78*
	$\Delta$ -SJ	0.78 (VL)*	0.61*
PT ♂	$\Delta$ -CMJ	0.85 (VL)*	0.72*
	$\Delta$ -SJ	0.77 (VL)*	0.59*
CTr ♂	$\Delta$ -CMJ	0.88 (VL)*	0.78*
	$\Delta$ -SJ	0.83 (VL)*	0.68*

**Table 3:** Pearson's correlation ( $r \leq 0.1$  = trivial,  $> 0.1 - 0.3$  = small,  $> 0.3 - 0.5$  = moderate,  $> 0.5 - 0.7$  = large,  $> 0.7 - 0.9$  = very large,  $> 0.9 - 1.0$  = almost perfect) and coefficient of determination (R<sup>2</sup>), between the magnitude of improvement for CMJ height ( $\Delta\%$ -CMJ) and SJ ( $\Delta\%$ -SJ) height with the magnitude of improvement for *sauté* ( $\Delta\%$ -*Sauté*) in female (♀) and male (♂) dancers for the plyometric training group (PT) and combined training group (CTr). Associations statistically significant ( $p \leq 0.05$ ) denoted in bold and \*.




Figure 1: Coefficient of determination between the magnitude of improvement for CMJ height ( $\Delta\%$ -CMJ) and SJ ( $\Delta\%$ -SJ) height with the magnitude of improvement for sauté ( $\Delta\%$ - Sauté) in female (♀) and male (♂) dancers for the plyometric training group (PT) and combined training group (CTr).

605x697mm (38 x 38 DPI)