



UWS Academic Portal

Infection control implications of the laundering of ambulance staff uniforms and reusable mops

Mackay, W.G.; Whitehead, S.; Purdue, N.; Smith, M.R.; Redhead, N.; Williams, C.; Wilson, S.

Published in:
Journal of Hospital Infection

DOI:
[10.1016/j.jhin.2017.02.017](https://doi.org/10.1016/j.jhin.2017.02.017)

Published: 01/05/2017

Document Version
Peer reviewed version

[Link to publication on the UWS Academic Portal](#)

Citation for published version (APA):
Mackay, W. G., Whitehead, S., Purdue, N., Smith, M. R., Redhead, N., Williams, C., & Wilson, S. (2017). Infection control implications of the laundering of ambulance staff uniforms and reusable mops. *Journal of Hospital Infection*, 96(1), 59-62. <https://doi.org/10.1016/j.jhin.2017.02.017>

General rights

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact pure@uws.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Infection control implications of the laundering of ambulance staff uniforms and reusable mops

Running title: laundering of ambulance staff uniforms and reusable mops

Mackay WG ^{1*}, Whitehead S ², Purdue N ³, Smith M ¹, Redhead N ⁴, Williams C ¹, Wilson S ³.

1. Institute of Healthcare Policy and Practice, School of Health, Nursing and Midwifery, University of the West of Scotland and University Hospital Crosshouse, UK
2. Department of Microbiology, Monklands District General Hospital, Airdrie, UK
3. Scottish Ambulance Service, Edinburgh, UK
4. National Services Scotland, Glasgow, UK

Corresponding author:

Email: w.mackay@uws.ac.uk, Tel: 07828669984

Summary

There is a lack of published studies on laundering in ambulance services. We performed bacterial culture on soiled and unsoiled uniforms and reusable mop heads artificially contaminated with *Escherichia coli*, *Staphylococcus aureus* and *Clostridium difficile* spores. Current laundering processes used for routine cleans in the ambulances appears, from our simulations, to be effective at reducing vegetative pathogenic bacteria to undetectable levels, $<3.398 \log_{10}$ CFU (*S. aureus* and *E. coli*). Reduced levels of *Clostridium difficile* were still detected after laundering but the risk this poses for infection is unknown, as background levels of these spores in the environment are unknown.

Keywords: Cleaning, disinfection, decontamination, ambulance

Introduction

Most patients arrive at hospital in an ambulance, yet little has been published on infection control in the pre-hospital setting. During 2014/2015 in Scotland there were 740,631 emergency and urgent incidents involving ambulances.[1] During 2013/2014 2.87 million calls to the ambulance service in England resulted in a response from an emergency vehicle, and in Victoria (Australia) there were 823,278 ambulance callouts during 2012/2013.[2, 3] Patients and healthcare personnel may be exposed to pathogens while in ambulances and this may be a route for the spread of pathogens between healthcare facilities, yet little information is currently available in the literature.

The Scottish Ambulance Service have laundering facilities on site that are used by the majority of staff to wash uniforms and also central facilities to wash reusable mop heads used to clean ambulances. The uniforms of healthcare staff and the environments they work in can become contaminated with potential human pathogens.[4] However, there are few published, peer-reviewed scientific studies on the infection control risks of ambulance staff uniforms or the microbiological quality of their laundering. The results of a cross-sectional study on the microbiological contamination of ambulance staff uniforms has provided direct evidence that uniforms can become contaminated with common potential human pathogens.[5] Evidence also exists for the contamination of ambulances with MRSA.[6]

We examined the effectiveness of ambulance station laundering in reducing bacterial contamination of ambulance staff uniforms, and the effectiveness of laundering of reusable mop heads used to clean the ambulances.

Methods

Bacterial strains were supplied by the National Collection of Type Cultures (London, UK). *Escherichia coli* (*E. coli*) (NCTC 9001), *Staphylococcus aureus* (*S. aureus*) (NCTC 8178) and *Clostridium difficile* (NCTC 11209) spores were used in this study. *E. coli* and *S. aureus* were cultivated overnight in broth culture. *C. difficile* was supplied as a spore stock by Blutest Laboratories Ltd, UK and prepared after the method described in Fraise et al. [7] The Scottish Ambulance Service provided a clean, unused, plastic-wrapped ambulance staff uniform and reusable mop heads. Ninety millimetre circular uniform swatches and reusable mop heads were sterilised by autoclaving at 121°C for 15 minutes. Swatches and mop strands (4 cm) were placed within sterile plastic 90 mm Petri dishes (Fisher Scientific, Loughborough, UK) in a class 1 biosafety cabinet (MDH Ltd, Andover, UK). The swatches and mop strands, in triplicate, were artificially contaminated with bacteria (4.685 – 7.304 log₁₀ CFU for *E. coli*, 4.749 - 6.816 log₁₀ CFU for *S. aureus*, and 6.426 - 7.495 log₁₀ CFU for *C. difficile* spores) with soiling (defibrinated horse blood) or without soiling (phosphate buffered saline) (100µL) covered and allowed to air dry at room temperature for 90 minutes. Unwashed controls were also analysed to determine the inoculum CFU, and a control without bacterial challenge washed to seek evidence for transfer of bacteria within the wash. Swatches were identified by punching holes near the edge prior to sterilisation – different numbers corresponding to different challenges. The swatches were washed together (Miele PW6055 machine / Cleanline biological washing powder) at 30°C (38 min cycle) or 60°C (49 min cycle) and reusable mop heads at 30°C (38 min cycle) or 95°C (62 min cycle).

All Swatches and mop strands were processed at the same time as follows: placed separately in 50mL of sterile buffered peptone water (Oxoid, Basingstoke, UK), and processed in a stomacher (Stomacher 400, Seward, Worthing, UK) (4 min) using the 'high' setting. The liquid was removed and analysed using the Miles and Misra plate counting method.[8] *E. coli* and *S. aureus* were assayed on CLED plates incubated at 37°C for 24 hours. *C. difficile* spores were assayed on *Clostridium difficile* agar (CDA) plates (*Clostridium difficile* agar base, D-cycloserine [250mg/L], Cefoxitin [8mg/L]) (Oxoid, Basingstoke, UK) and 7% defibrinated horse blood (Fisher Scientific, Loughborough, UK). CDA plates were incubated at 37°C for 72 hours

under anaerobic conditions in gas jars (Becton, Dickinson and Company, Oxford, UK).

We compared the log₁₀ CFU counts reported for the washed test materials to the unwashed controls (log₁₀ reduction). Where the CFU counts for the washed test materials were below the lower limit of detection for the Miles and Misra assay the log₁₀ reduction was estimated based on the distance between the log₁₀ CFU value for the unwashed controls and the lower limit of detection of the assay (3.398 log₁₀ CFU), and reported as 'greater than' values.

Results

E. coli were not recovered from any of the unsoiled or soiled swatches. *S. aureus* were only recovered from swatches washed at 30°C. *C. difficile* spores were recovered from all swatches washed at 30°C and 60°C (Table 1). Neither *E. coli* or *S. aureus* were recovered from washed unchallenged swatches. However, *C. difficile* spores, washed at 30°C and 60°C, were recovered from unchallenged swatches at the lower limit of detection of the assay (3.398 log₁₀ CFU).

E. coli were recovered from all mop strands washed at 30°C but not at 95°C. *S. aureus* were recovered from two of three unsoiled mop strands and all soiled mop strands washed at 30°C but none at 95°C. *C. difficile* spores were recovered from all mop strands washed at 30°C and one of three unsoiled mop strands washed at 95°C (Table 2). Neither *E. coli*, *S. aureus* or *C. difficile* spores, washed at 30°C and 95°C, were recovered from unchallenged mop strands.

Discussion

We examined the reduction in bacterial counts following standard laundering practice for staff uniforms and the reduction in bacterial counts following standard laundering practice for reusable mop heads.

Higher temperatures were effective at reducing *E. coli* and *S. aureus* to undetectable levels on uniform swatches and mop strands. However, *C. difficile* spores were still detectable after washing at 60°C both on the deliberately contaminated (challenged) (Table 1) and unchallenged uniform swatches, and one mop strand washed at 95°C but not the rest (Table 2). These data suggest that a proportion of *C. difficile* spores survived the wash and remained within the washing machine or rinse water.

Evidence exists that healthcare workers' uniforms can become contaminated with bacteria during use, with a mixture of bacteria from the wearers own flora (mostly non-pathogenic in nature) and potential pathogens from handling patients.[9] Nurse uniforms in hospitals have been shown to harbour MRSA, *C. difficile* and VRE both before and after shifts.[4] The ability of laundering to kill *Enterococcus faecalis* has been shown to be temperature dependant with a greater than 4 log₁₀ reduction at 65°C and greater than 8 log₁₀ reduction at 85°C (20 minute contact time).[10] We found that the log₁₀ reduction was heavily influenced by the temperature of the wash for *S. aureus*, *E. coli* and *C. difficile* spores.

Little information is available on the potential survival of *C. difficile* spores during laundering. Our data show that *C. difficile* spores could potentially survive laundering at temperatures as high as 60°C and perhaps as high as 95°C. These results are important and have implications in any environment where patients and staff are at risk of coming into contact with *C. difficile*. We also observed that *C. difficile* spores were transferred to the sterile unchallenged materials when uniform swatches were washed at 30 and 60°C, which suggests that there may be a risk of other garments becoming contaminated if washed along with materials contaminated with *C. difficile* spores. However, we do not know whether the reduced levels of *C. difficile* spores that may theoretically remain after washing are likely to pose a cross contamination risk to patients. We are not aware of any solid evidence that ambulances become contaminated with *C. difficile*. There are relatively few studies looking at the transmission of *C. difficile* from clothes, and the assessment of the impact of hygiene practices such as laundry hygiene is difficult to assess in isolation from other factors.[11] As a result, the possible presence of small numbers of *C. difficile* spores on some materials post laundering is not considered to be a high risk for the spread of infection. These results support the current practice and processes in place for

laundering of both ambulance staff uniforms and mop heads in the ambulance setting; where there is no evidence of contamination.

Conclusions

The laundering of ambulance staff uniforms and mop heads used for routine cleans in the ambulances appears, from our simulations, to be effective at reducing vegetative pathogenic bacteria to undetectable levels at higher temperatures. Laundering also reduces the number of *C. difficile* spores but they are still detectable on ambulance staff uniform swatches washed at 60°C and possibly mop strands washed at 95°C. These results support the current practice and processes in place for laundering of both uniforms and mop heads in ambulance stations; where there is no evidence of contamination. Although the study demonstrated that laundering processes were also effective when samples were contaminated with blood; in line with NHS Scotland guidance, contaminated uniforms and mop heads continue to be disposed of.

Acknowledgements

The Scottish Ambulance Service funded this work.

We are particularly grateful to the ambulance stations and staff in Paisley, Greenock and Vale of Leven (UK) for the use of the facilities that allowed us to complete the study. We are grateful to Blutest for supplying the *C. difficile* spore stock.

References

- [1] Scottish Ambulance Service. Scottish Ambulance Service Annual Report 2014/2015. Edinburgh, UK: Scottish Ambulance Service 2015.
- [2] Workforce and Facilities Team. Ambulance Services, England 2013-14. Health and Social Care Information Centre 2014.
- [3] Ambulance Victoria. Ambulance Victoria 2012-2013 Annual Report. Doncaster, Victoria: Ambulance Victoria 2014.
- [4] Perry C, Marshall R, Jones E Bacterial contamination of uniforms. *Journal of Hospital Infection* 2001; **48**: 238-41.

- [5] Vikke H, Giebner M UniStatus - a cross-sectional study on the contamination of uniforms in the Danish ambulance service. *BMC Research Notes* 2015; **8**: 95.
- [6] Eibicht SJ, Vogel U Meticillin-resistant Staphylococcus aureus (MRSA) contamination of ambulance cars after short term transport of MRSA-colonised patients is restricted to the stretcher. *Journal of Hospital Infection* 2011; **78**: 221-5.
- [7] Fraise AP, Wilkinson MAC, Bradley CR *et al.* Development of a sporicidal test method for Clostridium difficile. *Journal of Hospital Infection* 2015; **89**: 2-15.
- [8] Miles AA, Misra SS, Irwin JO The estimation of the bactericidal power of the blood. *The Journal of Hygiene* 1938; **38**: 732-49.
- [9] Wilson JA, Loveday HP, Hoffman PN, Pratt RJ Uniform: an evidence review of the microbiological significance of uniforms and uniform policy in the prevention and control of healthcare-associated infections. Report to the Department of Health (England). *Journal of Hospital Infection* 2007; **66**: 301-7.
- [10] Orr KE, Holliday MG, Jones AL, Robson I, Perry JD Survival of enterococci during hospital laundry processing. *Journal of Hospital Infection* 2002; **50**: 133-9.
- [11] Bloomfield S, Exner M, Signorelli C, Nath K, Scott E. The infection risks associated with clothing and household linens in home and everyday life settings, and the role of laundry. International Scientific Forum on Home Hygiene 2013.